**Age-related Differences in Implicit Sequence Learning and Consolidation
across the Human Life Span:
Implications for the Functioning of the Fronto-Striatal Circuitry**

**PhD Thesis
Karolina JANACSEK**

**Institute of Psychology
University of Szeged**

Supervisor: Dezso NEMETH, PhD.

**Szeged
2012**

Papers the thesis is based on:

I. **Janacsek, K.***, & Nemeth, D. (2012). Predicting the future: from implicit learning to consolidation. *International Journal of Psychophysiology*, 83(2), 213-221. DOI: 10.1016/j.ijpsycho.2011.11.012. Impact factor: 2.378.

II. **Janacsek, K.***, Fiser J., & Nemeth, D. (in press). The best time to acquire new skills: Age-related differences in implicit sequence learning across life span. *Developmental Science*. DOI: 10.1111/j.1467-7687.2012.01150.x. Impact factor: 3.53.

III. Nemeth, D., **Janacsek, K.***, Polner, B., & Kovacs, Z. A. (in press). Boosting human learning by hypnosis. *Cerebral Cortex*. DOI: 10.1093/cercor/bhs068. Impact factor: 6.844.

IV. Nemeth, D., & **Janacsek, K.*** (2011). The dynamics of implicit skill consolidation in young and elderly adults. *Journal of Gerontology Series B: Psychological Sciences*, 66B(1), 15-22. DOI: 10.1093/geronb/gbq063. Impact factor: 1.963.

Cumulative impact factor of papers the thesis based on: 14.715

Cumulative impact factor of Karolina Janacsek's all papers: 38.25

Papers related to the thesis:

I. Hallgató, E., Györi-Dani, D., Pekár, J., **Janacsek, K.***, & Nemeth, D. (in press). Perceptual – Motor learning debate in skill acquisition: the role of consolidation. *Cortex*. DOI: 10.1016/j.cortex.2012.01.002. Impact factor: 7.251.

II. Nemeth, D., **Janacsek, K.***, Csifcsak, G., Szvoboda, G., Howard, J. H. Jr., & Howard, D. V. (2011). Interference between sentence processing and probabilistic implicit sequence learning. *PLoS ONE*, 6(3): e17577. DOI: 10.1371/journal.pone.0017577. Impact factor: 4.411.

III. Nemeth, D., **Janacsek, K.***, Balogh, V., Londe, Zs., Mingesz, R., Fazekas, M., Jambori, Sz., Dányi, I., & Vetró, Á. (2010). Learning in Autism: implicitly superb. *PLoS ONE*, 5(7), e11731. Impact factor: 4.411.

IV. Nemeth, D., **Janacsek, K.***, Londe Zs., Ullman, M. T., Howard D. V., & Howard, J. H. Jr. (2010). Sleep has no critical role in implicit motor sequence learning in young and old adults. *Experimental Brain Research*, 201(2), 351-358. DOI : 10.1007/s00221-009-2024-x. Impact factor: 2.296.

V. Nemeth, D., Hallgato, E., **Janacsek, K.***, Sandor, T., & Londe Zs. (2009). Perceptual and motor factors of implicit skill learning. *Neuroreport*, 20, 1654-1658. Impact factor: 1.805.

* - first author or equivalent (equal contribution)

Abstracts related to the thesis:

- I. **Janacsek, K., & Nemeth, D.** (2011). Well-established skills are resistant to disruption – evidence from a dual-task paradigm. *ICOM-5, International Conference on Memory*, 31 July – 06 August, York, UK.
- II. **Janacsek, K., & Nemeth, D.** (2011). The time-course of implicit skill consolidation in young and elderly adults. *III Dubrovnik Conference on Cognitive Science*, 12-15 May, Dubrovnik, Croatia.
- III. **Janacsek K., & Németh D.** (2011). Consolidation and age-related changes in implicit learning. *XXXth National Conference of Hungarian Psychological Association*. 25-27 May, Budapest, Hungary.
- IV. **Nemeth D. & Janacsek, K.** (2010). Age-Related Changes in Implicit Learning Across Human Life Span. *Psychonomic Society: The 51st Annual Meeting*, 18-21 November, St. Louis, USA.
- V. **Janacsek, K., Londe, Zs., Ullman, M.T., Howard, D. V., Howard, J. H. Jr. & Nemeth, D.** (2010). One ring does not rule them all - Sleep has no critical role in implicit motor sequence learning in young and old adults. *IBRO International Workshop 2010*, 21-23 January, Pécs, Hungary.
- VI. **Janacsek, K., Vízi, I. & Németh, D.** (2009). Implicit learning and consolidation in old age. *MAKOG XVII – XVIIth Conference of Hungarian Cognitive Science Association*. 7-9 May, Budapest, Hungary.
- VII. **Janacsek, K., Vízi, I. & Németh, D.** (2009). The effect of aging on implicit skill learning. *Congress of Hungarian Psychiatric Association*. 28-31 January, Debrecen, Hungary.

Table of contents

TABLE OF CONTENTS.....	4
SUMMARY	5
ÖSSZEFOGLALÁS	6
INTRODUCTION.....	10
MEASURES OF IMPLICIT SEQUENCE LEARNING.....	11
CONSOLIDATION OF SEQUENCE KNOWLEDGE.....	14
AGE-RELATED DIFFERENCES IN IMPLICIT LEARNING.....	15
BIOLOGICAL BACKGROUND OF IMPLICIT SEQUENCE LEARNING AND CONSOLIDATION	16
INTERACTION BETWEEN LEARNING/MEMORY SYSTEMS.....	18
AIMS OF THE STUDIES	19
EXPERIMENT I	20
METHODS.....	21
PARTICIPANTS	21
TASK AND PROCEDURE.....	22
STATISTICAL ANALYSIS.....	23
RESULTS.....	24
DISCUSSION OF EXPERIMENT I	26
EXPERIMENT II	31
METHODS.....	32
PARTICIPANTS	32
TASKS	33
DESIGN AND PROCEDURE	33
STATISTICAL ANALYSIS.....	34
RESULTS.....	34
DISCUSSION OF EXPERIMENT II	36
EXPERIMENT III.....	37
METHODS.....	39
PARTICIPANTS	39
PROCEDURE	39
TASK.....	40
STATISTICAL ANALYSIS.....	40
RESULTS.....	41
ONLINE LEARNING.....	41
OFFLINE SEQUENCE-SPECIFIC LEARNING	43
OFFLINE GENERAL SKILL LEARNING	43
DISCUSSION OF EXPERIMENT III.....	44
GENERAL DISCUSSION	46
CONCLUSIONS	48
ACKNOWLEDGEMENTS.....	49
REFERENCES.....	50
APPENDIX.....	62

Summary

Implicit sequence learning occurs when information is acquired from an environment of complex stimuli without conscious access either to what was learned or to the fact that learning occurred. In everyday life, this learning mechanism is crucial for adapting to the environment and for predicting events unconsciously. Despite the growing interest in implicit learning in the past decades, there has been relatively little research on life-long development of implicit sequence learning and on offline processing of implicitly learned information (i.e., consolidation). Here, we present three studies investigating these issues. In Experiment I, we investigated implicit sequence learning from 4 to 85 years of age and found a marked decrease in learning performance - measured by raw reaction time (RT) – around age of 12. This decrement can be explained by a competition between two fundamentally different forms of learning (model-free and model-based) suggesting that after adolescence frontal lobe-mediated model-based learning has larger effect on the expression of implicit sequence knowledge, while before adolescence basal ganglia-dependent model-free learning is more influencing. As a growing body of data has shown that frontal lobe-mediated processes are disrupted in hypnosis, we tested this assumption in Experiment II by comparing implicit sequence learning in hypnosis and in waking alert state. We found that hypnosis boosted sequence learning in young adults. In addition, this boosting effect was mediated by frontal lobe related executive functions. Finally, we investigated consolidation of implicit sequence knowledge in young and elderly adults after 12-, 24-hour or a 1-week delay period in order to determine age-related differences not only in online learning, but also in offline processing of the learned material (Experiment III). We found that consolidation is not a single process, rather there are multiple mechanisms (e.g., sequence-specific, general skill learning) which are differentially affected by aging and the course of time. Our results contribute not only to the better understanding of learning on a behavioral level, but also to understanding the age-related changes in brain plasticity in healthy participants across the human life span. In addition, these findings can help better understand neurodevelopmental (e.g., autism, dyslexia), neurodegenerative (e.g., Parkinson's Disease, Huntington's Disease) and age-related disorders where related brain structures are affected. Finally, our findings can lead to the development of more effective diagnostic tools, training methods and rehabilitation programs.

Összefoglalás

Az implicit tanulás alapvető a mindennapjainkban, ugyanis a környezetünkben fellelhető mintázatok, szabályszerűségek többségét implicit módon sajátítjuk el (Hawkins, George, & Niemasik, 2009; Janacsek & Nemeth, 2012). Erre a tanulási formára jellemző, hogy nem tudatos, nehezen verbalizálható, és viszonylag lassú az explicit tanuláshoz képest (Cleeremans, Destrebecqz, & Boyer, 1998; Reber, 1993). Agyi hátterét tekintve elsősorban a kéreg alatti struktúrák és azon belül is a striatum szerepe jelentős (pl. Doyon, Bellec et al., 2009; Hikosaka, Nakamura, Sakai, & Nakahara, 2002). Újabb kutatások azonban rávilágítottak arra, hogy több, egymással kooperatív, ugyanakkor néha kompetitív viszonyban álló folyamat is jelen van a tanulás során, melyeket a striatum frontális lebenyhez fűződő kapcsolatai alapoznak meg. Ennek következtében bizonyos helyzetekben az implicit tanulás javítható vagy gyengíthető a frontális lebenyhez kötődő funkciók manipulálásával (pl. Filoteo, Lauritzen, & Maddox, 2010; Galea, Albert, Ditye, & Miall, 2010).

Az implicit tanulás képezi az alapját számos mozgásos (pl. autóvezetés, sportolás), nyelvi-zenei (pl. olvasás, idegen nyelvek ismerete) és szociális (pl. társas interakciók, társas státusz) készségünknek (Kaufman et al., 2010; Lieberman, 2000; Nemeth et al., 2011; Romano Bergstrom, Howard, & Howard, 2011; Ullman, 2004), melyek meghatározóak a mindennapjainkban gyerekkortól egészen időskorig. Annak ellenére, hogy az utóbbi évtizedekben megnövekedett a tudományos érdeklődés az implicit tanulás iránt, viszonylag kevés kutatás foglalkozott az implicit tanulás egész életen át tartó fejlődésének kérdésével illetve azzal, hogy hogyan történik az implicit módon elsajátított ismeretek 'offline' feldolgozása, vagyis konszolidációja. Jelen disszertáció három kutatást mutat be, melyek ezeket a kérdéseket vizsgálják meg.

Az implicit tanulás életkori változásaival foglalkozó kutatások többnyire egyetértenek abban, hogy idős korban gyengül a tanulási képesség (pl. D. V. Howard et al., 2004; Nemeth, Janacsek, Londe et al., 2010), azonban a gyerekkori teljesítmény kapcsán vita bontakozott ki. Az egyik nézőpont szerint a tanulás hátterében álló agyi struktúrák már viszonylag korai életkorban érettek, így a tanulás már gyerekkorban hasonló szintű, mint felnőttkorban (pl. Meulemans, Van der Linden, & Perruchet, 1998; Vinter & Perruchet, 2000). A másik nézőpont képviselői szerint felnőttkorban a leghatékonyabb a tanulás (pl. J. Fletcher, Maybery, & Bennett, 2000; Maybery, Taylor, & O'Brien-Malone, 1995), és ez összefüggésbe hozható a fronto-striális hálózatok jelentős fejlődésével egészen felnőttkorig (pl. Thomas et al., 2004). Eddig azonban nem született olyan kutatás, amely ugyanazzal a módszerrel

vizsgálta volna egy széles életkori spektrumban az implicit tanulást. Ezért az 1. kísérlet célja az volt, hogy az implicit tanulás egész életen át tartó változásait térképezze fel. A vizsgálatban összesen 421 személy vett részt. Az életkor 4-től 85 éves korig terjedt. A személyeket 9 életkori csoportba soroltuk. Az implicit tanulás mérésére a kognitív idegtudományban egyre népszerűbb alternáló szeriális reakcióidő (ASRT) feladatot használtuk (J. H. Howard, Jr. & Howard, 1997), melyben a vizsgálati személyeknek a számítógép képernyőjén megjelenő ingerekre kell minél gyorsabban és pontosabban válaszolniuk a megfelelő gombok megnyomásával. Az ingerek megjelenésének sorrendjében van egy rejtett mintázat (pl. 2r3r1r4r, ahol a számok az ingerek megjelenési helyét mutatják, az 'r' pedig egy random hely a négy lehetséges pozíció közül). A személyeknek erről nincs explicit/deklaratív tudásuk; implicit módon tanulják meg a rejtett mintázatot - ezt nevezzük a továbbiakban szekvencia-specifikus tanulásnak. A feladat előnye, hogy ezt a szekvencia-specifikus tanulást el tudjuk különíteni az általános készségtanulástól, vagyis attól, hogy a vizsgálati személyek egyre nagyobb rutint szereznek a válaszadás során, azonban ez független az ingerek megjelenésének szekvenciájától (lásd pl. Song, Howard, & Howard, 2007). Az ASRT feladat ebben a vizsgálatban 20 blokkból állt.

Az 1. kísérlet eredményei alapján megállapítható, hogy a 4-től 12 éves korig terjedő életkori csoportok mutatták a legnagyobb szekvencia-specifikus tanulást (a reakcióidő adatok alapján), a későbbi életkori csoportokban pedig valamivel gyengébb tanulás jelentkezett. Ennek a 12 éves kor körüli váltásnak a hátterében két alapvetően különböző (modell-függő és modell-független) tanulási forma versengése állhat (lásd pl. Daw, Niv, & Dayan, 2005). Ez a feltételezés alapján serdülőkor előtt a tanulás inkább modell-független, ami azt jelenti, hogy a környezetben fellelhető szabályszerűségek észlelése befolyásolja elsősorban a tanulási teljesítményt. Ez a modell-független tanulási forma a bazális ganglionokhoz köthető (lásd pl. Daw et al., 2005). A későbbi életkorokban pedig a frontális lebeny működésén alapuló modell-függő tanulás válik dominánssá, vagyis ekkor már a tanulást nagyobb mértékben befolyásolják a környezetünkről kialakított belső modellek, és kevésbé támaszkodunk a környezet statisztikai együttjárásainak, szabályszerűségeink észlelésére. A modell-függő tanulási forma hatékony lehet a már megtanult készségek használata során, illetve a környezet újabb eseményeinek bejósolására, azonban az alapvetően új szabályszerűségek, készségek megtanulását akár hátráltathatja is. A modell-függő és modell-független tanulás hátterében álló feltételezett mechanizmusok és agyi struktúrák összhangban állnak a különböző emlékezeti és tanulási folyamatok kooperatív-kompetitív viszonyát hangsúlyozó kutatásokkal (pl. Filoteo et al., 2010; Galea et al., 2010).

Mivel egyre több bizonyíték van arra, hogy a hipnózis megzavarja a frontális lebenyhez kötődő funkciókat (Egner, Jamieson, & Gruzelier, 2005; Fingelkurts, Kallio, & Revonsuo, 2007; Oakley & Halligan, 2009), ezért a 2. kísérletben megvizsgáltuk azt, hogy hogyan változik az elsősorban striatumhoz kötődő implicit szekvenciatanulás hipnózisban. 14 magas hipnabilitású személy vett részt a vizsgálatban. Egy 15 blokkból álló ASRT feladatot használtuk egy éber és egy hipnózisos feltételben. A két adatfelvétel között egy hónap telt el. A feltételek sorrendjét személyenként változtattuk, és különböző szekvenciák szerepeltek a két feltétel során. Emellett a frontális lebenyhez köthető végrehajtó funkciókat is felmértük Wisconsin kártyaszortírozási és verbális fluencia feladatokkal. A vizsgálati személyek nagyobb mértékű szekvenciatanulást mutattak hipnózis alatt szemben az éber feltétellel. Továbbá, a végrehajtó funkciók szintje interakcióban állt ezzel a serkentő hatással. Azoknak a vizsgálati személyeknek, akik jobb végrehajtó funkciókkal rendelkeztek, gyengébb volt a tanulásuk az éber feltételben és javult hipnózis alatt. Ezzel szemben a gyengébb végrehajtó funkciókkal rendelkező személyek hasonló mértékben tanultak a két feltételben. Az eredmények alátámasztják azt a feltételezést, hogy a frontális lebeny-függő folyamatok kompetitív viszonyban állnak a bazális ganglionokhoz köthető tanulási folyamatokkal: ha a frontális lebenyhez köthető funkciókat megzavarjuk, jobb implicit szekvenciatanulási teljesítmény érhető el. Ezek az eredmények összhangban állnak a modell-függő és modell-független tanulási folyamatok megkülönböztetésével és feltételezett szerepével a tanulásban.

Az első két kutatás az implicit szekvenciatanulás első, online szakaszára fókuszált, azonban az elsajátított ismeretekben jelentős változások következhetnek be a gyakorlás utáni, offline szakaszban is (Krakauer & Shadmehr, 2006; Robertson, 2009; Song, 2009). Eddig viszonylag kevés kutatás foglalkozott az implicit szekvenciatanulás konszolidációjának feltérképezésével, és ezeknek a kutatásoknak a többsége nem vizsgálta külön a szekvencia-specifikus és általános készségtanulást (pl. Robertson, Press, & Pascual-Leone, 2005; Shadmehr & Brashers-Krug, 1997; Walker & Stickgold, 2004), továbbá főleg csak a fiatal felnőtt életkori csoportra fókuszáltak. Ezért a 3. kísérlet az implicit szekvenciatanulás konszolidációját vizsgálta egészséges fiatal (N=58) és idős felnőtteknél (N=71). Annak érdekében, hogy pontosabb képet kapjunk a konszolidáció idői dinamikájáról, 3 késleltetési időt vizsgáltunk a tanulási és a tesztelési fázis között: 12-, 24 óra vagy pedig 1 hét telt el a két adatfelvétel között. Az első adatfelvétel alkalmával egy hosszabb (25 blokkos) ASRT feladatot használtunk, a második alkalommal pedig egy rövidebbet (5 blokk), ami által lehetőség nyílt annak tesztelésére, hogy mennyire szilárdult meg a korábban megszerzett tudás az adott konszolidációs idő alatt. Külön elemezve a szekvencia-specifikus és általános

készségtanulás offline változásait, azt találtuk, hogy a szekvencia-specifikus tudás megszilárdult a fiatal felnőtt csoportok esetében, tehát ugyanúgy teljesítettek a 2. adatfelvételkor, mint az 1. adatfelvétel végén. Ezzel szemben az idős csoport kis mértékű felejtést mutatott a szekvencia-specifikus tudásban a 2. adatfelvételkor. Ezeket az eredményeket nem befolyásolta a késleltetési idő egyik életkori csoportban sem. Ezzel szemben az általános készségtanulás esetében nagyobb mértékű javulás következett be, minél kevesebb idő telt el a tanulási és a tesztelési szakasz között. Az idősebb felnőtteknél csak a 12 órás késleltetés után jelentkezett javulás. Ezek az eredmények alapján megállapíthatjuk, hogy a konszolidáció nem egy egységes folyamat, hanem eltérően hat a szekvencia-specifikus és az általános készségtanulásra. Továbbá ezeket a folyamatokat különböző módon befolyásolja a késleltetés és az életkori változások. A szekvencia-specifikus tanulás offline stabilizációja feltételezhetően a tanulási szakasz után rövidebb idő alatt következik be (az első 12 órában), majd további, viselkedésben mérhető változás nem történik. Ezzel szemben az általános készségtanulás esetében az offline változások időben elhúzódóbbak, még 1 hét után is tapasztalható javulás a korábbi szakaszban nyújtott teljesítményhez képest.

Összefoglalva a három kísérlet eredményeit, elmondható, hogy az implicit szekvenciatanulás gyerekkorban (12 éves kor előtt) a leghatékonyabb, ezután pedig egy váltás következik be a tanulási stratégiában. A felnőttek feltételezhetően jobban támaszkodnak a már frontális lebenyhez kötődő belső modelljeikre a tanulás során szemben. Ezzel szemben gyerekkorban a bazális ganglionok által mediált modell-független tanulás a dominánsabb, mely hatékonyabban tudja detektálni a környezet szabályszerűségeit egy új készség elsajátítása során. Ezt a feltételezést alátámasztja az is, hogy ha hipnózissal megzavarjuk a frontális lebeny működését, akkor jobb lesz az implicit tanulás. Időskorban pedig mind az online tanulásban, mind a konszolidációban gyengülés következik be. Annak eldöntésére azonban, hogy ennek hátterében pontosan milyen folyamatok állnak, még további vizsgálatok szükségesek.

A három kísérlet eredményei nem csak a tanulási mechanizmusok viselkedéses szintű megértéséhez járulhatnak hozzá, hanem segíthetnek megérteni az agyi plaszticitás életkori változásait gyerekkortól idősorig. Emellett az eredmények segíthetnek olyan fejlődési rendellenességek (pl. autizmus, diszlexia), neurodegeneratív (pl. Parkinson-kór, Huntington-kór) és életkorral járó zavarok (pl. demencia) jobb megértéséhez, melyekben a vizsgált funkciók hátterében álló agyi struktúrák érintettek. Végül pedig eredményeink segítségül szolgálhatnak hatékonyabb diagnosztikus eszközök, fejlesztő és rehabilitációs programok kidolgozásához.

Introduction

Prediction is one of the most fundamental functions of the brain. During every moment of our waking life, the brain is trying to anticipate future sensations. In order to recognize time-based patterns and predict subsequent events, storing and recalling of sequences are required (Hawkins et al., 2009; Janacsek & Nemeth, 2012). Without these skills, it would be impossible to carry out evolutionary adaptive behaviors. Most predictions are based on the implicit learning that occurs when information is acquired from an environment of complex stimuli, without conscious access either to what was learned or to the fact that learning occurred (Cleeremans et al., 1998; Reber, 1993). Implicit sequence learning underlies not only motor, but also cognitive and social skills (Kaufman et al., 2010; Lieberman, 2000; Nemeth et al., 2011; Romano Bergstrom et al., 2011; Ullman, 2004); it is therefore an important aspect of life from infancy to old age. Implicit sequence learning is essential for learning languages, as well as learning to operate appliances, computer applications or musical instruments (D. V. Howard et al., 2004; Romano, Howard, & Howard, 2010). Social skills appear in compound behaviors (including series of perceptions, emotions as well as motor actions) realized in proper sequences and under appropriate circumstances. These skills—for example, dialogues, decision making in social context, communication of emotions, predicting others' behavior based on previous verbal and nonverbal social communication, and adjusting our own behavior based on these predictions—are needed for normal social functioning in various sorts of situations: in the workplace, in the family, in the neighborhood, during recreation, shopping, or in the context of medical and mental care (Heerey & Velani, 2010; Lieberman, 2000; Nemeth & Janacsek, 2011). Furthermore, these skills are crucial for effective participation in educational, training, and rehabilitation programs, for instance in relearning how to walk, reach for objects, and speak after brain injury (D. V. Howard et al., 2004; Nemeth, Janacsek, Balogh et al., 2010).

Despite the growing interest in implicit learning, there has been relatively little or no research focusing on life-long development of implicit sequence learning and on offline processing of implicitly learned information (i.e., consolidation). Here, we present three studies examining 1) implicit sequence learning across life-span, 2) the relationship between frontal-lobe functions and implicit sequence learning, and 3) the consolidation of implicitly learned material in healthy young and elderly adults.

In the theoretical background, first we present the most popular implicit sequence learning tasks which are frequently used in cognitive neuroscience. After, we discuss the

different stages of implicit sequence learning (i.e., online and offline learning). Then, we sum up the results of development and aging studies including what we know to date about the biological background of implicit sequence learning and consolidation in different age groups. Finally, we discuss recent studies highlighting that different learning and memory processes interact in cooperative and sometimes competitive ways in optimizing information processing performance.

Measures of implicit sequence learning

In experimental settings, implicit learning is defined as the acquisition of co-occurrence/dependencies between stimuli or trials, and is expressed only through performance (Peter A. Frensch, 1998; D. V. Howard et al., 2004; Rieckmann & Bäckman, 2009). In the past decades, several tasks have been developed to tap into implicit learning. These tasks can be organized into two main groups based on whether the covariation or the temporal sequence of stimuli has predictive information. We focus now only on the sequence learning tasks (for details of covariation learning see Janacsek & Nemeth, 2012)

A widely used sequence learning task is the finger tapping task (Figure 1). Here, participants are instructed to produce a particular sequence of finger movements either on a response box or by opposing their fingers to their thumb (Doyon et al., 2002; Karni et al., 1995). Performance is measured by the number of correctly produced sequences over a certain time interval (e.g., 30 s). The main problem with this task is that it can be difficult to classify as truly implicit, because participants are aware of the sequence they have to generate. However, it is based on the learning of sequences and the learning is expressed through performance; therefore, we discuss the results based on this task where it is relevant.

A) Task types

B) Sequence types

Deterministic:

Length: shorter, e.g., 23142 (Finger Tapping)
 longer, e.g., 124314234132 (SRT)

Probabilistic: 2r3r4r1r (ASRT)

	High-frequency triplets	Low-frequency triplets
1 – R – 2	e.g., 1 – 4 – 2	(always high)
R – 4 – R	e.g., 1 – 4 – 2	e.g., 1 – 4 – 3

C) Measuring the performance

Figure 1. A) Examples of the most commonly used sequence learning paradigms. B) The main distinction in the case of sequence type is whether it is deterministic or probabilistic. C) Typical examples of performance measures in different sequence learning paradigms. In the finger tapping task, performance is measured by the number of correctly produced sequences over a certain time interval (e.g., 30 sec). In the classical SRT task, sequence learning is measured as the RT difference between sequence (S) and random (R) blocks. In the probabilistic sequence learning tasks (e.g., the ASRT task), sequence-specific learning is measured by comparing responses to the random and sequence elements or more and less predictable events in all blocks.

In the past decade, the serial reaction time (SRT) task (Nissen & Bullemer, 1987) and its modification, the alternating serial reaction time (ASRT) task (J. H. Howard, Jr. & Howard, 1997; Nemeth, Janacek, Londe et al., 2010), have become the most popular implicit sequence learning tasks. In the original version of the SRT task, a stimulus appears at one of four possible locations on the screen, and subjects have to press the button corresponding to

that location. They are unaware that the sequence of subsequent locations (and correspondingly, the sequence of the responses) follows a predetermined order (Janacsek & Nemeth, 2012). For example, in classical SRT tasks, the structure of the sequence is deterministic, with the stimuli following a simple repeating pattern as in the series 213412431423, where the numbers refer to distinct events. Without becoming aware of the sequence, subjects learn the regularity – and as they learn, they produce faster and more accurate responses. When the sequence is changed to a random series of stimuli, subjects become slower and less accurate in responding. In this task, sequence learning is measured by the reaction time difference between sequence and random blocks.

The modified version of SRT is the alternating serial reaction time (ASRT) task (J. H. Howard, Jr. & Howard, 1997; Nemeth, Janacsek, Londe et al., 2010). Here, repeating events alternate with random ones in an eight-element sequence so that the location of every second stimulus in the stream is determined randomly. If, for instance, the sequence is 3214, where the numbers represent locations on the screen, the sequence of the stimuli will be 3R2R1R4R, with R representing a random element. The sequence is thus better hidden than in the classical SRT task. This structure is referred to as a probabilistic second-order dependency (Remillard, 2008). The structure is second-order in that for pattern trials, event $n-2$ predicts event n . It is probabilistic in that these pattern trials occur amid randomly determined ones. In addition, participants do not generally become aware of the alternating structure of the sequences, even after extended practice, and sensitive recognition tests indicate that people do not develop explicit knowledge of the event sequences that are more likely to occur (D. V. Howard et al., 2004; J. H. Howard, Jr. & Howard, 1997; Song et al., 2007).

Because sequence and random stimuli alternate in the ASRT task, some sequences of three events (called “triplets”) occur more frequently than others. In this task we can separate general skill learning from sequence-specific learning, where *general skill learning* refers to increasing speed as the result of practice. In contrast, *sequence-specific learning* refers to the acquisition of sequence-specific knowledge, resulting in relatively faster responses for more predictable high-frequency events compared to less predictable low-frequency events. In the classical SRT task, sequence learning is measured as the RT difference between sequence and random blocks. However, in this difference score, sequence-specific and general skill learning are mixed, because participants also exhibit general skill improvement on sequential blocks. Thus, the higher difference between these two types of blocks could be attributed to sequence-specific and general skill learning together, and we cannot determine the extent of these two types of learning. ASRT allows these types of learning to be separated. Another advantage of

the ASRT task compared to the classical SRT task is that in the ASRT, it is possible to track sequence-specific learning continuously by comparing responses to the random and sequence elements or more and less predictable events in all blocks. In summary, recent research favors the ASRT task, because 1) it is more implicit, 2) it can separate sequence-specific and general skill learning, and 3) it can continuously track these two types of learning across all blocks.

Consolidation of sequence knowledge

Sequence learning does not occur only during practice, in the so-called online periods, but also between practice periods, during the so-called offline periods. The process that occurs during the offline periods is referred to as consolidation, which denotes the stabilization of a memory trace after the initial acquisition; this can result in increased resistance to interference or even improvement in performance following an offline period (Krakauer & Shadmehr, 2006; Nemeth, Janacek, Londe et al., 2010; Robertson, 2009; Song, 2009).

The consolidation can include the integration of recently acquired information with past experiences (memory association), the anatomical reorganization of memory representations (memory translocation), reconsolidation of memory representations after recall (memory reconsolidation), and even the erasure of memory representations, all of which appear to occur outside of awareness and without additional training. These processes can be time dependent or sleep dependent (Stickgold, Fosse, & Walker, 2002; Walker, Brakefield, Seidman et al., 2003; Walker & Stickgold, 2004). Hence, it is essential to differentiate between time-dependent and sleep-dependent consolidation. *Time-dependent* consolidation reflects the stabilization or even improvement of the memory trace after an offline period, irrespective of whether sleep occurred in this period or not. Thus, in this type of consolidation, sleep is not an essential component (Nemeth, Csábi, Janacek, Várszegi, & Mari, in press; Nemeth, Janacek, Londe et al., 2010; Song et al., 2007).

In recent decades, special attention has been given to the role of sleep; for instance references are made to *sleep-dependent* consolidation (Walker & Stickgold, 2004) suggesting that performance improves more when the offline period includes sleep than when it does not. Several studies have shown the critical role of sleep in skill learning consolidation (Fischer, Hallschmid, Elsner, & Born, 2002; Stickgold et al., 2002; Walker, Brakefield, Morgan, Hobson, & Stickgold, 2002). Nonetheless, the results concerning consolidation of sequence knowledge have been mixed, and recent findings indicate that whether or not offline improvements occur at all, and whether they are sleep dependent, varies with factors such as

awareness, the length of the offline period, the type of information to be learned, and the age of the participants (Hallgató, Győri-Dani, Pekár, Janacsek, & Nemeth, *in press*; Nemeth, Hallgato, Janacsek, Sandor, & Londe, 2009; Nemeth & Janacsek, 2011; Nemeth, Janacsek, Londe et al., 2010; Press, Casement, Pascual-Leone, & Robertson, 2005; Robertson, Pascual-Leone, & Press, 2004; Song et al., 2007; Spencer, Gouw, & Ivry, 2007).

Studies on the time course of skill consolidation indicate that there is a “critical period” after the learning phase, which is necessary for the stabilization of memory traces. This time period depends on the task demand, and it varies from 1-2 hours (Robertson et al., 2005), to 5 hours (Shadmehr & Brashers-Krug, 1997; Shadmehr & Holcomb, 1997), or 6 hours (Walker, Brakefield, Hobson, & Stickgold, 2003). For example, using the SRT task, a study found that the offline enhancement increased with the length of delay (Press et al., 2005). In this SRT study no offline enhancement was found 1 hour after the learning phase, but significant enhancement was observed after 4 hours, which further increased after 12 hours. These results suggest that consolidation of sequence knowledge may be a dynamic process. However, this study examined only a shorter stretch of time, so the question can be raised, what happens in skill consolidation after more than 12 hours. In addition, age-related differences in implicit skill learning and consolidation have not yet been comprehensively characterized.

Age-related differences in implicit learning

Two main approaches to implicit learning emerged in developmental neuroscience with a different assessment of how learning abilities change with age: 1) the developmental invariance model and 2) the age-related changes model. Studies supporting the developmental invariance model of implicit learning failed to find significant age-related differences in learning (Meulemans et al., 1998; Vinter & Perruchet, 2000). In support of this view, infant studies have shown that adult-like implicit learning mechanisms exist even in very early infancy (Clohessy, Posner, & Rothbart, 2001; Saffran, Aslin, & Newport, 1996). Developmental invariance models explain this age-independence by linking implicit (or procedural) learning to evolutionarily primitive brain regions, such as the basal ganglia and the cerebellum. These regions are characterized as early-maturation regions and are relatively resistant to neurological impairments (Reber, 1993).

By contrast, the age-related changes models posit that considerable developmental differences can be observed in implicit learning. Several of these studies found that older

children and young adults showed stronger learning effects compared to very young participants (J. Fletcher et al., 2000; Kirkham, Slemmer, Richardson, & Johnson, 2007; Maybery et al., 1995; Thomas et al., 2004). These models accept the fronto-striatal origin of such learning, but they focus on evidence of continued development of these regions that form the basis of the behavioral changes with age (e.g., Thomas et al., 2004).

For example, in an SRT study, Meulemans et al. (1998) found that 6- and 10-year-old children showed similar degrees of learning as young adults. In contrast, Thomas et al. (2004) found that the learning performance of young adults was better than 7- to 10-year-old children. Studies investigating implicit skill learning at older ages also revealed inconsistent results. For example, several studies have demonstrated that, for simple repeating patterns (in the SRT task), the extent of implicit sequence learning in elderly adults was comparable to young adults (P. A. Frensch & Miner, 1994; D. V. Howard & Howard, 1992; D. V. Howard & Howard, 1989). Moreover, in a recent study, Gaillard et al. (2009) found that young (22-year-old), middle-aged (45-year-old), and elderly (71-year-old) participants performed at the same level.

The studies mentioned above used fixed (deterministic) sequences, which can be easily learned, making it less possible to detect age-related differences in learning. Furthermore, they cannot purely determine the acquired sequence-specific knowledge because these tasks (finger-tapping, classical SRT) confound general improvements with sequence-specific learning. Using the ASRT task, which enables to measure the “pure” sequence-specific learning distinguished from general improvements, recent studies have shown that, although elderly adults can also learn the higher-order structure of these complex sequences, they showed age-related deficits (D. V. Howard et al., 2004; J. H. Howard, Jr. & Howard, 1997; Nemeth, Janacsek, Londe et al., 2010). Both young and elderly adults were able to learn third-order dependencies (e.g., 1RR2RR3RR) although the elderly participants performed at a lower level than the younger participants (Bennett, Howard, & Howard, 2007). Whereas several studies investigated implicit learning in children using the ASRT task (Barnes et al., 2008; Barnes, Howard, Howard, Kenealy, & Vaidya, 2010; Nemeth, Janacsek, Balogh et al., 2010), no child-adult comparison of implicit skill learning performance has yet been reported.

Biological background of implicit sequence learning and consolidation

Most models and empirical studies of skill learning highlight the role of the striatum which is part of the basal ganglia (BG) system (Dennis & Cabeza, 2011; Doyon, Bellec et al.,

2009; Hikosaka et al., 1999; Hikosaka et al., 2002; Keele, Ivry, Mayr, Hazeltine, & Heuer, 2003; Kincses et al., 2008; Rieckmann, Fischer, & Bäckman, 2010; Sefcsik et al., 2009). In addition, the cerebellum, motor cortical areas and frontal associative regions in some cases are also active during skill learning (Doyon, Bellec et al., 2009). Recent studies also found hippocampus activation during implicit sequence learning (Albouy et al., 2008; Schendan, Searl, Melrose, & Stern, 2003). However, an ample body of previous research and also classical memory models highlight that the medial temporal lobe (MTL) and especially hippocampus plays a critical role in explicit/conscious/contextual memory formation (Bird & Burgess, 2008; Scoville & Milner, 1957; Squire & Knowlton, 2000; Squire, Stark, & Clark, 2004; Squire & Zola-Morgan, 1991; Zola-Morgan, Squire, & Amaral, 1986). Therefore, to date, the role of the hippocampus remains inconclusive.

Most experiments investigated the brain activity during online sequence learning, and not the consolidation processes (Dennis & Cabeza, 2011; N. Raz et al., 2005; Rieckmann & Bäckman, 2009). Therefore, we have relatively less information about consolidation of sequence knowledge. Some studies highlight that the striatum remains active, while the involvement of other associated brain regions decreases after a consolidation period (Albouy et al., 2008; Doyon, Bellec et al., 2009).

Similarly, developmental differences in implicit skill learning and its biological background have not yet been comprehensively characterized. As previously mentioned, BG were described as early-maturation regions in childhood (Reber, 1993). However, recent studies suggest substantial changes later in childhood, as well; they found an inverted U-shaped developmental trajectory, peaking at around 10-14 years of age (Blakemore & Choudhury, 2006; Giedd, 2008; Giedd et al., 1999). The shape of the BG developmental trajectory is more similar to that of frontal cortical areas than temporal, supporting the view that brain regions that share widespread connections also share similar developmental courses (Giedd, 2008). In an SRT study of Thomas et al. (2004), activity in the caudate (which is a part of the striatum) correlated significantly with behavioral measures of implicit learning for both children and adults, although adults showed greater signal change than children. It is in line with the behavioral results, as adults showed higher sequence learning magnitude compared to children. They found hippocampus activation, as well, with greater signal change in adults.

Regarding the changes in older ages, age-related structural and biochemical losses are pronounced in the striatal complex and connected frontal areas in the adulthood (Dennis & Cabeza, 2011; N. Raz et al., 2005; Rieckmann & Bäckman, 2009). In a recent fMRI study,

Dennis and Cabeza (2011) showed that older adults recruited the MTL for implicit learning, and this activation was significantly greater, while striatal activity decreased in older people compared with young adults during implicit learning. Rieckmann, Fisher & Backman (2010) found similar results: in young adults during the learning session, the activation of the striatum increased, but that of the MTL decreased. By contrast, in older adults, sequence learning positively related to activation increases in both the striatum and MTL. Furthermore, a recent diffusion tensor imaging aging study by Bennett et al. (2011) found that the caudate–dorsolateral prefrontal cortex (DLPFC) and hippocampus-DLPFC tract integrity were related to ASRT sequence learning. The caudate-DLPFC tract integrity decreased in the older ages, mediating age-related differences in sequence learning.

A major approach to this research is through brain imaging and neuropsychological studies; in addition to these, experiments investigating the effects of pharmacological agents provide an opportunity for the better understanding of the biological background of implicit learning (for review see Uddén, Folia, & Petersson, 2010). For example, a study by Frank et al. (2006) showed that the benzodiazepine midazolam, which inactivates the hippocampus, causes explicit memory deficits in healthy participants, but enhances implicit learning. In contrast, a more recent study found impaired implicit learning after the exogenous administration of the stress hormone cortisol (Römer, Schulz, Richter, Lass-Hennemann, & Schächinger, 2011). The engagement of specific brain structures in these phenomena needs to be clarified.

Interaction between learning/memory systems

Recent studies showed that manipulations, such as a demanding secondary task (Filoteo et al., 2010; Foerde, Knowlton, & Poldrack, 2006; Fu & Anderson, 2008), a distractor task inserted between the learning sessions (Brown & Robertson, 2007a), or a neuropharmacological blockage (Frank et al., 2006) had no detrimental effect or even led to performance improvements in implicit learning tasks. Galea, Albert, Ditye, and Miall (2010) also found improvements in implicit sequence learning after the disruption of the DLPFC using theta burst stimulation. In contrast, strengthening the reliance on explicit, conscious strategies during learning resulted in impaired sequence learning (P. C. Fletcher et al., 2005; Darlene V. Howard & Howard, 2001) with greater PFC activity during the acquisition (P. C. Fletcher et al., 2005). These results highlight that learning and memory processes interact in

cooperative and sometimes competitive ways in optimizing information processing performance (Brown & Robertson, 2007a; Poldrack et al., 2001; Poldrack & Packard, 2003).

In line with these findings, recent studies proposed that people use internally stored structured models of the world together with learning the statistical probabilities of the event occurrences in the environment in order to adapt more efficiently to future events (Orban, Fiser, Aslin, & Lengyel, 2008; Tenenbaum, Kemp, Griffiths, & Goodman, 2011). In this framework, as the internal models develop with experiences, interpretations of events become more elaborate and less directly related to the event probabilities of the environment. A recent study argued that existence of multiple learning mechanisms in the brain (cf. model-free vs. model-based learning) with an uncertainty-based arbitration between them would be computationally optimal (Daw et al., 2005). Anchoring this hypothesis biologically, the presumed mechanisms related to these two types of learning (model-free and model-based) were suggested to be mediated by the basal ganglia (e.g., striatum), and frontal- and temporal lobe of the cortex, respectively (Daw et al., 2005). This assumption is consistent with previous studies finding improvement in basal ganglia-dependent model-free learning by reducing the reliance on the PFC/MTL-dependent learning based on internal models (Brown & Robertson, 2007a; Filoteo et al., 2010; Foerde et al., 2006; Frank et al., 2006; Fu & Anderson, 2008; Galea et al., 2010). We plan to interpret our empirical results using this new approach proposed by Daw et al. (2005).

Aims of the studies

Here, we present three studies investigating these questions. In Experiment I, we compared implicit sequence learning from 4 to 85 years of age to determine age-related differences across human life span (Table 1). In Experiment II, we compared implicit sequence learning in hypnosis and in waking alert state to investigate the potentially competitive interaction between two fundamentally different forms of learning: the frontal lobe-mediated model-based learning and basal ganglia-dependent model-free learning. Finally, we investigated consolidation of implicit sequence knowledge in young and elderly adults in order to determine age-related differences not only in online learning, but also in offline processing of the learned material (Experiment III).

	Main question of the study	Participants	Methods	Results
Exp. I*	Age-related changes of implicit sequence learning across life span	N=421, from 4 to 85 years of age, clustered into 9 age groups	20 blocks of the ASRT task to measure implicit sequence learning	A gradual decline of sequence learning was found, with the highest learning performance in the 4- to 12-year-old age groups.
Exp. II**	How the disruption of frontal lobe functions by hypnosis affects implicit sequence learning?	N=14, highly hypnotizable university students (mean age=22 yrs, mean education=15 yrs)	15 blocks of the ASRT task to measure implicit sequence learning, separately in hypnotic and waking alert state (within-subject design)	Hypnosis boosted implicit sequence learning. This boosting effect was mediated by frontal lobe related executive functions.
Exp. III***	The time course of implicit sequence consolidation in healthy young and elderly adults	N=129, 71 young and 58 elderly adults; clustered into three conditions	25 blocks (Session 1) and 5 blocks (Session 2) of the ASRT task; between the two sessions there was a delay of either 12-, 24-hour or 1-week	Sequence-specific knowledge stabilized in young adults, but decreased in elderly, regardless of delay. General skill consolidation was time-dependent, with higher offline improvement after shorter delay in both age groups.

Table 1. The outline of the three experiments. *Janacsek, K., Fiser J., & Nemeth, D. (in press). *Developmental Science*. **Nemeth, D., Janacsek, K., Polner, B., & Kovacs, Z. A. (in press). *Cerebral Cortex*. ***Nemeth, D., & Janacsek, K. (2011). *Journal of Gerontology Series B: Psychological Sciences*, 66B(1), 15-22.

Experiment I

It is widely accepted that children should begin learning sports, music or languages early in their life if they are to develop a high proficiency, because late learners seldom become true champions or famous musicians or gain command of a second language similar to that of a native speaker. These observations contradict traditional measures of the ability of factual learning of declarative memories, which showed that humans become increasingly better at many learning tasks up until their late twenties (Craik & Bialystok, 2006). However, an important component of developing new abilities is related to implicit unconscious statistical learning processes (Hikosaka et al., 2002; Keele et al., 2003) that underlie the acquisition of not only motor but also cognitive and social skills (Doyon, Bellec et al., 2009; Hikosaka et al., 2002; Lieberman, 2000; Poldrack et al., 2005; Ullman, 2001). Thus, to understand complex skill acquisition, the characteristics of both explicit/declarative and implicit learning, such as the differences in their efficiency across the lifespan, must be clarified. In contrast to declarative memory (Tulving & Craik, 2000), the ontogenetic changes

in humans' implicit learning abilities have not yet been comprehensively characterized, and, thus, their role in acquiring new skills efficiently during development is unknown. The main goal of our study was to examine age differences in implicit learning across the human lifespan using the same task for all groups.

The computational underpinnings and the neural substrates of these different kinds of learning mechanisms are also controversial (Henke, 2010). Explicit learning has been linked more closely to medial temporal lobes of the cortex (Dennis & Cabeza, 2011; Squire & Zola, 1996). By contrast, implicit skill learning often requires fine-tuning of the perceptual-motor system based on experience; therefore, most models of implicit skill learning emphasize the role of the basal ganglia and the cerebellum (Cohen, Pascual-Leone, Press, & Robertson, 2005; Dennis & Cabeza, 2011; Doyon, Bellec et al., 2009; Hikosaka et al., 1999; Hikosaka et al., 2002), whereas the role of the hippocampus remains inconclusive (Albouy et al., 2008; Schendan et al., 2003). However, these models focused mostly on motor skill-related learning with less emphasis on more complex skills that could involve learning abstract cognitive dependencies implicitly. The second goal of our study was to relate our behavioral results to the various computational models of explicit and implicit learning. We compared our empirical results using a new approach to the problem of multiple neural substrates of learning proposed by Daw et al. (2005).

In summary, previous studies have addressed the development and aging in implicit skill learning, but no studies have examined age-related differences from childhood to old age with identical methods. Furthermore, in contrast to general skill improvements, using a probabilistic sequence learning task (ASRT) can help us to reveal the age-related differences of the underlying mechanisms of complex skill learning by measuring the sensitivity to raw probabilities of high and low frequency events (i.e., triplets). Therefore, in this study, we compared the implicit probabilistic sequence learning across the age range of 4-85 years.

Methods

Participants

There were 421 participants in the experiment between the ages of 4 and 85 that were clustered into nine age groups (Table 2). None of them suffered from any developmental, psychiatric or neurological disorders. All subjects gave signed informed consent (parental consent was obtained for children), and they received no financial compensation for

participation. All experimental procedures were approved by the Ethics Committee of the University of Szeged.

Group	Age	Gender	Education	Mean RT	Mean Accuracy
4-6-year-old (n=30)	5.31 (0.98)	17 M / 13 F	-	960.06 (214.67)	90.09 (6.34)
7-8-year-old (n=55)	7.09 (0.56)	31 M / 24 F	1.18 (0.39)	773.24 (159.29)	90.66 (7.03)
9-10-year-old (n=35)	9.89 (0.58)	14 M / 21 F	3.2 (0.96)	602.84 (121.03)	93.44 (4.21)
11-12-year-old (n=29)	11.5 (0.5)	21 M / 8 F	4.66 (0.67)	544.15 (95.00)	92.52 (4.23)
14-17-year-old (n=62)	14.89 (1.06)	46 M / 15 F	8.23 (1.02)	452.52 (67.06)	95.44 (2.92)
18-29-year-old (n=63)	23.09 (3.67)	40 M / 23 F	15.45 (2.6)	401.79 (50.85)	95.47 (2.45)
30-44-year-old (n=59)	35 (4.24)	24 M / 35 F	16.64 (3.1)	419.85 (58.68)	95.85 (2.98)
45-59-year-old (n=36)	50.8 (5.07)	12 M / 24 F	14.18 (3.58)	526.7 (112.99)	97.4 (3.45)
60-85-year-old (n=52)	69.85 (6.16)	16 M / 36 F	13.39 (3.04)	634.37 (126.54)	96.92 (2.38)

Table 2. Demographic data, such as mean age, gender (Male, Female), and mean education (in years), are shown in the table. In addition, mean reaction time (RT) in milliseconds and accuracy (percentages) are also presented for all age groups. In all columns, numbers in parentheses represent standard deviation.

Task and procedure

We used the ASRT task (J. H. Howard, Jr. & Howard, 1997; Nemeth, Janacek, Londe et al., 2010) where a stimulus (a dog's head) appeared in one of the four empty circles arranged in a line on a computer screen. The participants were instructed to respond to different stimulus events by pressing the corresponding response keys (Z, C, B or M) as fast and accurately as possible. The ASRT task consisted of 20 blocks with 85 key presses in each block. The first five responses of each stimulus block served for practice only, and then the eight-element alternating sequence (e.g., 2R1R3R4R, where numbers represent stimulus locations, and R represent random) was repeated ten times within a block. Stimuli were presented 120 ms after the response to the previous stimulus. Between blocks, the participants received feedback on the screen about their overall reaction time (RT) and accuracy. The computer program generated a different repeating ASRT sequence of the 4 locations for each participant using a permutation rule such that each of the six unique permutations of the 4 repeating events occurred with equal probability.

To determine the amount of explicit knowledge the participants acquired about the task, a short questionnaire was administered after the experimental session (Song et al., 2007).

This questionnaire included increasingly specific questions, such as “Have you noticed anything special regarding the task?”, “Have you noticed some regularity in the sequence of stimuli?”. The experimenter rated participants’ answers on a 5-point scale where 1 denoted “Nothing noticed” and 5 denoted “Total awareness”. None of the participants reported noticing the hidden repeating sequence.

Statistical analysis

The ASRT allows a comparison between responses to high- and low-probability events. For example, if the eight-element sequence is 2R1R3R4R, 2x1, 1x3, 3x4, and 4x2 would occur often (high frequency triplets) because two consecutive stimuli of the repeating sequence (e.g., 231 consisting 2R1) as well as two consecutive random elements by chance (e.g., the same 231 consisting R3R) could form these triplets. By contrast, 2x3 or 4x1 would occur less frequently (low frequency triplets) because they could never be obtained consisting two consecutive sequence elements. Of the 64 possible triplets, sixteen triplets were high frequency triplets, occurring 62.5% of the time, whereas the remaining 48 triplets were low frequency triplets, occurring 37.5% of the time. Thus, each low frequency triplet occurred in approximately 0.8% of the total number of trials, whereas each high frequency triplet occurred about 5 times more often, in approximately 4% of the trials. For each response, we defined whether it was in response to a high- or a low frequency element, depending on whether the element was more or less predictable based on the previous two items.

Following the method of previous studies (D. V. Howard et al., 2004; Song et al., 2007), two kinds of low frequency triplets were excluded from our analyses: repetitions (e.g., 222, 333) and trills (e.g., 212, 343). Repetitions and trills were low frequency for all participants, and in previous studies, the participants often showed pre-existing response tendencies towards them (D. V. Howard et al., 2004; Soetens, Melis, & Notebaert, 2004). After eliminating these triplets, previous studies have found that, following the practice, participants responded more quickly to the high than to the low frequency triplets, revealing a sequence learning effect (D. V. Howard et al., 2004; J. H. Howard, Jr. & Howard, 1997; Nemeth, Janacsek, Londe et al., 2010; Song et al., 2007). In addition, general skill learning can be observed in the ASRT task in the overall increase of the response speed, irrespective of the triplet type. Thus, we were able to separately measure sequence-specific and general skill learning in the ASRT task.

We calculated the mean accuracy of all trials and the median reaction time (RT) of correct responses separately for high and low frequency triplets. The accuracy and RT measures were analyzed by mixed-model analysis of variance (ANOVA) with TRIPLET TYPE (high vs. low frequency) as the within-subject factor, and AGE (9 groups) as the between-subject factor. All significant results are reported together with the η_p^2 effect size and Greenhouse-Geisser ε correction factors where applicable. Post-hoc analysis was conducted by Fisher's LSD pairwise comparisons.

Results

Overall RT's significantly differed among the age groups (main effect of AGE: $F(8,412)=107.11$, $p<0.001$, $\eta_p^2=0.675$; Table 2). The RT decreased significantly between each group from 4-6 to 18-29 years of age ($p's<0.04$), they were similar between the age groups of 18-29 and 30-44 ($p>0.38$) and significantly increased after 44 years of age ($p's<0.001$) (Figure 2a). The accuracy monotonically increased over the years (main effect of AGE: $F(8,412)=16.94$, $p<0.001$, $\eta_p^2=0.25$) (Figure 2b).

The comparison of RT to high vs. low probability triplets showed a surprising pattern of implicit sequence learning across the age groups. Even though there was a significant sequence learning at all ages because the RT's were faster for high frequency than to low frequency triplets (main effect of TRIPLET TYPE: $F(1,412)=333.7$, $p<0.001$, $\eta_p^2=0.45$) (Figure 2c), the magnitude of this difference was not uniform (TRIPLET TYPE x AGE interaction: $F(8,412)=6.79$, $p<0.001$, $\eta_p^2=0.12$). The post-hoc test revealed that learning was significantly higher in the 4- to 12-year-old groups than in any other group in the 14-85 range ($p's<0.02$). There was no difference in learning between the 14-59 years of age ($p's>0.37$), whereas the magnitude of learning decreased significantly in the 60-85-year-old group ($p's<0.02$). Thus, sequence learning was uniformly effective until the age of 12 where it reduced significantly and remained at a lower level until the age of 60 (Figure 2c).

Figure 2. Sequence learning in all groups. Reaction time (A) and accuracy (B) for high and low frequency triplets are plotted. Learning measures of RT (C) and accuracy (D) represents the RT/ACC difference between low- and high-frequency triplets. Error bars indicate standard error of mean (SEM).

However, it is a long-standing issue in developmental and aging studies how to compare groups with different baseline speeds. A customary approach to this problem is to analyze the data using z-transformation. Therefore, we calculated the z-scores within each participant (thus, each participant's own mean and SD was used to transform that participant's data (see, for example, Christ, White, Mandernach, & Keys, 2001) and conducted an ANOVA based on these z-scores (Figure 3). ANOVA revealed significant sequence-specific learning (main effect of TRIPLET TYPE: $F(1,412)=320.12$, $p<0.001$, $\eta_p^2=0.44$), but the extent of learning differed across groups (TRIPLET TYPE $\times$ AGE interaction: $F(8,412)=8.91$, $p<0.001$, $\eta_p^2=0.15$). We found that the participants from 9 years of age showed similar extent of sequence learning as the adult groups to 44 years of age (p 's >0.25), but the learning in 4-8 years of age was smaller compared to these adult groups (p 's <0.014). At older ages, there was a decline in the sequence learning, with both the 45-59 and 60-85-year-old group differing significantly from the groups between the 11-44 years of age (p 's <0.025).

Z-transformed RT data by age groups

Figure 3. Sequence learning measured by the z-transformed RT data in all groups. Error bars indicate SEM.

The analysis of the response accuracy further enhanced the picture emerging from the results with RTs. We found a significantly greater accuracy for high- than low-frequency triplets (main effect of TRIPLET TYPE: $F(1,412)=217.14$, $p<0.001$, $\eta_p^2=0.345$). Although all age groups older than 6 showed significant sequence learning (p 's <0.011), the age groups differed significantly in the strength of the sequence learning (TRIPLET TYPE $\times$ AGE interaction: $F(8,412)=3.73$, $p<0.001$, $\eta_p^2=0.07$). Whereas groups between 7 and 44 years of age showed similar degrees of learning, this was significantly higher than the youngest (4-6) and the two oldest (45-59 and 60-85) groups (p 's <0.03) (Figure 2d).

Discussion of Experiment I

The goal of the present study was to investigate the differences in implicit skill learning across the human life span. This work extends previous studies (Bennett et al., 2007; Gaillard et al., 2009; J. H. Howard, Jr. & Howard, 1997; Meulemans et al., 1998; Thomas et al., 2004) in two ways: 1) it examined a wide range of ages between 4-85 years, and 2) it used a probabilistic task, which enabled us to measure the “pure” sequence-specific learning defined by the sensitivity to raw probabilities of high and low frequency events. We found that the 4- to 12-year-old age groups showed the strongest learning effect measured by the raw RT difference scores. Around the age of 12, we found a striking transition to less pronounced sequence-specific learning, as measured by smaller differences between the responses to high and low frequency triplets. Confirming earlier results (D. V. Howard et al., 2004; J. H. Howard, Jr. & Howard, 1997; Nemeth, Janacsek, Londe et al., 2010), we found

that this learning capacity was significantly reduced in the oldest age group. Thus, in contrast to the developmental invariance (Reber, 1993) and the age-related changes approaches (Meulemans et al., 1998; Vinter & Perruchet, 2000), our results demonstrate a gradual decline in learning across the lifespan.

Sequence learning scores based on the accuracy and raw reaction time showed different curves: the former one is a bell-shaped curve, whereas the latter is a gradually declining curve (Figure 2c-d). Hence, these two types of learning scores can reflect different underlying mechanisms and brain systems. The accuracy learning score may be more related to attention, mainly voluntary attention, whereas the RT learning score may be related to involuntary attention and intuitive processes (Burgess, Gilbert, & Dumontheil, 2007; Prinzmetal, McCool, & Park, 2005). In other words, we propose that more effective response selection processes are essential to achieve a higher general accuracy and also sequence-specific learning measured by accuracy. As response selection is mediated by frontal and prefrontal areas (e.g., anterior cingulate cortex, Aarts & Roelofs, 2011; Dum & Strick, 1993; Medford & Critchley, 2010; Roelofs, Van Turenout, & Coles, 2006), the relatively weaker performance in children and elderly groups may be due to the underdeveloped/deteriorating attentional brain circuits connected to the frontal lobe. It is also in line with this assumption, that elderly groups could maintain a high general accuracy rate only with a trade-off in reaction time.

Our study raised a methodological issue, which affects the developmental studies in general. It is a long-standing issue in the literature how to compare groups with different baseline speeds. It could be argued that the youngest groups in our study have larger raw RT learning scores (more difference between high and low frequency triplets) because they have more room to do so, given their longer RTs. However this argument does not seem to hold in our study, because the oldest group showed equally long RTs as 9-10 year-olds, yet they had a four-fold reduced learning score (Figure 2). Along a similar line of argument, the accuracy measures of the youngest group (4-6 year-olds) and the oldest groups (45-85 year-olds) showed a marked difference (~8%) in general accuracy, giving more room for the young group to produce larger differences between high and low frequency triplets, yet the actual sequence-learning (i.e. high-low differences in accuracy) showed no difference. In contrast, the two smallest age groups had almost identical general accuracy yet there was a more than two-fold increase in sequence learning in the second youngest group (7-8 year-olds). Thus, our data suggests no linear relationship between the general magnitude of reaction time/accuracy and the learning measures.

A second customary approach to the problem of comparing groups with different baseline speeds is to analyze the data using z-transformation. Z-scores of our results show a different picture than raw the RT data analysis: the learning performances from 9 to 44 years of age are similar with weaker performance in the younger and older age groups. The z-transformation is often used to control general processing speed across different age groups in developmental studies. However, the main function of z-transformation is not to control the processing speed, but to normalize the distribution of responses. Thus, z-transformation has mathematical assumptions about the form of distributions and is therefore not theory independent (Yap, Balota, Sibley, & Ratcliff, 2012). Z scoring fully adjusts for processing speed only if all participants have the same type of distributions. Therefore using z-scores in developmental studies might be misleading. Furthermore, it is unclear how general processing speed and variability contributes to learning and performance in different ages. In recent years, several studies analyzed the variability and noise across a wide age-range (Der & Deary, 2006; Hultsch, MacDonald, Hunter, Levy-Bencheton, & Strauss, 2000; Rabbitt, Osman, Moore, & Stollery, 2001). For example, Rabbitt et al. (2001) found that people's fastest RT's were relatively unaffected by age, but the number of unnecessarily slow responses was higher in older ages, and, thus, the increase in the mean RT was a result of increasing variability, which was an important component of cognitive aging. Moreover, several studies outline that the noise and the performance variability enables adaptive plasticity of motor skills (Slifkin & Newell, 1998, 1999; Turner & Brainard, 2007) and high variability can support effective learning and performance (Sanger, 2010). Thus, based on previous studies and on our analyses, we suggest that both the processing speed (mean RT) and variability are inherent aspects of development and aging. We believe that the z-transformation eliminates these inherent aspects of learning, making the transformed results difficult to understand and explain.

Based on the reasoning above, we propose that, the raw RT difference between the high and low frequency triplets in the ASRT task is a measure of human sensitivity to the relative raw probabilities of events observed implicitly in their environment. Thus, our results show a marked decrease in this sensitivity around the age of 12, which is in contrast to the traditional view of a steady improvement of cognitive learning abilities until late in adulthood (Craik & Bialystok, 2006). However, this discrepancy might be explained based on a shift in the structural development of implicit learning. Although the raw probabilities of the sensory environment are important for learning and both infants (Aslin, Saffran, & Newport, 1998; Fiser & Aslin, 2002; Saffran et al., 1996; Saffran, Johnson, Aslin, & Newport, 1999) and

adults (Fiser & Aslin, 2001; Hunt & Aslin, 2001) are highly sensitive to them, there is an ongoing debate on how using these simple probabilities can lead to highly complex knowledge of the world, such as sensory invariances and development of a language (Gomez & Gerken, 1999; Marcus, Vijayan, Rao, & Vishton, 1999). Recent studies proposed that using an internally stored structured model of the world (model-based learning) together with probabilistic model-free learning could help to address this issue and also provide evidence that humans might implement such a strategy during implicit learning (Orban et al., 2008; Tenenbaum et al., 2011). Support for the separated and complementary nature of the prefrontal- (PFC) and medial temporal lobe (MTL)-dependent learning based on internal models vs. BG-dependent model-free learning comes from various studies investigating learning under specific conditions. These studies showed that obstructing the PFC and/or MTL by a demanding secondary task (Foerde et al., 2006) do not adversely affect implicit learning. Other studies found that inserting a task between the learning sessions (Brown & Robertson, 2007a, 2007b), performing a working memory and an implicit learning task simultaneously (Filoteo et al., 2010), or a neuropharmacological blockage (Frank et al., 2006) even had a positive effect on performance in implicit learning task. Importantly, it is known that the cortical areas connected to the internal models related to model-based learning become truly functional late in the development, around age of 12 (Blakemore & Choudhury, 2006; Giedd et al., 1999), which is about the age when we found the sudden decrement in sensitivity to the relative raw probabilities. We propose, that this enhanced functionality signals the shift when the system adapts efficiently to more complex aspects of the world by relying more on internal model-based interpretations, while somewhat neglecting the raw probabilities of the sensory input (Figure 4a-b), and therefore, decreasing the ability to develop and stabilize fundamentally new basic competences. Thus the seemingly paradoxical result of gradually becoming less sensitive to basic statistics, if timed appropriately, could be the optimal strategy for human implicit learning in general.

Figure 4. Competition between model-based and model-free neurocognitive subsystems of skill learning. (A) Before adolescence, underdeveloped internal models (dashed boundary) have little influence on interpretations of detected statistical probabilities of events in the environment (dashed arrows). Skill learning performance is determined by detection of probabilities. (B) From adolescence, well-developed internal models (solid boundary) strongly modulate the interpretations of observed statistics. This helps extracting complex relations but relatively impairs measuring and utilizing raw probabilities in skill learning (dotted arrow). (C) In older ages, skill learning performance decreases. This decline could be caused by the reduced sensitivity to statistical probabilities, increasingly rigid internal models and/or weaker connection between these systems.

Our results did not reveal any differences between the young adults and middle-age groups. Salthouse's (1996) "simultaneity mechanism" theory of cognitive aging predicts the age-related deficits in probabilistic sequence learning (Curran, 1997; J. H. Howard, Jr. & Howard, 1997). Feeney, Howard & Howard (2002) found age-related deficits in pattern sensitivity in "older" (mean age: 49.4) compared with "younger" (mean age: 41.4) middle-

aged groups. These different results could be related to that Feeney used a smaller sample size, a longer version of the ASRT and different method of analysis.

What are the underlying mechanisms of the decreased performance of the elderly group? Several studies have found both structural and functional impairments in the fronto-striatal circuitry in older ages (Bennett et al., 2011; Giorgio et al., 2010) and there was also evidence for recruiting MTL to learn sequences implicitly (Dennis & Cabeza, 2011; Rieckmann et al., 2010). Within the computational framework proposed by Daw and colleagues (2005), these findings can be interpreted as a deterioration in three mechanisms that contribute to the age-related decline in skill learning: 1) reduced detection of probabilities, 2) rigidity of internal models and/or 3) more restricted connections between internal models and probability detection (Figure 4c). Thus, not only the model-free, but also the model-based learning, might be limited in older ages. Future studies are needed to systematically examine the underlying neural mechanisms of age-related differences in skill learning.

In summary, based on our raw RT results we suggest that acquiring fundamentally new skills that cannot be derived from skills already possessed is the most effective before adolescence, and it might be largely based on the fronto-striatal circuitry, in agreement with skill learning- (Doyon, Bellec et al., 2009; Hikosaka et al., 1999; Hikosaka et al., 2002) and computational learning models (Daw et al., 2005). This result can be explained by a competition between two fundamentally different forms of learning (model-free and model-based) suggesting that after adolescence frontal lobe-mediated model-based learning dominates in the expression of implicit sequence knowledge, while before adolescence basal ganglia-dependent model-free learning is more influencing. As a growing body of data has shown that frontal lobe-mediated processes are disrupted in hypnosis, we tested this assumption in Experiment II by comparing implicit sequence learning in hypnosis and in waking alert state.

Experiment II

As rapid and reversible changes of cognitive processing are encountered in hypnosis, this phenomenon is an excellent tool of research in the cognitive neurosciences (Egner et al., 2005; A. Raz & Shapiro, 2002). Regarding the neural background of hypnosis, studies demonstrated that people (especially with high susceptibility to hypnosis) show decreased performance on some frontal lobe-related tasks in hypnosis (Kaiser, Barker, Haenschel, Baldeweg, & Gruzelier, 1997; Kallio, Revonsuo, Hamalainen, Markela, & Gruzelier, 2001;

Wagstaff, Cole, & Brunas-Wagstaff, 2007). More recent studies suggest reduced functional brain connectivity between cortical areas in hypnosis, and this is especially typical for frontal areas (Fingelkurts et al., 2007; Oakley & Halligan, 2009). Hypnosis temporarily disconnects certain frontal areas from the anterior cingulate cortex and other brain areas, disturbing the frontal attentional control and executive system (Egner et al., 2005; Gruzelić, 2006; Kaiser et al., 1997).

In our experiment, we used hypnosis as a tool to reduce the competition between frontal lobe-related model-based and striatum-related model-free systems. The main question of the study was how the disruption of frontal lobe functions by hypnosis affects sequence learning. Highly hypnotizable young adults performed the ASRT task both in waking alert and hypnotic state. In addition, executive functions were assessed by the Wisconsin Card Sorting Test (Anokhin, Golosheykin, Grant, & Heath, 2010; Heaton, Chelune, Talley, Kay, & Curtiss, 1993) and Verbal Fluency Task (Spreen & Strauss, 1991) in order to investigate the possible interactions between frontal lobe functions and the effect of hypnosis on sequence learning.

Methods

Participants

Fourteen healthy, right-handed students from the University of Szeged participated in the experiment (12 females; mean age: 22.70, SD: 1.70; mean years of education: 15.50, SD: 1.58). All participants provided signed informed consent. They received no financial compensation for participating in the study. The study was approved by the Psychology Ethics Committee at the University of Szeged, Institute of Psychology.

Participants were selected from a pool of pre-tested subjects, on the basis of their hypnotic susceptibility. Hypnotizability was measured using the Hungarian version of the Harvard Group Scale of Hypnotic Susceptibility: Form A (HGSHS:A, Shor, Orne, & Press, 1962). Scoring procedure was based on the original English version (scores ranging from 0 to 12). Similarly to previous studies (Halsband, 2006; Kallio et al., 2001) we defined high hypnotizability as having 8 or higher score on the HGSHS:A. The mean hypnotizability score of the participants was 9.07 (SD = 0.997; range from 8 to 12).

Tasks

Sequence learning task – We used the same ASRT task as in the Experiment I. The task consisted of one practice block with random stimuli, and 15 blocks with the alternating pattern described above.

Fluency task – In this task, participants are instructed to produce as many words belonging to the same category (animals, supermarket) as possible in 60 seconds, without repetitions, synonyms, or generated forms of the same word (Spreen & Strauss, 1991). The average number of correct words was used as the performance score. Higher score reflects better frontal lobe functions (Baldo, Schwartz, Wilkins, & Dronkers, 2006).

Wisconsin Card Sorting Test (WCST) – This task is one of the most specific tests of prefrontal functions (Anokhin et al., 2010; Heaton et al., 1993). Participants are required to derive a correct card sorting rule based on a trial-by-trial feedback. As the rule changes without warning, the participant has to modify the previously learned response strategy on the basis of the feedback information. A key indicator of cognitive flexibility is the number of perseverative errors that occur when the participant persists in using the old strategy despite the negative feedback. A lower number of perseverative errors indicate better frontal lobe functions.

Design and procedure

All tests were conducted on an individual basis. Participants performed the ASRT task in both the alert and the hypnotic state, with the same standard instructions. The order of the two conditions was counterbalanced across participants. The delay between the two sessions was 1 month. Fluency and WCST task was administered once, in a third session in alert state. Two participants did not take part in the third session due to time schedule problems.

A skilled hypnotist therapist (the author, Z.A.K.), who has extensive experience with hypnosis, tape-recorded the induction, instructions, and dehypnotizing phases (similarly to Szendi et al.'s study (2009)). This recording was played to each participant. The type of hypnosis induction, similarly to that of the hypnotizability scale, was essentially relaxational.

The induction took approximately 14 minutes. After the induction, the hypnosis session began. When participants had completed half of the ASRT task, we played an approximately 30 s long induction in order to maintain their hypnotic state (in the wake condition, subjects had a 30 s rest). In the hypnosis condition, after the task had ended, the dehypnosis instruction was played.

Statistical analysis

Since the participants' accuracy remained very high throughout the test (as is typical, the average was over 92% for both conditions; (J. H. Howard, Jr. & Howard, 1997; Nemeth, Janacek, Londe et al., 2010), we focused on RT for the analyses reported. We calculated medians for correct responses only, separately for high- and low-frequency triplets and for each participant and each third of the stimulus blocks (1-5, 6-10 and 11-15).

To compare sequence learning between hypnosis and alert condition, and between groups with high and low executive functions, we conducted repeated measures and mixed design ANOVAs with LSD post hoc tests. We reported the relevant effect sizes: η_p^2 for main effects and interactions, and Cohen's *d* measures for post hoc tests.

Results

A repeated measures ANOVA was conducted with TRIPLET TYPE (2: high vs. low), BLOCK (3: 1-5, 6-10 and 11-15) and CONDITION (2: alert and hypnosis) as within-subjects factors. ANOVA revealed significant sequence-specific learning (main effect of TRIPLET TYPE: $F(1,13)=16.21$, $p=0.001$, $\eta_p^2=0.55$), which increased with practice (TRIPLET TYPE x BLOCK interaction: $F(2,26)=9.36$, $p=0.001$, $\eta_p^2=0.42$). The two states differed significantly from each other (TRIPLET TYPE x CONDITION interaction: $F(1,13)=7.08$, $p=0.02$, $\eta_p^2=0.35$): sequence learning was 2.5-times higher under hypnosis than in the waking alert state (Figure 5). Independently from sequence learning, general reaction time decreased with practice (main effect of BLOCK: $F(2,26)=4.93$, $p=0.034$, $\eta_p^2=0.27$). Other main effects and interactions were not significant (all p 's > 0.34), thus the general reaction time was similar in the waking alert and hypnotic state (main effect of CONDITION: $F(1,13)=0.12$, $p=0.73$, $\eta_p^2=0.009$). In addition, the rate of sequence learning process was also similar between the two conditions (TRIPLET TYPE x BLOCK x CONDITION: $F(2,26)=0.42$, $p=0.63$, $\eta_p^2=0.03$). Thus, hypnosis affected only sequence-specific learning (the difference between RTs for low and high probability events), and not the general reaction time. As we used a within-subject design with two learning sessions, a further analysis was conducted to test the possible effect of whether hypnosis was in the first or in the second session and ANOVA revealed no order effect on sequence learning.

Figure 5. Sequence learning across blocks is plotted for waking alert (A) and hypnotic state (B), separately. Sequence learning performance (measured by the reaction time differences between high- and low-predictability events) was higher in the hypnotic state compared to the waking alert condition (C). Error bars represent SEM.

To calculate a composite score for executive function, first we transformed measures of fluency task and WCST into z-scores. Then, we averaged these two transformed data into a composite score. Based on the median of this composite measure, we assigned half of the participants to the higher and other half to the lower executive function group. To compare sequence learning between the high- and the low-executive-function groups, in the alert and the hypnosis conditions (see Figure 6), a mixed design ANOVA was conducted with TRIPLET TYPE (2: high vs. low), BLOCK (3: 1-5, 6-10 and 11-15) and CONDITION (2: alert and hypnosis) as within-subjects factors and GROUP (2: high vs. low executive function) as a between-subject factor.

The general reaction time was similar in the two groups (main effect of GROUP: $F(1,10)=1.5$, $p=0.25$, $\eta_p^2=0.13$), and it was not affected differently by the two conditions (CONDITION x GROUP interaction: $F(1,10)=0.71$, $p=0.42$, $\eta_p^2=0.07$). The TRIPLET TYPE x CONDITION interaction almost reached the significance ($F(1,10)=4.5$, $p=0.06$, $\eta_p^2=0.31$) replicating that participants exhibited greater sequence learning under the hypnosis condition compared to the waking alert condition. The overall sequence learning was similar in the high and low executive function groups (TRIPLET TYPE x GROUP: $F(1,10)=3.07$, $p=0.11$, $\eta_p^2=0.24$). Although the TRIPLET TYPE x CONDITION x GROUP interaction did not reach significance ($F(1,10)=1.72$, $p=0.219$, $\eta_p^2=0.15$), the LSD post hoc tests revealed that participants with higher executive functions showed smaller sequence learning in the waking alert state compared to the hypnotic condition (Figure 6C; $p=0.03$, $d=0.94$), while participants

with lower executive functions showed similar extent of sequence learning (Figure 6E; $p=0.58$, $d=0.25$). In addition, the learning performance of the high executive function group was significantly smaller compared to the low executive function group in the waking alert state ($p=0.04$, $d=1.31$), while it was similar in the hypnotic state ($p=0.51$, $d=0.396$).

Figure 6. Sequence learning across blocks for the high executive function group in waking alert (A) and hypnotic state (B), as well as sequence learning across blocks for the low executive function group in waking alert (D) and hypnotic state (E) is plotted. Participants with high executive functions showed smaller sequence-specific learning (measured by reaction time differences between high and low probability events) in the alert state compared to the hypnosis condition (C), while participants with lower executive functions showed similar extent of sequence learning in the waking alert and hypnotic state (F). Error bars represent SEM.

Discussion of Experiment II

Taken together, we found enhanced sequence learning in hypnosis. Our results provide support for the idea that learning and memory processes may not only involve the engagement of specific neuroplastic mechanisms, but may also rely upon the disengagement of interacting systems (Brown & Robertson, 2007a, p. 149). Our finding is in line with previous studies

demonstrating that manipulations reducing the reliance on frontal lobe-dependent processes improved BG-dependent learning performance (e.g., Filoteo et al., 2010; Galea et al., 2010).

The improved sequence learning in hypnosis could be attributed to the disruption of attentional control and executive system (Kaiser et al., 1997; Kallio et al., 2001; Wagstaff et al., 2007) by weakening the engagement of the frontal lobe and/or the interconnectivity between related brain areas (Egner et al., 2005; Fingelkurts et al., 2007; Gruzelier, 2006; Oakley & Halligan, 2009). This could diminish the competition between two fundamentally incompatible modes of learning: 1) PFC/MTL-mediated model-based, attention-dependent processes vs. 2) BG-dependent model-free learning (Ashby, Alfonso-Reese, Turken, & Waldron, 1998; Filoteo et al., 2010; Henke, 2010; Poldrack et al., 2001). Reducing the reliance on the former could have improved sequence learning capacity by heightening the sensitivity to statistical probabilities, which is essential for automatic, procedural mechanisms (Daw et al., 2005). This interpretation is consistent with the result that participants with better executive functions showed decreased sequence learning in the alert condition, while subjects with lower executive functions, suggesting that in the alert state relying more on attentional processes prevented the learning of statistical contingencies to a greater extent (P. C. Fletcher et al., 2005). In the hypnotic state, participants with higher executive functions shifted from relying on frontal lobe-related attentional processes to automatic, procedural-based mechanisms, resulting in enhanced sequence learning. However, future neuroimaging studies need to corroborate these results providing direct evidence for the underlying brain systems.

In sum, we found that hypnosis substantially boosted sequence learning in healthy young adults. This result supports the proposed model of Experiment I highlighting the competition between PFC/MTL-mediated model-based and BG-dependent model-free learning systems. By disrupting the former system we can improve learning performance in adults, probably through strengthening the reliance on model-free processes. In addition, our finding sheds light not only on the competitive nature of brain systems in cognitive processes, but also could have important implications for training and rehabilitation programs, especially for developing new methods to improve human skill learning.

Experiment III

Skill learning does not occur only during practice, in the online periods, but also between practice periods, during the offline periods. The process that occurs during the offline periods is referred to as consolidation which means stabilization of a memory trace after the

initial acquisition (Krakauer & Shadmehr, 2006; Nemeth, Janacsek, Londe et al., 2010; Robertson, 2009; Song, 2009). Studies on the time course of skill consolidation indicate that there is a “critical period” after the learning phase, which is necessary for the stabilization of memory traces. This time period depends on the task demand, and it varies from 1-2 hours (Robertson et al., 2005), to 5 hours (Shadmehr & Brashers-Krug, 1997; Shadmehr & Holcomb, 1997), or 6 hours (Walker, Brakefield, Hobson et al., 2003). These results suggest that offline learning may be a dynamic process. However, these studies examined only a shorter stretch of time, so the question can be raised, what happens in skill consolidation after more than 12 hours. Therefore, the aim of the present study was to investigate the time course of implicit skill consolidation after a relatively longer stretch of time.

In addition, several studies demonstrated that older adults show age-related deficits in online skill learning (Curran, 1997; D. V. Howard et al., 2004; J. H. Howard, Jr. & Howard, 1997; Janacsek, Fiser, & Nemeth, *in press*). However, little is known about the effects of aging on skill consolidation and its time course (e.g., Siengsukon & Boyd, 2009a; Spencer et al., 2007). Therefore, we planned to compare skill consolidation in healthy young and elderly adults.

Previous research on consolidation using the ASRT task focused primarily on whether sleep affects skill consolidation and has concluded that these processes are not influenced by sleep (Nemeth, Janacsek, Londe et al., 2010; Song et al., 2007). For example, in a recent study, Nemeth et al. (2010) investigated implicit skill learning after 12-hour offline period. They showed that general skill learning improved offline in both the young and older groups, with the young group improving more than the old group. However, the improvement was not sleep-dependent, in that it was not relevant whether the 12-hour offline period included sleep or not. They did not find sequence-specific offline improvement in either age group, and similarly to general skill learning it was not influenced by sleep. However, this study did not examine the consolidation of skills after more than 12-hour in healthy aging.

The present study goes beyond Nemeth et al (2010) and other previous studies (Press et al., 2005; Siengsukon & Boyd, 2009a; Spencer et al., 2007) by comparing the performance after 12-, 24-hour and 1 week delays from the initial learning session. We focused on the consolidation of implicit sequence-specific learning and general skill learning, separately, in healthy young and elderly adults. Exploring the time course of skill consolidation in different age groups can help us not only in better understanding memory and learning systems, but also in understanding age-related changes in brain plasticity, and thus developing more effective training and rehabilitation programs.

Methods

Participants

Seventy-one young and 58 elderly right-handed adults participated in the experiment. They were randomly assigned to the 12-hour, 24-hour or 1-week delay group (seventy percent of the 12-hour groups were also presented in the study of Nemeth et al. (2010)). The age and education of participants were controlled (for detailed data see Table 3). Participants did not suffer from any developmental, psychiatric or neurological disorders, did not have sleeping disorders, and all reported having 7-8 hours of sleep a day. Participants were recruited from Szeged and Bekescsaba (Hungary). They gave informed written consent after the aims and procedures of the experiment were explained to them, and received no financial compensation for participation.

Group		Age	Education	Gender
Young	12 hrs (n=23)	20.83 (1.11)	13.48 (1.27)	8 M / 15 F
	24 hrs (n=31)	21.74 (4.15)	14.77 (2.13)	12 M / 19 F
	1 week (n=17)	19.88 (1.27)	13.65 (1.32)	3 M / 14 F
Elderly	12 hrs (n=23)	66.43 (6.18)	12.90 (3.26)	7 M / 14 F
	24 hrs (n=22)	67.36 (5.30)	13.84 (2.44)	5 M / 17 F
	1 week (n=13)	65.15 (4.14)	14.23 (2.00)	6 M / 7 F

Table 3: Mean age, mean education in years and gender (female – F, male – M) of young and elderly groups. In all columns, numbers in parentheses represent standard deviation.

Procedure

There were two sessions in the experiment to examine the offline changes of implicit skill learning: a learning phase (Session 1) and a testing phase (Session 2) separated by a 12-, 24-hour or 1-week interval offline period (see Figure 7). Previous studies with similar tasks and experimental designs showed no time of day effect either on general reaction times or on learning measures (Nemeth, Janacsek, Londe et al., 2010; Press et al., 2005; Robertson et al., 2004; Song et al., 2007), the time of testing was however controlled in our study.

Figure 7: The design of the experiment. There were two sessions in the experiment: a learning and a testing phase separated either by 12-, 24-hour or 1-week.

Task

We used the same ASRT task (J. H. Howard, Jr. & Howard, 1997) as in Experiment I and Experiment II. During Session 1 (learning phase), the ASRT task consisted of 25 blocks, while Session 2 (testing phase) consisted of only 5 stimulus blocks of the same type as in Session 1. The repeating sequence was identical between Session 1 and Session 2 for each participant. To explore how much explicit knowledge subjects acquired about the task, we administered a short questionnaire as in the previous experiments. None of the participants reported noticing the repeating sequence of the stimulus locations.

Statistical analysis

To facilitate data processing, stimulus blocks were organized into larger clusters (called *epochs*); where the first epoch contained blocks 1-5, the second epoch blocks 6-10, etc. (Nemeth, Janacek, Londe et al., 2010; Song et al., 2007). Consequently, Session 1 consisted of 5 epochs, whereas Session 2 consisted of 1 epoch.

The accuracy of responses remained very high throughout the test (average over 97% for all groups), resulting a ceiling effect. Therefore, we analyzed the median reaction time (RT) for correct responses only, calculated separately for high- and low-frequency triplets and for each epoch.

Results

Online learning

To be able to investigate the offline changes, the learning in Session 1 must be similar in the groups. From this point of view the end of Session 1 is crucial (Nemeth, Janacsek, Londe et al., 2010; Press et al., 2005; Song et al., 2007). Therefore, sequence-specific learning (SSL) score was computed for RTs on low minus high frequency triplets in Epoch 5 (Song et al., 2007; Song, Howard, & Howard, 2008). The larger this difference, the greater the sequence-specific learning was. This score was submitted to a Univariate ANOVA with AGE GROUP (young vs. older) and DELAY (12, 24 hours or 1 week) as between-subject factors. The ANOVA revealed significant sequence-specific learning ($F(1,123)=136.69$, $\eta_p^2=0.53$, $p<0.001$). Neither the main effects of AGE GROUP and DELAY, nor the AGE GROUP x DELAY interaction was significant ($p's>0.19$), thus there were no differences in sequence-specific learning among age groups or delays at the end of Session 1 (Figure 8AB).

Regarding the general skill, a Univariate ANOVA was conducted on overall RT's of Epoch 5 (collapsed across triplet types) with AGE GROUP (young vs. older) and DELAY (12, 24 hours or 1 week) as between-subject factors. The only significant effect was the main effect of AGE GROUP ($F(1,123)=409.89$, $\eta_p^2=0.77$, $p<0.001$), reflecting longer RT's for the elderly group (537 ms) compared with the younger participants (381 ms). The main effect of DELAY and AGE GROUP x DELAY interaction did not approach significance (all $p's>0.41$). Thus, participants responded with similar overall reaction times in all delay conditions (Figure 8CD).

Figure 8: Sequence-specific learning (SSL; measured by RT on Low- minus High-frequency triplets) at the end of Session 1 and at the beginning of Session 2 for young (A) and elderly adults (B). All groups showed significant SSL by the end of Session 1. Overall RT's are also plotted for young (C) and elderly groups (D). Young participants were generally faster than the old ones. Regarding the offline SSL (measured by the difference of the SSL scores at the end of Session 1 and the beginning of Session 2), young groups retained the previously acquired sequence-specific skill, while the elderly groups showed decrement in this skill. There were no differences across the 12, 24 hours and 1 week conditions (E). Offline general skill learning (measured by the overall RT changes between sessions) were obtained in all young groups and decreased gradually across delay conditions. Older adults showed offline improvement only after a 12-hour interval (F). These groups responded significantly faster at the beginning of Session 2 compared to the end of Session 1, whereas elderly participants did not speed up after 24 hours or 1 week delay. Error bars indicate SEM.

Offline sequence-specific learning

To define the index of offline sequence-specific learning, we calculated the SSL learning score at the first epoch of Session 2 similarly to the learning score in Session 1 (Epoch 5). The greater this score at the beginning of the Session 2 compared to the end of Session 1, the larger the offline sequence-specific learning was. These two SSL learning scores were submitted to a mixed-design ANOVA with SESSION (end of Session 1 vs. beginning of Session 2) as within-subject factor, and AGE GROUP (young vs. elder) and DELAY (12, 24 hours or 1 week) as between-subject factors. Thus, any offline change of sequence-specific learning would be revealed by main effects and/or interactions with SESSION.

The main effect of SESSION was not significant ($F(1,123)=2.28$, $\eta_p^2=0.02$, $p=0.13$). However, the ANOVA revealed significant main effect of AGE GROUP ($F(1,123)=15.01$, $\eta_p^2=0.11$, $p<0.001$) and significant SESSION $\times$ AGE GROUP interaction ($F(1,123)=5.35$, $\eta_p^2=0.04$, $p=0.02$), reflecting that age groups differed in offline sequence-specific learning. Sequence-specific knowledge decreased in the elderly groups (-6 ms) compared to the young, who retained the previously acquired skill (1 ms). There was no offline improvement of sequence-specific knowledge in either group.

Regarding the time course, there were no differences among delay conditions (neither the main effect, nor the interactions with DELAY was significant, p 's >0.38), thus the consolidation of sequence-specific knowledge was not affected by the elapsed time between the learning and the testing session (Figure 8E).

Offline general skill learning

Offline general skill learning was tested by comparing the overall RT's (collapsed across triplet types) between the last epoch of Session 1 and the first epoch of Session 2 (Figure 8F). The greater the decrease from Session 1 to Session 2, the larger the offline general skill learning was. These two variables were submitted to a mixed-design ANOVA with SESSION (end of Session 1 vs. beginning of Session 2) as within-subject factor, and AGE GROUP (young vs. older) and DELAY (12, 24 hours or 1 week) as between-subject factors. Similarly to previous analysis, any offline change of general skill would be revealed by main effects and/or interactions with SESSION.

This ANOVA revealed significant offline general skill improvement (indicated by the main effect of SESSION: $F(1,123)=27.88$, $\eta_p^2=0.19$, $p<0.001$), participants responding faster

at the beginning of Session 2 than at the end of Session 1. This offline improvement was larger for the young group than for the elderly participants (shown by the SESSION x AGE GROUP interaction: $F(1,123)=11.45$, $\eta_p^2=0.09$, $p=0.001$). In addition, the elapsed time between the two sessions influenced the improvement of general skill improvements as well (indicated by the SESSION x DELAY interaction: $F(2,123)=5.29$, $\eta_p^2=0.08$, $p=0.006$). Thus, participants' response speed improved more after the 12-hour than after the 24-hour (LSD post hoc test: $p=0.007$) or 1-week delay ($p=0.006$), whereas there was no difference between the 24-hour and 1-week delay conditions ($p=0.64$).

The subsequent post hoc tests conducted separately for all age and delay groups revealed that the offline improvement of general skill was significant in all young groups ($p's < 0.047$), while in the elderly groups only the 12-hour delay led to offline enhancement ($p=0.032$). We also compared the performance of young and elderly groups in all conditions separately. Young adults showed larger offline improvement than the elderly participants in the 12-hour ($p=0.019$) as well as in the 24-hour delay period ($p=0.006$), while there was no significant difference between groups in the 1-week condition ($p=0.44$).

Discussion of Experiment III

We studied the time course of implicit skill consolidation in young and elderly adults with probabilistic second-order regularity sequences (the ASRT task), which enabled us to separate general skill and sequence-specific learning. In the young adults, we found offline improvement of the general skill (overall RT) after the 12-, 24-hour and 1-week delay as well, with gradual decline among delays. The elderly adults showed offline improvement of general skill only after the 12-hour offline period and this improvement was weaker than that in the young group. Although the pattern in age groups is similar, these results suggest that the offline course of general skill learning may be affected by aging, since we did not find improvement either after 24-hour or 1-week delay in the elderly group. No offline improvement was found in sequence-specific learning in either age group with either the 12-, 24-hour or 1-week consolidation interval. Sequence-specific learning did not decrease significantly between sessions for young participants suggesting that sequence-specific knowledge was well consolidated in this group, whereas the older group showed weaker consolidation in all delay conditions compared to the young. So, according our results offline general skill learning is influenced both by the time course and aging, while the offline sequence learning is affected only by aging.

The significant offline general skill improvement after the 12-hour delay period is compatible with the results of Song et al. (2007) and Nemeth et al. (2010). It is also possible that the improvement in overall RTs after the delay period reflects a release from fatigue rather than consolidation *per se*. However, studies that have included a fatigue control group (Spencer, Sunm, & Ivry, 2006; Walker et al., 2002) make this interpretation unlikely. The current results confirm those of Song et al. (2007) and Nemeth et al. (2010) in finding no evidence of any offline improvement of sequence-specific performance and extend them to the 24-hour and 1 week consolidation period.

The offline general skill improvement after 12 hours in the elderly adults contrasts the findings of both Spencer et al. (2007) and Siengsukon and Boyd (2009b) who obtained no offline improvement in elderly adults. Due to the different paradigms used in the current and the two previous studies, it is difficult to identify the source of differences. We refer to the point that these previous studies did not distinguish between general skill and sequence-specific learning, and therefore the signal-to-noise ratio might be reduced making it difficult to detect the offline improvements for elderly participants.

The differences among the 12-, 24-hour and 1-week offline intervals suggest that the consolidation of general skill learning is time-dependent. In addition, older participants are more sensitive for this offline time course in that they showed no offline improvement even after 24-hour delay. These results are congruent with recent theories of motor skill consolidation (Press et al., 2005; Robertson et al., 2005; Shadmehr & Brashers-Krug, 1997; Walker, Brakefield, Seidman et al., 2003) that claim that memory stabilization occurs during the first 5-6 hours after learning. The observed strong offline improvement after 12 hours may reflect this first stabilization process of memory traces, including the previously mentioned critical time period. The differences among 12-, 24-hour and 1-week consolidation intervals suggest that during new skill acquisition it could be important to place the training sessions closer to each other for optimal performance, with shorter intervals for elderly participants.

The current results as well as previous findings (Doyon, Korman et al., 2009; Robertson et al., 2004) are compatible with the notion that skill consolidation processes may be different and they could be profoundly dependent on the nature of task demand, such as on the relative proportion of general skill and sequence-specific learning requirements of the task. Given that these different components of learning are usually not separated in classical sequence learning tasks, offline improvements in such studies could be falsely attributed to sequence-specific learning itself. Nevertheless, in the current study consolidation of sequence-specific information was similar in the 12-, 24-hour and 1-week offline periods, with a decline

for elderly compared to the young, independently of time course. These results suggest that stabilization of sequence-specific memory is a faster process, whereas offline changes of general skill are more influenced by a longer stretch of time.

Interpreting our results in the framework proposed in Experiment I, there are at least three mechanisms which may underlie the age-related decline in the consolidation of skill learning on the functional level: 1) reduced detection of probabilities, 2) rigidity of internal models and/or 3) more restricted connections between internal models and probability detection (cf. Figure 4c). As the initial level of learning performance was matched between young and elderly adults, the decrease in sequence knowledge after the consolidation period might be attributed to the rigidity of internal models or to the weaker connections than to the reduced probability detection itself. Thus, based on these consolidation results, we suggest that not only the model-free, but also the model-based learning is limited in older ages. On the neuronal level age-related decrement was observed both structurally and functionally in the basal ganglia (Chen et al., 2005; Dennis & Cabeza, 2011; Erixon-Lindroth et al., 2005; N. Raz et al., 2005), that is involved in skill learning. A recent neuroimaging study (Dennis & Cabeza, 2011) found that elderly participants recruited medial temporal lobe structures in implicit sequence learning tasks, suggesting that compensation mechanisms are present to perform to an optimal level in these tasks. However, future studies are still needed to systematically examine the background mechanisms of age-related differences in skill consolidation.

General Discussion

Prediction is one of the most fundamental functions of the brain. In order to recognize time-based patterns and predict subsequent events, storing and recalling of sequences are required (Hawkins et al., 2009; Janacsek & Nemeth, 2012). Most predictions are based on the implicit learning that occurs when information is acquired from an environment of complex stimuli, without conscious access either to what was learned or to the fact that learning occurred (Cleeremans et al., 1998; Reber, 1993). Despite the growing interest in implicit learning in the past decades, there has been relatively little research on life-long development of implicit learning and on offline processing of implicitly learned information (i.e., consolidation). Here, we presented three studies investigating these issues.

As implicit sequence learning is highly dependent on the integrity of the striatal network, and age-related structural and biochemical changes are pronounced in the striatal complex and connected prefrontal areas (Dennis & Cabeza, 2011; N. Raz et al., 2005;

Rieckmann & Bäckman, 2009), it is an important question how online and offline sequence learning is affected by development and aging. Therefore, in **Experiment I**, we compared implicit sequence learning from 4 to 85 years of age to determine age-related differences across human life span and found the highest sequence learning before adolescence measured by raw RT. This result can be explained by a competition between two fundamentally different forms of learning (model-free and model-based; based on Daw et al., 2005) suggesting that after adolescence frontal lobe-mediated model-based learning has larger effect on the expression of implicit sequence knowledge, while before adolescence basal ganglia-dependent model-free learning is more influencing.

Since a growing body of data has shown that frontal lobe-mediated processes are disrupted in hypnosis (Fingelkurts et al., 2007; Oakley & Halligan, 2009), we tested the previous assumption in **Experiment II** by comparing implicit sequence learning in hypnosis and in waking alert state. We found that hypnosis boosted sequence learning providing support for the idea that learning and memory processes may not only involve the engagement of specific neuroplastic mechanisms, but may also rely upon the disengagement of interacting systems (Brown & Robertson, 2007a, p. 149). Our finding is in line with previous studies demonstrating that manipulations reducing the reliance on frontal lobe-dependent processes improved basal ganglia-dependent learning performance (e.g., Filoteo et al., 2010; Galea et al., 2010). This interpretation is consistent with the result that participants with better frontal lobe related executive functions showed decreased sequence learning in the waking alert condition compared to the participants with lower executive functions. By contrast, in the hypnotic state, participants with higher executive functions shifted from relying on frontal lobe-related attentional processes to automatic, procedural-based mechanisms, resulting in enhanced sequence learning.

Finally, in **Experiment III** we investigated consolidation of implicit sequence knowledge by comparing the performance after 12-, 24-hour, and 1-week delays from the initial learning session in young and elderly adults in order to determine age-related differences not only in online learning, but also in offline processing of the learned material. This research went beyond previous ones (e.g., Robertson et al., 2005) (Shadmehr & Brashers-Krug, 1997; Shadmehr & Holcomb, 1997; Walker, Brakefield, Hobson et al., 2003) in that: 1) it used the ASRT task, which allowed to investigate sequence-specific and general skill learning separately, and 2) it compared three consolidation intervals to explore the time-dependent offline changes in more detail. In the young adults, we found offline improvement of the general skill after all delays, with gradual decline among them. The elderly adults

showed offline improvement of the general skill only after the 12-hour offline period, and this improvement was weaker than that in the young group. Although the pattern in age groups was similar, these results suggest that the offline course of general skill learning may be affected by aging, since we did not find improvement either after a 24-hour or 1-week delay in the elderly group. No offline improvement was found in sequence-specific learning in either age group with any of the consolidation intervals. Sequence-specific learning did not decrease significantly between sessions for young participants, suggesting that sequence-specific knowledge was well consolidated in this group, whereas the older group showed weaker consolidation in all delay conditions compared to the younger group. Thus, according to these results, offline general skill learning is influenced both by the time course and aging, while the offline sequence learning is affected only by aging.

Conclusions

The following conclusions can be drawn based on the studies presented in this thesis:

1. We found dissociation between accuracy and RT learning measures in Experiment I.
2. Accuracy learning measures showed a typical bell-shaped curve across life span. This measure of implicit sequence learning may be connected to attentional resources to a higher extent. Consequently, children and elderly people who have underdeveloped / deteriorating attentional functions compared to young adults, show a lower performance.
3. Based on raw RT results we suggest that acquiring fundamentally new skills implicitly (i.e., relying less on attentional resources) is most effective before adolescence. The reduced learning after adolescence indicates a shift to relying more on internal model-based interpretations (connected to PFC/MTL structures), while somewhat neglecting the raw probabilities of the sensory input (mediated primarily by the BG).
4. In Experiment II we have shown that hypnosis boosted sequence learning in young adults, providing support for the idea that manipulations which reduce the reliance on frontal lobe-dependent processes improve BG-dependent learning performance.
5. Executive functions moderated the boosting effect of hypnosis; it was more pronounced in participants with higher executive functions.
6. Regarding the consolidation of implicit knowledge, offline improvement of the general skill was found after 12-, 24-hour and 1-week delay in young adults, with a gradual decline among them. The elderly adults showed offline improvement of the

general skill only after the 12-hour offline period and this improvement was weaker than that in the young group.

7. No offline improvement was found in sequence-specific learning in either age group with any of the consolidation intervals. Sequence-specific learning did not decrease significantly between sessions for young participants, suggesting that sequence-specific knowledge was well consolidated in this group, whereas the older group showed weaker consolidation in all delay conditions compared to the younger group.
8. To sum up, our findings in Experiment III draw attention to the fact that the consolidation is not a single process; instead there are multiple mechanisms in offline learning (general skill, sequence-specific processes), which are differently influenced by aging and the course of time.

Our results contribute not only to the better understanding of learning and memory systems on a behavioral level, but also to understanding the age-related changes in brain plasticity in healthy participants across the human life span. In addition, these findings can help better understand neurodevelopmental (e.g., autism, specific language impairment, dyslexia), neurodegenerative (e.g., Parkinson's Disease, Huntington's Disease) and age-related disorders (e.g., dementia) where related brain structures (e.g., PFC, MTL, BG) are affected. Finally, our findings can lead to the development of more effective diagnostic tools, training methods and rehabilitation programs.

Acknowledgements

I would like to express my sincere gratitude to my supervisor Dezső Németh for his support throughout the years of my scientific work. I am also thankful to Attila Krajcsi, Darlene V. Howard, James H. Howard, Jr., Michael T. Ullman, and Gábor Orosz for numerous illuminating discussions. Their vast knowledge and attitude towards science have been extremely valuable for me. Thanks to my co-authors József Fiser, Zoltan Ambrus Kovács, Bertalan Polner, Emese Hallgató, Dóra Győri-Dani, Judit Pekár, Eszter Csábi, Mária Várszegi, Zoltán Mari, Gábor Csifcsák, Gábor Szvoboda, Virág Balogh, Márta Fazekas, Izabella Dányi, Ágnes Vetró, Szilvia Jámbori, Robert Mingesz, Tímea Sándor, and Zsuzsa Londe for fruitful collaborations. Thanks also to István Winkler, Ágnes Lukács, and Ágnes Kovács for their comments on the previous versions of the published manuscripts discussed in my PhD thesis. Thanks to BA and MA students in Szeged for their valuable assistance during data collection.

References

- Aarts, E., & Roelofs, A. (2011). Attentional control in anterior cingulate cortex based on probabilistic cueing. *Journal of cognitive neuroscience*, 23(3), 716-727.
- Albouy, G., Sterpenich, V., Balteau, E., Vandewalle, G., Desseilles, M., Dang-Vu, T., et al. (2008). Both the hippocampus and striatum are involved in consolidation of motor sequence memory. *Neuron*, 58(2), 261-272.
- Anokhin, A. P., Golosheykin, S., Grant, J. D., & Heath, A. C. (2010). Developmental and genetic influences on prefrontal function in adolescents: A longitudinal twin study of WCST performance. *Neuroscience Letters*, 472(2), 119-122.
- Ashby, F. G., Alfonso-Reese, L. A., Turken, A. U., & Waldron, E. M. (1998). A neuropsychological theory of multiple systems in category learning. *Psychological Review*, 105, 442-481.
- Aslin, R. N., Saffran, J. R., & Newport, E. L. (1998). Computation of conditional probability statistics by 8-month old infants. *Psychological Science*, 9(4), 321-324.
- Baldo, J. V., Schwartz, S., Wilkins, D., & Dronkers, N. F. (2006). Role of frontal versus temporal cortex in verbal fluency as revealed by voxel-based lesion symptom mapping. *Journal of the International Neuropsychological Society*, 12(06), 896-900.
- Barnes, K. A., Howard, J. H., Jr., Howard, D. V., Gilotty, L., Kenworthy, L., Gaillard, W. D., et al. (2008). Intact implicit learning of spatial context and temporal sequences in childhood autism spectrum disorder. *Neuropsychology*, 22(5), 563-570.
- Barnes, K. A., Howard, J. H., Jr., Howard, D. V., Kenealy, L., & Vaidya, C. J. (2010). Two forms of implicit learning in childhood ADHD. *Developmental Neuropsychology*, 35(5), 494-505.
- Bennett, I. J., Howard, J. H., Jr., & Howard, D. V. (2007). Age-Related Differences in Implicit Learning of Subtle Third-Order Sequential Structure. *Journal of Gerontology: Psychological Sciences*, 62B(2), 98-103.
- Bennett, I. J., Madden, D. J., Vaidya, C. J., Howard, J. H., Jr., & Howard, D. V. (2011). White matter integrity correlates of implicit sequence learning in healthy aging. *Neurobiology of Aging*, 32(12), 2317.e2301-e2312.
- Bird, C. M., & Burgess, N. (2008). The hippocampus and memory: insights from spatial processing. *Nature Reviews Neuroscience*, 9(3), 182-194.

- Blakemore, S. J., & Choudhury, S. (2006). Development of the adolescent brain: implications for executive function and social cognition. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, 47(3-4), 296-312.
- Brown, R. M., & Robertson, E. M. (2007a). Inducing motor skill improvements with a declarative task. *Nature Neuroscience*, 10(2), 148-149.
- Brown, R. M., & Robertson, E. M. (2007b). Off-line processing: Reciprocal interactions between declarative and procedural memories. *Journal of Neuroscience*, 27(39), 10468-10475.
- Burgess, P. W., Gilbert, S. J., & Dumontheil, I. (2007). Function and localization within rostral prefrontal cortex (area 10). *Philosophical Transactions of the Royal Society B: Biological Sciences*, 362(1481), 887-899.
- Chen, P. S., Yang, Y. K., Lee, Y. S., Yeh, T. L., Lee, I. H., Chiu, N. T., et al. (2005). Correlation between different memory systems and striatal dopamine D2/D3 receptor density: a single photon emission computed tomography study. *Psychol Med*, 35(2), 197-204.
- Christ, S. E., White, D. A., Mandernach, T., & Keys, B. A. (2001). Inhibitory control across the life span. *Developmental Neuropsychology*, 20(3), 653-669.
- Cleeremans, A., Destrebecqz, A., & Boyer, M. (1998). Implicit learning: news from the front. *Trends in Cognitive Sciences*, 2(10), 406-416.
- Clohessy, A. B., Posner, M. I., & Rothbart, M. K. (2001). Development of the functional visual field. *Acta Psychologica*, 106(1), 51-68.
- Cohen, D., Pascual-Leone, A., Press, D., & Robertson, E. (2005). Off-line learning of motor skill memory: A double dissociation of goal and movement. *Proceedings of the National Academy of Sciences*, 102(50), 18237-18241.
- Craik, F. I., & Bialystok, E. (2006). Cognition through the lifespan: mechanisms of change. *TRENDS in Cognitive Sciences*, 10(3), 131-138.
- Curran, T. (1997). Effects of aging on implicit sequence learning: accounting for sequence structure and explicit knowledge. *Psychological Research*, 60(1-2), 24-41.
- Daw, N. D., Niv, Y., & Dayan, P. (2005). Uncertainty-based competition between prefrontal and dorsolateral striatal systems for behavioral control. *Nature Neuroscience*, 8(12), 1704-1711.
- Dennis, N., & Cabeza, R. (2011). Age-related dedifferentiation of learning systems: an fMRI study of implicit and explicit learning. *Neurobiology of Aging*, 32(12), 2318.e2317-e2330. doi: 10.1016/j.neurobiolaging.2010.04.004

- Der, G., & Deary, I. J. (2006). Age and sex differences in reaction time in adulthood: Results from the United Kingdom Health and Lifestyle Survey. *Psychology and Aging, 21*(1), 62-73.
- Doyon, J., Bellec, P., Amsel, R., Penhune, V., Monchi, O., Carrier, J., et al. (2009). Contributions of the basal ganglia and functionally related brain structures to motor learning. *Behavioral Brain Research, 199*(1), 61-75.
- Doyon, J., Korman, M., Morin, A., Dostie, V., Hadj Tahar, A., Benali, H., et al. (2009). Contribution of night and day sleep vs. simple passage of time to the consolidation of motor sequence and visuomotor adaptation learning. *Exp Brain Res, 195*(1), 15-26.
- Doyon, J., Song, A. W., Karni, A., Lalonde, F., Adams, M. M., & Ungerleider, L. G. (2002). Experience-dependent changes in cerebellar contributions to motor sequence learning. *Proc Natl Acad Sci U S A, 99*(2), 1017-1022.
- Dum, R. P., & Strick, P. L. (1993). Cingulate motor areas. In B. A. Vogt & M. Gabriel (Eds.), *Neurobiology of cingulate cortex and limbic thalamus: A comprehensive handbook* (pp. 415-441). Boston: Birkhauser.
- Egner, T., Jamieson, G., & Gruzelier, J. (2005). Hypnosis decouples cognitive control from conflict monitoring processes of the frontal lobe. *Neuroimage, 27*(4), 969-978.
- Erixon-Lindroth, N., Farde, L., Wahlin, T. B., Sovago, J., Halldin, C., & Backman, L. (2005). The role of the striatal dopamine transporter in cognitive aging. *Psychiatry Res, 138*(1), 1-12.
- Feeney, J. J., Howard, J. H., & Howard, D. V. (2002). Implicit learning of higher order sequences in middle age. *Psychology and Aging, 17*(2), 351-355.
- Filoteo, J. V., Lauritzen, S., & Maddox, W. T. (2010). Removing the Frontal Lobes. *Psychological Science, 21*(3), 415-423.
- Fingelkurts, A. A., Kallio, S., & Revonsuo, A. (2007). Cortex functional connectivity as a neurophysiological correlate of hypnosis: An EEG case study. *Neuropsychologia, 45*(7), 1452-1462.
- Fischer, S., Hallschmid, M., Elsner, A. L., & Born, J. (2002). Sleep forms memory for finger skills. *PNAS, 99*(18), 11987-11991.
- Fiser, J., & Aslin, R. N. (2001). Unsupervised statistical learning of higher-order spatial structures from visual scenes. *Psychological Science, 12*(6), 499-504.
- Fiser, J., & Aslin, R. N. (2002). Statistical learning of new visual feature combinations by infants. *Proceedings of the National Academy of Sciences, 99*(24), 15822-15826. doi: 10.1073/pnas.232472899

- Fletcher, J., Maybery, M. T., & Bennett, S. (2000). Implicit learning differences: A question of developmental level? *Journal of Experimental Psychology: Learning, Memory, and Cognition*, 26(1), 246-252.
- Fletcher, P. C., Zafiris, O., Frith, C. D., Honey, R. A. E., Corlett, P. R., Zilles, K., et al. (2005). On the benefits of not trying: brain activity and connectivity reflecting the interactions of explicit and implicit sequence learning. *Cerebral Cortex*, 15(7), 1002-1015.
- Foerde, K., Knowlton, B. J., & Poldrack, R. A. (2006). Modulation of competing memory systems by distraction. *Proceedings of the National Academy of Science*, 103(31), 11778-11783.
- Frank, M. J., O'Reilly, R. C., & Curran, T. (2006). When Memory Fails, Intuition Reigns: Midazolam Enhances Implicit Inference in Humans. *Psychological Science*, 17(8), 700 - 707.
- Frensch, P. A. (1998). One concept, multiple meanings. In M. A. Stadler & P. A. Frensch (Eds.), *Handbook of implicit learning* (pp. 47-104). London: SAGE Publications.
- Frensch, P. A., & Miner, C. S. (1994). Effects of presentation rate and individual differences in short-term memory capacity on an indirect measure of serial learning. *Memory and Cognition*, 22(1), 95-110.
- Fu, W.-T., & Anderson, J. R. (2008). Solving the credit assignment problem: Explicit and implicit learning of action sequences with probabilistic outcomes. *Psychological Research/Psychologische Forschung*, 72(3), 321.
- Gaillard, V., Destrebecqz, A., Michiels, S., & Cleeremans, A. (2009). Effects of age and practice in sequence learning: A graded account of ageing, learning, and control. *European Journal of Cognitive Psychology*, 21(2), 255-282.
- Galea, J. M., Albert, N. B., Ditye, T., & Miall, R. C. (2010). Disruption of the dorsolateral prefrontal cortex facilitates the consolidation of procedural skills. *Journal of cognitive neuroscience*, 22(6), 1158-1164.
- Giedd, J. N. (2008). The teen brain: insights from neuroimaging. *Journal of Adolescent Health*, 42(4), 335-343.
- Giedd, J. N., Blumenthal, J., Jeffries, N. O., Castellanos, F. X., Liu, H., Zijdenbos, A., et al. (1999). Brain development during childhood and adolescence: A longitudinal MRI study. *Nature Neuroscience*, 2(10), 861-863.

- Giorgio, A., Santelli, L., Tomassini, V., Bosnell, R., Smith, S., De Stefano, N., et al. (2010). Age-related changes in grey and white matter structure throughout adulthood. *Neuroimage*, 51(3), 943-951.
- Gomez, R. L., & Gerken, L. (1999). Artificial grammar learning by 1-year-olds leads to specific and abstract knowledge. *Cognition*, 70(2), 109-135.
- Gruzelier, J. H. (2006). Frontal functions, connectivity and neural efficiency underpinning hypnosis and hypnotic susceptibility. *Contemporary Hypnosis*, 23(1), 15-32.
- Hallgató, E., Györi-Dani, D., Pekár, J., Janacsek, K., & Nemeth, D. (in press). The differential consolidation of perceptual and motor learning in skill acquisition. *Cortex*. doi: 10.1016/j.cortex.2012.01.002
- Halsband, U. (2006). Learning in trance: Functional brain imaging studies and neuropsychology. *Journal of Physiology-Paris*, 99(4-6), 470-482.
- Hawkins, J., George, D., & Niemasik, J. (2009). Sequence memory for prediction, inference and behaviour. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 364(1521), 1203-1209.
- Heaton, R., Chelune, G., Talley, J., Kay, G., & Curtiss, G. (1993). Wisconsin card sorting test manual: PAR.
- Heerey, E. A., & Velani, H. (2010). Implicit learning of social predictions. *Journal of Experimental Social Psychology*, 46(3), 577-581.
- Henke, K. (2010). A model for memory systems based on processing modes rather than consciousness. *Nature Reviews Neuroscience*, 11, 523-532.
- Hikosaka, O., Nakahara, H., Rand, M. K., Sakai, K., Lu, X., Nakamura, K., et al. (1999). Parallel neural networks for learning sequential procedures. *Trends in Neurosciences*, 22(10), 464-471.
- Hikosaka, O., Nakamura, K., Sakai, K., & Nakahara, H. (2002). Central mechanisms of motor skill learning. *Current Opinion in Neurobiology*, 12(2), 217-222.
- Howard, D. V., & Howard, J. H. (1992). Adult age differences in the rate of learning serial patterns: Evidence from direct and indirect tests. *Psychology and Aging*, 7(2), 232-241.
- Howard, D. V., & Howard, J. H. (2001). When it does hurt to try: Adult age differences in the effects of instructions on implicit pattern learning. *Psychonomic Bulletin and Review*, 8(4), 798-805.
- Howard, D. V., & Howard, J. H., Jr. (1989). Age differences in learning serial patterns: direct versus indirect measures. *Psychology and Aging*, 4(3), 357-364.

- Howard, D. V., Howard, J. H., Jr., Japikse, K., DiYanni, C., Thompson, A., & Somberg, R. (2004). Implicit sequence learning: effects of level of structure, adult age, and extended practice. *Psychology and Aging, 19*(1), 79-92.
- Howard, J. H., Jr., & Howard, D. V. (1997). Age differences in implicit learning of higher-order dependencies in serial patterns. *Psychology and Aging, 12*(4), 634-656.
- Hultsch, D. F., MacDonald, S. W. S., Hunter, M. A., Levy-Bencheton, J., & Strauss, E. (2000). Intraindividual variability in cognitive performance in older adults: Comparison of adults with mild dementia, adults with arthritis, and healthy adults. *Neuropsychology, 14*(4), 588-598.
- Hunt, R. H., & Aslin, R. N. (2001). Statistical learning in a serial reaction time task: access to separable statistical cues by individual learners. *Journal of Experimental Psychology: General, 130*(4), 658-680.
- Janacsek, K., Fiser, J., & Nemeth, D. (in press). The best time to acquire new skills: age-related differences in implicit sequence learning. *Developmental Science*.
- Janacsek, K., & Nemeth, D. (2012). Predicting the future: From implicit learning to consolidation. *International Journal of Psychophysiology, 83*(2), 213-221. doi: 10.1016/j.bbr.2011.03.031
- Kaiser, J., Barker, R., Haenschel, C., Baldeweg, T., & Gruzelier, J. H. (1997). Hypnosis and event-related potential correlates of error processing in a stroop-type paradigm: a test of the frontal hypothesis* 1. *International Journal of Psychophysiology, 27*(3), 215-222.
- Kallio, S., Revonsuo, A., Hamalainen, H., Markela, J., & Gruzelier, J. (2001). Anterior brain functions and hypnosis: A test of the frontal hypothesis. *International Journal of Clinical and Experimental Hypnosis, 49*(2), 95-108.
- Karni, A., Meyer, G., Jezard, P., Adams, M. M., Turner, R., & Ungerleider, L. G. (1995). Functional MRI evidence for adult motor cortex plasticity during motor skill learning. *Nature, 377*, 155-158.
- Kaufman, S. B., DeYoung, C. G., Gray, J. R., Jiménez, L., Brown, J., & Mackintosh, N. (2010). Implicit learning as an ability. *Cognition, 116*(3), 321-340.
- Keele, S. W., Ivry, R., Mayr, U., Hazeltine, E., & Heuer, H. (2003). The cognitive and neural architecture of sequence representation. *Psychological Review, 110*(2), 316-339.
- Kincses, T., Johansen-Berg, H., Tomassini, V., Bosnell, R., Matthews, P., & Beckmann, C. (2008). Model-free characterization of brain functional networks for motor sequence learning using fMRI. *Neuroimage, 39*(4), 1950-1958.

- Kirkham, N., Slemmer, J., Richardson, D., & Johnson, S. (2007). Location, location, location: Development of spatiotemporal sequence learning in infancy. *Child Development*, 78(5), 1559-1571.
- Krakauer, J. W., & Shadmehr, R. (2006). Consolidation of motor memory. *Trends Neurosci*, 29(1), 58-64.
- Lieberman, M. D. (2000). Intuition: a social cognitive neuroscience approach. *Psychological Bulletin*, 126(1), 109-137.
- Marcus, G. F., Vijayan, S., Rao, S. B., & Vishton, P. M. (1999). Rule Learning by seven-month-old infants. *Science*, 283, 77-80.
- Maybery, M., Taylor, M., & O'Brien-Malone, A. (1995). Implicit learning: Sensitive to age but not IQ. *Australian Journal of Psychology*, 47(1), 8-17.
- Medford, N., & Critchley, H. D. (2010). Conjoint activity of anterior insular and anterior cingulate cortex: awareness and response. *Brain Structure and Function*, 214(5), 535-549.
- Meulemans, T., Van der Linden, M., & Perruchet, P. (1998). Implicit sequence learning in children. *Journal of Experimental Child Psychology*, 69(3), 199-221.
- Nemeth, D., Csábi, E., Janacsek, K., Várszegi, M., & Mari, Z. (in press). Intact implicit probabilistic sequence learning in Obstructive Sleep Apnea. *Sleep Medicine*.
- Nemeth, D., Hallgato, E., Janacsek, K., Sandor, T., & Londe, Z. (2009). Perceptual and motor factors of implicit skill learning. *Neuroreport*, 20(18), 1654-1658. doi: 10.1097/WNR.0b013e3283333ba08
- Nemeth, D., & Janacsek, K. (2011). The dynamics of implicit skill consolidation in young and elderly adults. *Journal of Gerontology Psychological Science*, 66(1), 15-22. doi: 10.1093/geronb/gbq063
- Nemeth, D., Janacsek, K., Balogh, V., Londe, Z., Mingesz, R., Fazekas, M., et al. (2010). Learning in Autism: Implicitly Superb. *PLoS ONE*, 5(7), e11731.
- Nemeth, D., Janacsek, K., Csifcsak, G., Szvoboda, G., Howard, J. H., Jr., & Howard, D. V. (2011). Interference between Sentence Processing and Probabilistic Implicit Sequence Learning. *PloS one*, 6(3), e17577.
- Nemeth, D., Janacsek, K., Londe, Z., Ullman, M. T., Howard, D., & Howard, J. (2010). Sleep has no critical role in implicit motor sequence learning in young and old adults. *Experimental Brain Research*, 201(2), 351-358. doi: 10.1007/s00221-009-2024-x
- Nissen, M. J., & Bullemer, P. (1987). Attentional requirements of learning: Evidence from performance measures. *Cognitive Psychology*, 19, 1-32.

- Oakley, D. A., & Halligan, P. W. (2009). Hypnotic suggestion and cognitive neuroscience. *Trends in Cognitive Sciences*, 13(6), 264-270.
- Orban, G., Fiser, J., Aslin, R. N., & Lengyel, M. (2008). Bayesian learning of visual chunks by human observers. *Proceedings of the National Academy of Sciences*, 105(7), 2745-2750.
- Poldrack, R. A., Clark, J., Pare-Blagoev, E. J., Shohamy, D., Creso Moyano, J., Myers, C., et al. (2001). Interactive memory systems in the human brain. *Nature*, 414(6863), 546-550.
- Poldrack, R. A., & Packard, M. G. (2003). Competition among multiple memory systems: converging evidence from animal and human brain studies. *Neuropsychologia*, 41(3), 245-251.
- Poldrack, R. A., Sabb, F. W., Foerde, K., Tom, S. M., Asarnow, R. F., Bookheimer, S. Y., et al. (2005). The Neural Correlates of Motor Skill Automaticity. *Journal of Neuroscience*, 25, 5356 - 5364.
- Press, D. Z., Casement, M. D., Pascual-Leone, A., & Robertson, E. M. (2005). The time course of off-line motor sequence learning. *Brain Res Cogn Brain Res*, 25(1), 375-378.
- Prinzmetal, W., McCool, C., & Park, S. (2005). Attention: Reaction time and accuracy reveal different mechanisms. *Journal of Experimental Psychology: General*, 134(1), 73-92.
- Rabbitt, P., Osman, P., Moore, B., & Stollery, B. (2001). There are stable individual differences in performance variability, both from moment to moment and from day to day. *The Quarterly Journal of Experimental Psychology Section A*, 54(4), 981-1003.
- Raz, A., & Shapiro, T. (2002). Hypnosis and neuroscience: a cross talk between clinical and cognitive research. *Archives of General Psychiatry*, 59(1), 85-90.
- Raz, N., Lindenberger, U., Rodrigue, K. M., Kennedy, K. M., Head, D., Williamson, A., et al. (2005). Regional brain changes in aging healthy adults: general trends, individual differences and modifiers. *Cereb Cortex*, 15(11), 1676-1689.
- Reber, A. R. (1993). *Implicit learning and tacit knowledge: An essay on the cognitive unconscious* (Vol. 19). New York: Oxford University Press.
- Remillard, G. (2008). Implicit learning of second-, third-, and fourth-order adjacent and nonadjacent sequential dependencies. *The Quarterly Journal of Experimental Psychology*, 61(3), 400-424.
- Rieckmann, A., & Bäckman, L. (2009). Implicit learning in aging: Extant patterns and new directions. *Neuropsychology review*, 19(4), 490-503.

- Rieckmann, A., Fischer, H., & Bäckman, L. (2010). Activation in striatum and medial temporal lobe during sequence learning in younger and older adults: Relations to performance. *Neuroimage*, 50(3), 1303-1312.
- Robertson, E. M. (2009). From creation to consolidation: A novel framework for memory processing. *PLoS Biology*, 7(1), e1000019.
- Robertson, E. M., Pascual-Leone, A., & Press, D. Z. (2004). Awareness modifies the skill-learning benefits of sleep. *Current Biology*, 14(3), 208-212.
- Robertson, E. M., Press, D. Z., & Pascual-Leone, A. (2005). Off-line learning and the primary motor cortex. *J Neurosci*, 25(27), 6372-6378.
- Roelofs, A., Van Turenout, M., & Coles, M. G. H. (2006). Anterior cingulate cortex activity can be independent of response conflict in Stroop-like tasks. *Proceedings of the National Academy of Sciences*, 103(37), 13884-13889.
- Romano Bergstrom, J. C., Howard, J. H., Jr., & Howard, D. V. (2011). Enhanced implicit sequence learning in collage-age video game players and musicians. *Applied Cognitive Psychology*. doi: 10.1002/acp.1800
- Romano, J. C., Howard, J. H., & Howard, D. V. (2010). One-year retention of general and sequence-specific skills in a probabilistic, serial reaction time task. *Memory*, 18(4), 427-441.
- Römer, S., Schulz, A., Richter, S., Lass-Hennemann, J., & Schächinger, H. (2011). Oral cortisol impairs implicit sequence learning. *Psychopharmacology*, 215, 33-40.
- Saffran, J. R., Aslin, R. N., & Newport, E. L. (1996). Statistical learning by 8-month-old infants. *Science*, 274, 1926-1928.
- Saffran, J. R., Johnson, E. K., Aslin, R. N., & Newport, E. L. (1999). Statistical learning of tone sequences by human infants and adults. *Cognition*, 70(1), 27-52. doi: S0010-0277(98)00075-4
- Salthouse, T. A. (1996). The processing-speed theory of adult age differences in cognition. *Psychological Review*, 103(3), 403-428.
- Sanger, T. D. (2010). Controlling variability. *Journal of Motor Behavior*, 42(6), 401-407.
- Schendan, H., Searl, M., Melrose, R., & Stern, C. (2003). An fMRI study of the role of the medial temporal lobe in implicit and explicit sequence learning. *Neuron*, 37(6), 1013-1025.
- Scoville, W. B., & Milner, B. (1957). Loss of recent memory after bilateral hippocampal lesions. *Journal of Neurology, Neurosurgery, and Psychiatry*, 20, 11-21.

- Sefcsik, T., Nemeth, D., Janacsek, K., Hoffmann, I., Scialabba, J., Klivenyi, P., et al. (2009). The role of the putamen in cognitive functions—A case study. *Learning & Perception*, 1(2), 215-227.
- Shadmehr, R., & Brashers-Krug, T. (1997). Functional stages in the formation of human long-term motor memory. *Journal of Neuroscience*, 17(1), 409-419.
- Shadmehr, R., & Holcomb, H. H. (1997). Neural correlates of motor memory consolidation. *Science*, 277(5327), 821-825.
- Shor, R. E., Orne, E. C., & Press, C. P. (1962). *Harvard group scale of hypnotic susceptibility*: Consulting Psychologists Press.
- Siengsukon, C. F., & Boyd, L. A. (2009a). Does Sleep Promote Motor Learning? Implications for Physical Rehabilitation. *Physical Therapy*, 89(4), 370.
- Siengsukon, C. F., & Boyd, L. A. (2009b). Sleep to learn after stroke: implicit and explicit off-line motor learning. *Neuroscience Letters*, 451(1), 1-5.
- Slifkin, A. B., & Newell, K. M. (1998). Is variability in human performance a reflection of system noise? *Current Directions in Psychological Science*, 7(6), 170-177.
- Slifkin, A. B., & Newell, K. M. (1999). Noise, information transmission, and force variability. *Journal of Experimental Psychology: Human Perception and Performance*, 25(3), 837.
- Soetens, E., Melis, A., & Notebaert, W. (2004). Sequence learning and sequential effects. *Psychological Research*, 69(1), 124-137.
- Song, S. (2009). Consciousness and the consolidation of motor learning. *Behavioural Brain Research*, 196(2), 180-186.
- Song, S., Howard, J. H., Jr., & Howard, D. V. (2007). Sleep does not benefit probabilistic motor sequence learning. *Journal of Neuroscience*, 27(46), 12475-12483.
- Song, S., Howard, J. H. J., & Howard, D. V. (2008). Perceptual sequence learning in a serial reaction time task. *Experimental Brain Research*, 189(2), 145-158.
- Spencer, R. M., Gouw, A. M., & Ivry, R. B. (2007). Age-related decline of sleep-dependent consolidation. *Learn Mem*, 14(7), 480-484.
- Spencer, R. M., Sunm, M., & Ivry, R. B. (2006). Sleep-Dependent Consolidation of Contextual Learning. *Current Biology*, 16, 1001-1005.
- Spreen, O., & Strauss, E. (1991). Language Tests *A Compendium of Neuropsychological Tests* (pp. 268-275).

- Squire, L. R., & Knowlton, B. J. (2000). The medial temporal lobe, the hippocampus, and the memory systems of the brain. In M. S. Gazzaniga (Ed.), *The new cognitive neurosciences* (pp. 765-780). Cambridge, MA: MIT Press.
- Squire, L. R., Stark, C. E., & Clark, R. E. (2004). The Medial Temporal Lobe. *Annual Review of Neuroscience*, 27, 279-306.
- Squire, L. R., & Zola-Morgan, S. (1991). The medial temporal lobe memory system. *Science*, 253, 1380-1386.
- Squire, L. R., & Zola, S. M. (1996). Structure and function of declarative and nondeclarative memory systems. *Proceedings of the National Academy of Sciences USA*, 93, 13515-13522.
- Stickgold, R., Fosse, R., & Walker, M. P. (2002). Linking brain and behavior in sleep-dependent learning and memory consolidation. *Proc Natl Acad Sci U S A*, 99(26), 16519-16521.
- Szendi, I., Kovacs, Z. A., Szekeres, G., Galsi, G., Boda, K., Boncz, I., et al. (2009). Effects of a hypnotically altered state of consciousness on intensification of semantic processing. *International Journal of Clinical and Experimental Hypnosis*, 57(4), 382-401.
- Tenenbaum, J. B., Kemp, C., Griffiths, T. L., & Goodman, N. D. (2011). How to grow a mind: Statistics, structure, and abstraction. *Science*, 331(6022), 1279-1285.
- Thomas, K. M., Hunt, R. H., Vizueta, N., Sommer, T., Durston, S., Yang, Y., et al. (2004). Evidence of Developmental Differences in Implicit Sequence Learning: An fMRI Study of Children and Adults. *Journal of Cognitive Neuroscience*, 16(8), 1339-1351.
- Tulving, E., & Craik, F. I. M. (Eds.). (2000). *The Oxford handbook of memory*. Oxford: Oxford University Press.
- Turner, E. C., & Brainard, M. S. (2007). Performance variability enables adaptive plasticity of crystallized adult birdsong. *Nature*, 450(7173), 1240-1244.
- Uddén, J., Folia, V., & Petersson, K. M. (2010). The Neuropharmacology of Implicit Learning. *Current Neuropharmacology*, 8(4), 367-381.
- Ullman, M. T. (2001). A neurocognitive perspective on language: The declarative/procedural model. *Nature Reviews Neuroscience*, 2, 717-726.
- Ullman, M. T. (2004). Contributions of memory circuits to language: The declarative/procedural model. *Cognition*, 92(1-2), 231-270.
- Vinter, A., & Perruchet, P. (2000). Implicit learning in children is not related to age: evidence from drawing behavior. *Child Development*, 71(5), 1223-1240.

- Wagstaff, G. F., Cole, J. C., & Brunas-Wagstaff, J. (2007). Effects of hypnotic induction and hypnotic depth on phonemic fluency: A test of the frontal inhibition account of hypnosis. *International Journal of Psychology and Psychological Therapy*, 7(1), 27-40.
- Walker, M. P., Brakefield, T., Hobson, J. A., & Stickgold, R. (2003). Dissociable stages of human memory consolidation and reconsolidation. *Nature*, 425(6958), 616-620. doi: 10.1038/nature01930
- Walker, M. P., Brakefield, T., Morgan, A., Hobson, J. A., & Stickgold, R. (2002). Practice with sleep makes perfect: sleep-dependent motor skill learning. *Neuron*, 35(1), 205-211.
- Walker, M. P., Brakefield, T., Seidman, J., Morgan, A., Hobson, J. A., & Stickgold, R. (2003). Sleep and the time course of motor skill learning. *Learn Mem*, 10(4), 275-284.
- Walker, M. P., & Stickgold, R. (2004). Sleep-dependent learning and memory consolidation. *Neuron*, 44(1), 121-133.
- Yap, M. J., Balota, D. A., Sibley, D. E., & Ratcliff, R. (2012). Individual differences in visual word recognition: Insights from the English lexicon project. *Journal of Experimental Psychology: Human Perception and Performance*, 38(1), 53-79.
- Zola-Morgan, S., Squire, L. R., & Amaral, D. G. (1986). Human Amnesia and the medial temporal region: Enduring memory impairment following a bilateral lesion limited to field CA1 of the hippocampus. *J Neurosci*, 6(10), 2950-2967.

Appendix

I. Janacsek, K., & Nemeth, D. (2012). Predicting the future: from implicit learning to consolidation. *International Journal of Psychophysiology*, 83(2), 213-221. DOI: 10.1016/j.ijpsycho.2011.11.012. Impact factor: 2.378.

II. Janacsek, K., Fiser J., & Nemeth, D. (in press). The best time to acquire new skills: Age-related differences in implicit sequence learning across life span. *Developmental Science*. DOI: 10.1111/j.1467-7687.2012.01150.x. Impact factor: 3.53.

III. Nemeth, D., Janacsek, K., Polner, B., & Kovacs, Z. A. (in press). Boosting human learning by hypnosis. *Cerebral Cortex*. DOI: 10.1093/cercor/bhs068. Impact factor: 6.844.

IV. Nemeth, D., & Janacsek, K. (2011). The dynamics of implicit skill consolidation in young and elderly adults. *Journal of Gerontology Series B: Psychological Sciences*, 66B(1), 15-22. DOI: 10.1093/geronb/gbq063. Impact factor: 2.094.

Appendix I

Janacsek, K., & Nemeth, D. (2012).

Predicting the future: from implicit learning to consolidation.

International Journal of Psychophysiology, 83(2), 213-221.

DOI: 10.1016/j.ijpsycho.2011.11.012.

Impact factor: 2.378.

Review

Predicting the future: From implicit learning to consolidation ☆

Karolina Janacsek, Dezso Nemeth *

Institute of Psychology, University of Szeged, Szeged, Hungary

ARTICLE INFO

Article history:

Received 14 July 2011

Received in revised form 30 October 2011

Accepted 17 November 2011

Available online 5 December 2011

Keywords:

Implicit sequence learning

Memory consolidation

Sleep

Offline changes

Implicit predictions

Aging

Awareness

Perceptual–motor learning

ABSTRACT

Sequence learning can be differentiated according to phases (rapid and slower), modalities (perceptual and motor), and whether or not it is conscious (implicit and explicit). Implicit sequence learning occurs when information is acquired from an environment of complex stimuli without conscious access either to what was learned or to the fact that learning occurred. In everyday life, this learning mechanism is crucial for adapting to the environment and for predicting events unconsciously. Implicit sequence learning underlies not only motor, but also cognitive and social skills; it is therefore an important aspect of life from infancy to old age. Moreover, this kind of learning does not occur only during practice, in the so-called online periods, but also between practice periods, during the so-called offline periods. The process that occurs during the offline periods is referred to as consolidation, which denotes the stabilization of a memory trace after the initial acquisition; this can result in increased resistance to interference or even improvement in performance following an offline period. Understanding the multiple aspects and influencing factors of consolidation can help us to reveal the nature of memory and changes in brain plasticity. Our review focuses on how consolidation varies with factors such as awareness, the length of offline periods, the type of information to be learned, and the age of participants. We highlight that consolidation is not a single process; instead, there are multiple mechanisms in the offline period, which are differently influenced by these factors.

© 2011 Elsevier B.V. All rights reserved.

1. Introduction

Prediction is one of the most fundamental functions of the brain. During every moment of our waking life, the brain is trying to anticipate future sensations. In order to recognize time-based patterns and predict subsequent events, storing and recalling of sequences are required (Hawkins et al., 2009). Without these skills, it would be impossible to carry out evolutionary adaptive behaviors. Most predictions are based on the implicit learning that occurs when information is acquired from an environment of complex stimuli, without conscious access either to what was learned or to the fact that learning occurred (Cleeremans et al., 1998; Reber, 1993). Despite the growing interest in implicit learning in the past decades, there has been relatively little research on offline processing of implicitly learned information (i.e., consolidation). Here, we review recent work on implicit sequence learning and its consolidation, with an emphasis on the last 10 years. More selectively, we address four of the most important factors that influence the consolidation of this fundamental learning mechanism. These factors must be taken into consideration before

planning and performing brain imaging, psychophysiological, and behavioral studies on sequence learning and its consolidation.

Implicit sequence learning underlies not only motor, but also cognitive and social skills (Kaufman et al., 2010; Lieberman, 2000; Nemeth et al., 2011; Romano Bergstrom et al., in press; Ullman, 2004); it is therefore an important aspect of life from infancy to old age. Implicit sequence learning is essential for learning languages, as well as learning to operate appliances, computer applications or musical instruments (Howard et al., 2004; Romano et al., 2010). Social skills appear in compound behaviors (including series of perceptions, emotions as well as motor actions) realized in proper sequences and under appropriate circumstances. These skills – for example, dialogs, decision making in social context, communication of emotions, predicting others' behavior based on previous verbal and nonverbal social communication, and adjusting our own behavior based on these predictions – are needed for normal social functioning in various sorts of situations: in the workplace, in the family, in the neighborhood, during recreation, shopping, or in the context of medical and mental care (Heerey and Velani, 2010; Lieberman, 2000; Nemeth and Janacsek, 2011). Furthermore, these skills are crucial for effective participation in educational, training, and rehabilitation programs, for instance in relearning how to walk, reach for objects, and speak after brain injury (Howard et al., 2004; Nemeth et al., 2010a).

Most models and empirical studies of skill learning highlight the role of the basal ganglia and the cerebellum (Dennis and Cabeza, 2011; Doyon et al., 2009a; Hikosaka et al., 1999; Hikosaka et al.,

☆ The authors report no conflict of interest and have no financial disclosure.

* Corresponding author at: Institute of Psychology, University of Szeged, Hungary, Egyetem u. 2., 6722, Szeged, Hungary. Tel.: +36 62 544 692, +36 62 544 509; fax: +36 62 544 509.

E-mail address: nemethd@edpsy.u-szeged.hu (D. Nemeth).

URL: <http://www.memory-and-language.com/> (D. Nemeth).

2002b; Keele et al., 2003; Kincses et al., 2008; Rieckmann et al., 2010; Sefcsik et al., 2009); in contrast, the role of the hippocampus remains inconclusive (Albouy et al., 2008; Schendan et al., 2003). A major approach to this research is through brain imaging and neuropsychological studies; in addition to these, experiments investigating the effects of pharmacological agents provide an opportunity for the better understanding of the biological background of implicit learning (for review see Uddén et al., 2010). For example, a study by Frank et al. (2006) showed that the benzodiazepine midazolam, which inactivates the hippocampus, causes explicit memory deficits in healthy participants, but enhances implicit learning. In contrast, a more recent study found impaired implicit learning after the exogenous administration of the stress hormone cortisol (Römer et al., 2011). The engagement of specific brain structures in these phenomena needs to be clarified.

In experimental settings, implicit learning is defined as the acquisition of co-occurrence/dependencies between stimuli or trials, and is expressed only through performance (Frensch, 1998; Howard et al., 2004; Rieckmann and Bäckman, 2009). In the past decades, several tasks have been developed to tap into implicit learning. These tasks can be organized into two main groups based on whether the covariation or the temporal sequence of stimuli has predictive information. For example, in artificial grammar learning, participants are exposed to strings of letters. They are not informed that the strings follow a set of rules; yet, it has been found that they can apply these rules at a later stage of practice (Dienes et al., 1991; Reber, 1989). In the weather prediction task, individuals have to decide whether a specific combination of cards predicts rainy or sunny weather. They are unaware that each combination of cards is probabilistically related to a particular weather outcome. During the task, participants learn gradually which of two outcomes will occur, although they have no conscious knowledge of the rule (Gluck et al., 2002; Kemény and Lukács, 2009; Kincses et al., 2004; Poldrack and Rodríguez, 2004). Similarly, in the contextual cueing task, the global configuration of a display cues the location of a search target (Chun and Jiang, 1998; Howard et al., 2006).

In these tasks, the covariation of certain stimuli (e.g., in a letter string/a set of cards) has predictive information, in contrast to sequence learning tasks, where participants have to predict the onset of a stimulus based on the preceding stimuli (Rieckmann and Bäckman, 2009). Evidence suggests that the latter type of task has partly different underlying mechanisms and activates partly different brain structures (Greene et al., 2007; Jimenez and Vázquez, 2011; Poldrack et al., 2005); therefore, it is important to differentiate between these two types of tasks. In recent years, a growing body of data has emerged regarding the consolidation of implicit sequence learning, while covariation learning has received less attention. In our review, we focus on the perceptual–motor learning of sequences. First, we describe the sequence learning tasks in more details. We follow this with a discussion of the consolidation processes and its potentially influencing factors. Finally, we consider developing questions and future directions on this research field.

2. Measures of implicit sequence learning

A widely used sequence learning task is the finger tapping task (Fig. 1A). Here, participants are instructed to produce a particular sequence of finger movements either on a response box or by opposing their fingers to their thumb (Doyon et al., 2002; Karni et al., 1995). Performance is measured by the number of correctly produced sequences over a certain time interval (e.g., 30 s). Similarly to the previously mentioned tasks, participants' performance becomes better with practice. The main problem with this task is that it can be difficult to classify as truly implicit, because participants are aware of the sequence they have to generate. However, it is based on the learning of sequences and the learning is expressed through performance; therefore, we discuss the results based on this task where it is relevant to the question of consolidation.

In the past decade, the serial reaction time (SRT) task (Nissen and Bullemer, 1987) and its modification, the alternating serial reaction time (ASRT) task (Howard and Howard, 1997; Nemeth et al., 2010b), have become the most popular implicit sequence learning tasks. In the original version of the SRT task, a stimulus appears at one of four possible locations on the screen, and subjects have to press the button corresponding to that location (Fig. 1A). They are unaware that the sequence of subsequent locations (and correspondingly, the sequence of the responses) follows a predetermined order (Hallgató et al., *in press*). For example, in classical SRT tasks, the structure of the sequence is deterministic, with the stimuli following a simple repeating pattern as in the series 213412431423, where the numbers refer to distinct events (Fig. 1B). Without becoming aware of the sequence, subjects learn the regularity – and as they learn, they produce faster and more accurate responses. When the sequence is changed to a random series of stimuli, subjects become slower and less accurate in responding. In this task, sequence learning is measured by the reaction time difference between sequence and random blocks.

The modified version of SRT is the alternating serial reaction time (ASRT) task (Howard and Howard, 1997; Nemeth et al., 2010b). Here, repeating events alternate with random ones in an eight-element sequence so that the location of every second stimulus in the stream is determined randomly. If, for instance, the sequence is 3214, where the numbers represent locations on the screen, the sequence of the stimuli will be 3R2R1R4R, with R representing a random element. The sequence is thus better hidden than in the classical SRT task. This structure is referred to as a probabilistic second-order dependency (Remillard, 2008). The structure is second-order in that for pattern trials, event $n-2$ predicts event n . It is probabilistic in that these pattern trials occur amid randomly determined ones. In addition, participants do not generally become aware of the alternating structure of the sequences, even after extended practice, and sensitive recognition tests indicate that people do not develop explicit knowledge of the event sequences that are more likely to occur (Howard et al., 2004; Howard and Howard, 1997; Song et al., 2007).

Because sequence, and random stimuli alternate in the ASRT task, some sequences of three events (called “triplets”) occur more frequently than others (Fig. 1B). In this task we can separate general skill learning from sequence-specific learning, where *general skill learning* refers to increasing speed as the result of practice. In contrast, *sequence-specific learning* refers to the acquisition of sequence-specific knowledge, resulting in relatively faster responses for more predictable high-frequency events compared to less predictable low-frequency events. In the classical SRT task, sequence learning is measured as the RT difference between sequence and random blocks (Fig. 1C). However, in this difference score, sequence-specific and general skill learning are mixed, because participants also exhibit general skill improvement on sequential blocks. Thus, the bigger difference between these two types of blocks could be attributed to sequence-specific and general skill learning together, and we cannot determine the extent of these two types of learning. ASRT allows these types of learning to be separated. Another advantage of the ASRT task compared to the classical SRT task is that in the ASRT, it is possible to track sequence-specific learning continuously by comparing responses to the random and sequence elements or more and less predictable events in all blocks. In summary, recent research favors the ASRT task, because 1) it is more implicit, 2) it can separate sequence-specific and general skill learning, and 3) it can continuously track these two types of learning across all blocks.

3. Consolidation of sequence knowledge

Sequence learning does not occur only during practice, in the so-called online periods, but also between practice periods, during the so-called offline periods. The process that occurs during the offline

Fig. 1. A) Examples of the most commonly used sequence learning paradigms. B) The main distinction in the case of sequence type is whether it is deterministic or probabilistic. C) Typical examples of performance measures in different sequence learning paradigms. In the finger tapping task, performance is measured by the number of correctly produced sequences over a certain time interval (e.g., 30 s). In the classical SRT task, sequence learning is measured as the RT difference between sequence (S) and random (R) blocks. In the probabilistic sequence learning tasks (e.g., the ASRT task), sequence-specific learning is measured by comparing responses to the random and sequence elements or more and less predictable events in all blocks.

periods is referred to as consolidation, which denotes the stabilization of a memory trace after the initial acquisition; this can result in increased resistance to interference or even improvement in performance following an offline period (Krakauer and Shadmehr, 2006; Nemeth et al., 2010b; Robertson, 2009; Song, 2009).

This review will focus on factors that can determine the post-encoding stabilization and enhancement phases of consolidation. The consolidation can include the integration of recently acquired information with past experiences (memory association), the anatomical reorganization of memory representations (memory translocation), reconsolidation of memory representations after recall (memory reconsolidation), and even the erasure of memory representations, all of which appear to occur outside of awareness and without additional training. These processes can be time dependent or sleep dependent (Stickgold et al., 2002; Walker et al., 2003b; Walker and Stickgold, 2004). Hence, it is essential to differentiate between time-dependent and sleep-dependent consolidation. *Time-dependent* consolidation reflects the stabilization or even improvement of the memory trace after an offline period, irrespective of whether sleep occurred in this period or not. Thus, in this type of consolidation, sleep is not an essential component.

In recent decades, special attention has been given to the role of sleep; for instance references are made to *sleep-dependent* consolidation (Walker and Stickgold, 2004) suggesting that performance improves more when the offline period includes sleep than when it does not. Several studies have shown the critical role of sleep in skill learning consolidation (Fischer et al., 2002; Maquet et al., 2000; Peigneux et al., 2003; Stickgold et al., 2002; Walker et al., 2002). Nonetheless, the results concerning consolidation of sequence

knowledge have been mixed, and recent findings indicate that whether or not offline improvements occur at all, and whether they are sleep dependent, varies with factors such as awareness, the length of the offline period, the type of information to be learned, and the age of the participants (Hallgató et al., in press; Nemeth and Janacsek, 2011; Press et al., 2005; Robertson et al., 2004; Song et al., 2007; Spencer et al., 2007).

Although a growing body of research and fertile models advance our understanding of online learning (e.g., Cohen and Squire, 1980; Hawkins et al., 2009; Henke, 2010; Willingham, 1997), less is known about the offline processes, and there are yet unresolved contradictions between some of the related findings. Therefore, it is critical to determine which factors can potentially influence the consolidation of sequence learning. We discuss four factors that must be taken into account in consolidation research: 1) The role of awareness, 2) the length of the offline period, 3) perceptual and motor sequences, and 4) the effect of age. Considering these factors, we can organize the findings that emerge more appropriately and build more effective models of consolidation.

3.1. The role of awareness in consolidation

Ample research using the finger tapping task suggests that sleep plays an important role in the consolidation of sequence learning (Doyon et al., 2009b; Fischer et al., 2002; Korman et al., 2003; Walker et al., 2003b; Walker and Stickgold, 2004). Walker et al. (2003b), for example, found that finger tapping skill developed to the highest degree during the first night of sleep following training, but additional nights of sleep also offered continued improvements.

Other studies have suggested that sleep-dependent improvement is independent of whether sleep occurs during daytime or nighttime; even a 90-minute daytime nap, immediately after learning, resulted in robust gains (Fischer et al., 2002; Korman et al., 2007). Korman et al. (2003) found that time in sleep determined the time of expression of these delayed gains.

Regarding the consolidation-related neural mechanisms, Fischer et al. (2005) showed that sleep-dependent improvement was linked to greatly reduced brain activation in the prefrontal, premotor, and primary motor cortical areas, along with a stronger involvement of the left parietal cortical regions. Investigating the electroencephalographic characteristics of post-training sleep, Morin et al. (2008) found an increased number and duration of sleep spindles during the offline consolidation of sequential finger movements.

Using the SRT task, Robertson et al. (2004) found offline improvement after sleep as well, but only if participants gained declarative knowledge about the sequence. Robertson (2009) argued that the consolidation of explicit and implicit learning is differentially affected by sleep and wakefulness. In another study, Fischer et al. (2007) suggested that sleep leads to offline enhancement by increasing the explicit knowledge about the learned information. Hence, when participants are aware of the sequential structure, we can observe sleep-dependent offline improvement (as in the studies using finger tapping or explicit SRT tasks). In some cases, however, this overnight enhancement in explicit knowledge can interfere with performance – for example, when the learning is implicit (Brown and Robertson, 2007; Galea et al., 2010) – even resulting in an overnight decrease of implicit knowledge (e.g., Fischer et al., 2007).

According to Robertson (2009), in implicit learning when there is no declarative knowledge about the task, consolidation may occur during both wakefulness and sleep. In line with this prediction, recent SRT and ASRT studies found similar consolidation of implicit sequence learning during both sleep and wakefulness (Nemeth et al., 2010b; Robertson et al., 2004; Song et al., 2007). For instance, Song et al. (2007) studied offline learning in young adults with the ASRT task, which has proven to be the most implicit sequence learning task because of its probabilistic properties (Howard et al., 2004). Participants were tested in three sessions, with an equivalent period of wake or sleep between sessions. There was no evidence of offline improvement in sequence-specific learning following either a period of sleep or wakefulness. The performance in the testing session, however, was the same as in the learning phase, suggesting that the sequential knowledge was well consolidated. Nemeth et al. (2010b) found similar results in elderly participants with the same method.

Summarizing these findings, when participants are aware of the sequence structure, offline improvement can occur only after sleep (for detailed explanation see Robertson, 2009). This claim is also in accordance with results from finger tapping tasks (Fischer et al., 2002; Walker et al., 2003b; Walker and Stickgold, 2004) and explicit SRT tasks (Robertson et al., 2004), because in those cases participants are aware of the sequence knowledge. In contrast, consolidation of implicit learning is independent of sleep. Thus, awareness of the sequential structure can modify the process of consolidation: When people gain explicit knowledge about the task, sleep-dependent consolidation occurs, while in the case of implicit learning, consolidation is only time dependent. Yet, it is important to note that there are other factors that may influence the effect of sleep in these tasks. For example, in finger tapping tasks, sequences are shorter and deterministic. In contrast, SRT tasks usually consist of longer deterministic sequences, while in ASRT tasks the sequences are probabilistic. Thus, the consolidation of implicit sequence knowledge might also depend on the type, length, and complexity of sequences. Nevertheless, to our knowledge, no study has comprehensively controlled and manipulated these factors.

We can interpret the role of awareness more deeply if we take into consideration that sleep affects evolutionarily newer brain structures,

such as the frontal lobe, compared to other brain structures (Hobson, 2009; Muzur et al., 2002). Studies on the relationship between cognitive functions and normal or disrupted sleep suggest that sleep has a greater impact on cognitive functions connected to the frontal lobe and other cortical structures than on those connected to subcortical structures (Bearpark et al., 1987; Bedard et al., 1993; Bedard et al., 1991; Cosentino et al., 2000; Engelman et al., 2000; Gosselin et al., 2006; Hoekema et al., 2007; Naeyegele et al., 2006; Nemeth et al., in press; Nemeth et al., 2010b; Pierobon et al., 2008; Robertson et al., 2004; Song et al., 2007). Furthermore, the awareness of sequential structure can be related to cortical structures, primarily to the frontal lobe (Gomez Beldarrain et al., 2008; Gomez Beldarrain et al., 2002; Squire et al., 1993; Squire et al., 1990). These claims can explain the findings that in finger tapping and explicit SRT tasks, sleep-dependent consolidation can be obtained, in contrast to ASRT studies where participants gain no explicit knowledge about the task. Consequently, we can conclude that implicit learning connected to subcortical structures does not benefit from sleep.

3.2. The length of the offline period

The previously mentioned studies used a 12- or 24-hour delay between sessions to specifically investigate the role of sleep. However, another important question is what happens during consolidation in a much shorter or a much longer time interval. Studies on the time course of skill consolidation indicate that there is a “critical period” after the learning phase that is necessary for the stabilization of memory traces (Nemeth and Janacsek, 2011). This time period depends on the task demand, and it varies from 1–2 h (Robertson et al., 2005) to 5 h (Shadmehr and Brashers-Krug, 1997; Shadmehr and Holcomb, 1997) or 6 h (Walker et al., 2003a). Before this time interval passes (e.g., immediately or 30 min after the learning session), the newly acquired skill is more sensitive to interference effects and no offline improvement is observed compared to longer delays. For example, using the SRT task, one study found that the gain of consolidation increased with the length of delay (Press et al., 2005). In this SRT study, no enhancement was found 1 h after the learning phase, but significant enhancement was observed after 4 h, and this further increased after 12 h. These results suggest that offline learning may be a dynamic process. However, these studies examined only a shorter stretch of time, so the question remains of what happens in consolidation after more than 12 h.

A recent ASRT study investigated the time course of implicit sequence learning by comparing the performance after 12-, 24-hour, and 1-week delays from the initial learning session (Nemeth and Janacsek, 2011). The researchers focused on the consolidation of implicit sequence-specific learning, and separately, general skill learning between young and elderly adults. The aim of the study was to determine a time point in a longer stretch of time at which improvement can still be observed in consolidation, and analyze its possible age-related differences. In the young adults, the researchers found offline improvement of the general skill after all three delay periods, with a gradual decline among delays. Although no offline improvement was found in sequence-specific learning with any of the consolidation intervals, it did not decrease significantly between sessions for young participants, suggesting that sequence-specific knowledge consolidated well. Thus, according to these results, offline general skill learning is influenced by the time course. In contrast, offline sequence-specific learning is not affected by the length of the offline period, since the consolidation of sequential knowledge had already occurred after a 12-hour delay, and did not differ from the 24-hour and 1-week delay conditions.

These results are congruent with recent theories of consolidation (Press et al., 2005; Robertson et al., 2005; Shadmehr and Brashers-Krug, 1997; Shadmehr and Holcomb, 1997; Walker et al., 2003a), which claim that memory stabilization occurs during the first 5–6 h

after learning. The strong offline improvement of general skill that was observed after 12 h may reflect this first stabilization process of memory traces, including the previously mentioned critical time period (Nemeth and Janacek, 2011). In addition, consolidation of sequence-specific information was similar in the 12-, 24-hour, and 1-week offline periods, independently of time course. These results suggest that the stabilization of sequence-specific memory is a faster process, because it had already occurred after 12 h and did not differ from the other delay conditions. In contrast, in the case of general skill learning, further changes were observed after longer time intervals (e.g., 24-hour and 1-week delays) compared to the 12-hour delay condition (Nemeth and Janacek, 2011). In another recent study, retention of the sequence-specific knowledge was found after a full year (Romano et al., 2010), suggesting that sequence knowledge without further practice is stable even after much longer periods.

3.3. *Perceptual and motor factors of sequence learning*

Another major debate in the field of sequence learning is whether we rely on “our hands” or on “our eyes” during learning (Deroost and Soetens, 2006; Keele et al., 2003; Kemény and Lukács, 2011; Mayr, 1996; Nemeth et al., 2009; Song et al., 2008; Ziessler and Nattkemper, 2001). In the classical sequence learning paradigms, such as the SRT and ASRT tasks, exactly what the participants learn is not clear: They might learn the sequence of the stimuli (perceptual learning), the sequence of their own eye movements (oculomotor learning), the sequence of response locations (response-based learning), or the sequence of given finger movements (effector-based learning) (Cohen et al., 1990; Remillard, 2003; Willingham, 1999). These factors determine not only the online learning process, but also the consolidation of sequence knowledge.

In an SRT study, Willingham (1999) used two conditions to examine the perceptual and the motor factors of learning. In one condition, the stimulus–response mapping was changed in the transfer (test) phase that followed the learning phase, so that half of the participants had to press the same sequence of keys as in the learning phase but saw new stimuli (motor condition), whereas the other half had to press a different sequence of keys as in the learning phase but saw the same stimuli as before (perceptual condition). Willingham (1999) found that transfer was shown only when the motor sequence was kept constant, and not when the perceptual sequence was constant. In a previous study, Nemeth et al. (2009) compared the magnitude of perceptual and motor components of implicit sequence learning using a modification of the ASRT-task in a similar design.

In the ASRT-Race task, the stimuli were the left, right, up, and down arrows, which appeared in the center of the screen (minimizing oculomotor movements). In the learning phase, participants had to mentally rotate the arrows by 90° to the right, and press the button corresponding to this rotated arrow. In the transfer phase, the stimulus–response mapping changed and participants no longer had to rotate the arrows (rather, they had to press the left button when seeing the left arrow). Half of the participants were assigned to the perceptual and the other half to the motor condition. In the perceptual condition, the perceptual sequence was the same, but the motor sequence (key presses) changed compared to the previously practiced sequence. Conversely, key presses followed the previously learned sequence and the perceptual information (the sequence of the stimuli displayed on the screen) changed in the motor condition. The goal of comparing the participants' performance between these two conditions was to determine whether the contribution of perceptual and motor component was the same as or different from the learning (for more details see Nemeth et al., 2009). This task involves second-order probabilistic sequences, whereas classical SRT tasks use deterministic sequences. ASRT-Race also allows “pure” sequence learning to be measured separately from general skill improvements, where sequence learning is reflected in the difference between the reaction times to more as opposed to less

predictable events. In addition, this task eliminates the possibility of oculomotor learning, as stimuli always appear in the same central position on the screen. In contrast to Willingham's findings, Nemeth et al. (2009) demonstrated that not only motor, but also perceptual learning of second-order probabilistic sequences is possible. Furthermore, Nemeth et al. (2009) showed that the two types of learning do not differ significantly in magnitude.

Although several studies have investigated the perceptual and motor components of online sequence learning (Deroost and Soetens, 2006; Mayr, 1996; Nemeth et al., 2009; Remillard, 2003, 2009; Song et al., 2008), to our knowledge, the effect of consolidation on the perceptual and motor characteristics of learning has received less empirical attention. The goal of the Hallgató et al.'s (in press) study was to fill this gap by investigating how the offline period modifies motor and perceptual components of implicit sequence learning. This issue is of particular interest because it deals with the question of whether sequence learning and consolidation are mediated primarily by perceptual or by motor brain networks (Deroost and Soetens, 2006; Goschke, 1998). Hallgató et al.'s study investigated the role of 12-hour and 24-hour delay on the perceptual and motor components of implicit sequence learning using the ASRT-Race task, while eliminating oculomotor learning. In addition, they aimed to explore the role of sleep in offline consolidation of these two factors of learning. Therefore, a 12-hour delay was administered between the learning phase and the transfer phase of the experiment, during which participants either slept (night group) or stayed awake (day group). They found significant sequence learning in the learning phase. Moreover, after the 12-hour and the 24-hour offline period, they found a significant learning effect in both the perceptual and the motor conditions; however, the transfer in the motor condition was more effective compared to the perceptual condition. They did not find any sleep effect on sequence learning in either condition.

The weaker consolidation of the perceptual component of sequence learning is in agreement with the results of Deroost and Soetens (2006) and Willingham (1999), who found evidence of perceptual learning only under specific conditions. According to several studies, perceptual learning only takes place when the structure of the sequence is simple, but in case of deterministic sequences with second-order dependencies and probabilistic sequences with first-order dependencies, perceptual learning is absent or only weakly present (Deroost and Soetens, 2006; Mayr, 1996; Remillard, 2003). In addition, previous studies found perceptual learning in explicit conditions (Russeler and Rosler, 2000) and when a motor sequence was learned concurrently (Mayr, 1996). In Hallgató et al.'s (in press) study, participants had no conscious awareness at all of the structure of the sequence. The only condition that met Deroost and Soetens (2006) criteria is that in the learning phase, participants learned the perceptual and motor components concurrently. Compared to Nemeth et al. (2009), who found similar magnitudes of perceptual and motor learning immediately after the learning phase, Hallgató et al. (in press) found a weaker perceptual learning effect in the transfer phase both after a 12-hour and a 24-hour delay. As the only difference was the 12-hour or the 24-hour delay, Hallgató et al. supposed that the differences between the results of the two studies can be related to the consolidation period only. Thus, this criterion, where in the Learning Phase, participants learned the perceptual and motor components concurrently, can be enough to infer that significant perceptual learning occurs immediately after the learning phase (Meier and Cock, 2010; Nemeth et al., 2009; Weiermann et al., 2010). However, it might also result in weaker consolidation after the delay period. One potential explanation for this is that brain structures underlying the perceptual and motor components of sequence learning are disconnected in the offline periods, and the perceptual component might be more sensitive to interference effects. To put the puzzle together, we propose that the consolidation period has a differential effect on the motor and perceptual

components of sequence learning, so that in the transfer phase, the motor component is larger than the perceptual one. However, more investigations are needed to determine the potential background mechanisms of this phenomenon.

Beyond the question of the perceptual and motor components of learning, Hallgató et al.'s study has relevance for the sleep debate in consolidation as well (Debas et al., 2010; Doyon et al., 2009b; Gerván and Kovács, 2007; Karni, 1994; Robertson, 2009; Song, 2009; Stickgold and Walker, 2005; Walker et al., 2002). Hallgató et al. also found that sleep does not benefit sequence learning. In addition, the role of sleep in the consolidation of motor and perceptual factors of implicit sequence learning exhibits no difference. Another plausible explanation besides those previously mentioned (e.g., awareness, the separation of general skill and sequence-specific learning) is that in the probabilistic sequence learning task used in this study, in addition to the primary sensory and motor brain regions, the subcortical structures and cerebellum are more involved (Doyon, 2008; Hikosaka et al., 1999; Hikosaka et al., 2002a). This contrasts with the more basic finger tapping tasks, where sleep-dependent improvement has usually been found (Walker et al., 2002).

3.4. The effect of aging on consolidation

As implicit sequence learning is highly dependent on the integrity of the striatal network, and age-related structural and biochemical losses are pronounced in the striatal complex and connected prefrontal areas (Dennis and Cabeza, 2011; Raz et al., 2005; Rieckmann and Bäckman, 2009), how online and offline sequence learning is affected by aging is an important question.

Several studies have shown that older adults exhibit online implicit sequence-specific learning comparable to young adults for simple repeating patterns in the SRT task (Brown et al., 2009; Fraser et al., 2009; Frensch and Miner, 1994; Howard and Howard, 1989; Howard and Howard, 1992). However, more recent studies have reported that although older adults can learn higher-order sequence structure, they show age-related deficits in doing so (Curran, 1997; Howard et al., 2004; Howard and Howard, 1997; Nemeth and Janacsek, 2011; Nemeth et al., 2010b). For example, using a version of the ASRT task, Bennett et al. (2007) found that old persons were able to learn even third-order dependencies (1RR2RR3 where R is a random element), although they learned less than the young control group.

Only a few studies have investigated the consolidation of implicit knowledge in older persons (Brown et al., 2009; Fraser et al., 2009; Nemeth and Janacsek, 2011; Nemeth et al., 2010b; Siengsukon and Boyd, 2009a, b; Spencer et al., 2007). Spencer et al. (2007), for example, used an implicit contextual version of the SRT task in order to specifically examine the effect of sleep. In a previous study, they found sleep-related offline improvement in this version of the task among young adults (probably because it's contextual component; Spencer et al., 2006). In older subjects, however, neither offline improvement nor a sleep effect was shown (Spencer et al., 2007). Compared to young adults, elderly participants showed deficits in consolidation. Similarly, Brown et al. (2009) found age-related consolidation deficits using the classical version of the SRT task. Nevertheless, neither Spencer et al. (2007) nor Brown et al. (2009) could distinguish general skill learning from sequence-specific learning in their tasks. As the ASRT task has been shown to yield offline general skill learning, but not offline sequence-specific learning in young adults (Song et al., 2007), it is important to differentially analyze these two aspects of implicit sequence learning in older adults as well.

Nemeth et al. (2010b) investigated implicit sequence learning after a 12-hour offline period. The novelty of this research in comparison to previous studies of consolidation in older adults (Brown et al., 2009; Siengsukon and Boyd, 2009a; Spencer et al., 2007) was that 1) it used probabilistic second-order sequences, and 2) it dealt separately with general skill and sequence-specific learning. The researchers

showed that general skill learning improved offline in both the young and older groups, with the young group improving more than the old group. However, the improvement was not sleep dependent, in that it was not relevant whether the 12-hour offline period included sleep or not. In the case of sequence-specific learning, they found no offline improvement in either group.

A more recent study investigated the consolidation of implicit sequence learning by comparing the performance after 12-, 24-hour, and 1-week delays from the initial learning session in young and elderly adults (Nemeth and Janacsek, 2011). In the young adults, the researchers found offline improvement of the general skill after all delays, with gradual decline among them. The elderly adults showed offline improvement of the general skill only after the 12-hour offline period, and this improvement was weaker than that in the young group. Although the pattern in age groups is similar, these results suggest that the offline course of general skill learning may be affected by aging, since Nemeth and Janacsek did not find improvement either after a 24-hour or 1-week delay in the elderly group. No offline improvement was found in sequence-specific learning in either age group with any of the consolidation intervals. Sequence-specific learning did not decrease significantly between sessions for young participants, suggesting that sequence-specific knowledge was well consolidated in this group, whereas the older group showed weaker consolidation in all delay conditions compared to the younger group. Thus, according to these results, offline general skill learning is influenced both by the time course and aging, while the offline sequence learning is affected only by aging.

The differences between the young and old groups suggest that older participants are more sensitive to the time course in general skill learning, in that they showed no offline improvement even after the 24-hour delay. Regarding the practical consequences, the differences among the 12-, 24-hour, and 1-week consolidation intervals suggest that during new skill acquisition, it could be important to hold the training sessions closer together for optimal performance, with shorter intervals for elderly participants (Nemeth and Janacsek, 2011).

In the case of sequence-specific learning, older people show deficits both in online learning when the sequence structure is more complex and in the consolidation of sequence knowledge. A recent neuroimaging study (Dennis and Cabeza, 2011) found that elderly participants recruited medial temporal lobe structures in implicit sequence learning tasks, suggesting that compensation mechanisms are present to perform to an optimal level in these tasks. However, further studies are needed to precisely determine these compensation mechanisms and their potential role in age-related dementia and rehabilitation programs.

4. Conclusion and remaining questions

In view of the above, we can conclude that consolidation is not a single process; instead, there are multiple mechanisms in the offline period (e.g. general skill, sequence-specific processes), which are differently influenced by the task demand, awareness of the sequence, the length of the delay period, perceptual and motor factors, and the age of the participant (Table 1). Contradictions in this field can occur due to low or absent control of these factors of sequence learning. For example, sleep does not benefit sequence learning performance in the case of probabilistic sequences (Nemeth et al., 2010b; Song et al., 2007), because such sequences are more complex and implicit compared to simple and explicit sequence structures such as those used in finger tapping tasks. Consequently, the awareness of the sequence structure can modify the role of sleep in the consolidation process (Robertson, 2009; Robertson et al., 2004; Song, 2009).

The length of the delay period between the learning and testing phase is also an important factor that differentially modifies general skill and sequence-specific learning. In the case of general skill learning, the length of the delay is critical: The highest offline

Table 1

Factors that influence consolidation of sequence learning and related studies.

Factors	Related studies
Awareness (explicit vs. implicit)	Fischer et al. (2007); Fischer et al. (2002); Korman et al. (2007); Korman et al. (2003); Nemeth et al. (2010a, 2010b); Robertson et al. (2004); Song et al. (2007); Walker et al. (2003b)
Length of delay period	Nemeth and Janacsek (2011); Press et al. (2005); Robertson et al. (2005); Romano et al. (2010); Shadmehr and Brashers-Krug (1997); Shadmehr and Holcomb (1997); Walker et al. (2003a)
Perceptual vs. motor	Hallgató et al. (in press)
Age of participants	Brown et al. (2009); Fraser et al. (2009); Nemeth and Janacsek (2011); Nemeth et al. (2010a, 2010b); Spencer et al. (2007)

improvement is observed after shorter delay periods (Nemeth and Janacsek, 2011). In contrast, the delay period has less effect on sequence-specific learning: Retention of the previously acquired knowledge has been observed after one week (Nemeth and Janacsek, 2011) and even one year (Romano et al., 2010), but without additional offline improvement. Separating the perceptual and motor factors of learning can further elucidate this issue in that consolidation of motor sequence learning is more robust than perceptual factor of sequence learning. Nevertheless, more investigations are needed to replicate this finding and further detail its potential background mechanisms. Finally, one of the most important factors that must be taken into consideration is the age of participants, since several studies found significant age-related deficits in the consolidation of both sequence-specific and general skill learning in elderly adults (Brown et al., 2009; Nemeth and Janacsek, 2011).

Further studies of sequence consolidation should take these factors into consideration and investigate the following issues: 1) How consolidation affects implicit sequence learning in childhood in the separate cases of general skill learning and sequence-specific learning; 2) whether there is differential consolidation of the perceptual and motor factors of sequence learning in older ages; 3) how the length of delay modifies the consolidation of explicit sequence knowledge in the case of general skill and sequence specific learning; and finally 4) whether the length of delay has a differential effect on perceptual and motor consolidation in the case of explicit sequence learning.

Ultimately, we know a lot about the biological background of on-line implicit sequence learning (Albouy et al., 2008; Dennis and Cabeza, 2011; Doyon et al., 2009a; Keele et al., 2003; Kincses et al., 2008; Poldrack et al., 2005; Rieckmann et al., 2010; Sefcsik et al., 2009). However, there is a huge gap in our knowledge related to brain plasticity during consolidation. Future neuroscientific investigations must address this question while controlling for the above-mentioned factors, especially the differentiation between general skill and sequence-specific learning. These distinctions can contribute to developing a more sophisticated picture of brain-consolidation-behavior interaction.

Acknowledgment

Thanks to Gabor Orosz for helpful comments on the manuscript. This review was supported by OTKA K 82068 (Hungarian Scientific Research Fund).

References

- Albouy, G., Sterpenich, V., Baletau, E., Vandewalle, G., Desseilles, M., Dang-Vu, T., Darsaud, A., Ruby, P., Luppi, P.H., Degueldre, C., Peigneux, P., Luxen, A., Maquet, P., 2008. Both the hippocampus and striatum are involved in consolidation of motor sequence memory. *Neuron* 58, 261–272.
- Bearpark, H., Grunstein, R., Touyz, S., Channon, L., Sullivan, C., 1987. Cognitive and psychological dysfunction in sleep apnea before and after treatment with CPAP. *Journal of Sleep Research* 16, 303–310.

- Bedard, M.A., Montplaisir, J., Richer, F., Rouleau, I., Malo, J., 1991. Obstructive sleep apnea syndrome: pathogenesis of neuropsychological deficits. *Journal of Clinical and Experimental Neuropsychology* 13, 950–964.
- Bedard, M.A., Montplaisir, J., Malo, J., Richer, F., Rouleau, I., 1993. Persistent neuropsychological deficits and vigilance impairment in sleep apnea syndrome after treatment with continuous positive airways pressure (CPAP). *Journal of Clinical and Experimental Neuropsychology* 15, 330–341.
- Bennett, I.J., Howard Jr., J.H., Howard, D.V., 2007. Age-related differences in implicit learning of subtle third-order sequential structure. *The Journals of Gerontology. Series B: Psychological Sciences* 62B, 98–103.
- Brown, R.M., Robertson, E.M., 2007. Off-line processing: reciprocal interactions between declarative and procedural memories. *The Journal of Neuroscience* 27, 10468–10475.
- Brown, R.M., Robertson, E.M., Press, D.Z., 2009. Sequence skill acquisition and off-line learning in normal aging. *PLoS One* 4, e6683.
- Chun, M.M., Jiang, Y., 1998. Contextual cueing: implicit learning and memory of visual context guides spatial attention. *Cognitive Psychology* 36, 28.
- Cleeremans, A., Destrebecqz, A., Boyer, M., 1998. Implicit learning: news from the front. *Trends in Cognitive Sciences* 2, 406–416.
- Cohen, N.J., Squire, L.R., 1980. Preserved learning and retention of pattern-analyzing skills in amnesia: dissociation of knowing how and knowing that. *Science* 210, 207–210.
- Cohen, A., Ivry, R.J., Keele, S.W., 1990. Attention and structure in sequence learning. *Journal of Experimental Psychology. Learning, Memory, and Cognition* 16, 17–30.
- Cosentino, F.I.L., Bosco, P., Drago, V., Prestianni, G., Lanuzza, B., Iero, I., Tripodi, M., Spada, R.S., Toscano, G., Caraci, F., Ferri, R., 2000. The APOE e4 allele increases the risk of impaired spatial working memory in obstructive sleep apnea. *Sleep Medicine* 9, 831–839.
- Curran, T., 1997. Effects of aging on implicit sequence learning: accounting for sequence structure and explicit knowledge. *Psychological Research* 60, 24–41.
- Debas, K., Carrier, J., Orban, P., Barakat, M., Lungu, O., Vandewalle, G., Tahar, A.H., Bellec, P., Karni, A., Ungerleider, L.G., Benali, H., Doyon, J., 2010. Brain plasticity related to the consolidation of motor sequence learning and motor adaptation. *Proceedings of the National Academy of Sciences of the United States of America* 107 (41), 17839–17844.
- Dennis, N., Cabeza, R., 2011. Age-related dedifferentiation of learning systems: an fMRI study of implicit and explicit learning. *Neurobiology of Aging* 32, 2318.e17–2318.e30.
- Deroost, N., Soetens, E., 2006. Perceptual or motor learning in SRT tasks with complex sequence structures. *Psychological Research* 70, 88–102.
- Dienes, Z., Broadbent, D., Berry, D., 1991. Implicit and explicit knowledge bases in artificial grammar learning. *Journal of Experimental Psychology. Learning* 17, 875–887.
- Doyon, J., 2008. Motor sequence learning and movement disorders. *Current Opinion in Neurology* 21, 478–483.
- Doyon, J., Song, A.W., Karni, A., Lalonde, F., Adams, M.M., Ungerleider, L.G., 2002. Experience-dependent changes in cerebellar contributions to motor sequence learning. *Proceedings of the National Academy of Sciences of the United States of America* 99, 1017–1022.
- Doyon, J., Bellec, P., Amel, R., Penhune, V., Monchi, O., Carrier, J., Lehericy, S., Benali, H., 2009a. Contributions of the basal ganglia and functionally related brain structures to motor learning. *Behavioural Brain Research* 199, 61–75.
- Doyon, J., Korman, M., Morin, A., Dostie, V., Hadj Tahar, A., Benali, H., Karni, A., Ungerleider, L.G., Carrier, J., 2009b. Contribution of night and day sleep vs. simple passage of time to the consolidation of motor sequence and visuomotor adaptation learning. *Experimental Brain Research* 195, 15–26.
- Engelman, H.M., Kingshott, R.N., Martin, S.E., Douglas, N.J., 2000. Cognitive function in the sleep apnea/hypopnea syndrome (SAHS). *Sleep* 23, 102–108.
- Fischer, S., Hallschmid, M., Elsner, A.L., Born, J., 2002. Sleep forms memory for finger skills. *Proceedings of the National Academy of Sciences of the United States of America* 99, 11987–11991.
- Fischer, S., Nitschke, M.F., Melchert, U.H., Erdmann, C., Born, J., 2005. Motor memory consolidation in sleep shapes more effective neuronal representations. *The Journal of Neuroscience* 25, 11248–11255.
- Fischer, S., Fischer, S., Wilhelm, I., Born, J., 2007. Developmental differences in sleep's role for implicit off-line learning: comparing children with adults. *Journal of Cognitive Neuroscience* 19, 214.
- Frank, M.J., O'Reilly, R.C., Curran, T., 2006. When memory fails, intuition reigns: Midazolam enhances implicit inference in humans. *Psychological Science* 17, 700–707.
- Fraser, S.A., Li, K.Z.H., Penhune, V.B., 2009. A comparison of motor skill learning and retention in younger and older adults. *Experimental Brain Research* 195, 419–427.
- Frensch, P.A., 1998. One concept, multiple meanings. In: Stadler, M.A., Frensch, P.A. (Eds.), *Handbook of Implicit Learning*. SAGE Publications, London, pp. 47–104.
- Frensch, P.A., Miner, C.S., 1994. Effects of presentation rate and individual differences in short-term memory capacity on an indirect measure of serial learning. *Memory & Cognition* 22, 95–110.
- Galea, J.M., Albert, N.B., Ditye, T., Miall, R.C., 2010. Disruption of the dorsolateral prefrontal cortex facilitates the consolidation of procedural skills. *Journal of Cognitive Neuroscience* 22, 1158–1164.
- Gerván, P., Kovács, I., 2007. Sleep dependent learning in contour integration. *Journal of Vision* 7, 48.
- Gluck, M.A., Shohamy, D., Myers, C., 2002. How do people solve the “weather prediction” task?: individual variability in strategies for probabilistic category learning. *Learning & Memory* 9, 408–418.
- Gomez Beldarrain, M., Gafman, J., Ruiz De Valesco, V., Pascual-Leone, A., Garcia-Monco, J.C., 2002. Prefrontal lesions impair the implicit and explicit learning of sequences on visuomotor tasks. *Experimental Brain Research* 142, 529–538.

- Gomez Beldarrain, M., Astorgano, A.G., Gonzalez, A.B., Garcia-Monco, J.C., 2008. Sleep improves sequential motor learning and performance in patients with prefrontal lobe lesions. *Clinical Neurology and Neurosurgery* 110, 245–252.
- Goschke, T., 1998. Implicit learning of perceptual and motor sequences: evidence for independent learning systems. In: Stadler, M.A., Frensch, P.A. (Eds.), *Handbook of Implicit Learning*. SAGE Publications, London, pp. 401–444.
- Gosselin, N., Mathieu, A., Mazza, S., Petit, D., Malo, J., Montplaisir, J., 2006. Attentional deficits in patients with obstructive sleep apnea syndrome: an event-related potential study. *Clinical Neurophysiology* 117, 2228–2235.
- Greene, A.J., Gross, W.L., Elsinger, C.L., Rao, S.M., 2007. Hippocampal differentiation without recognition: an fMRI analysis of the contextual cueing task. *Learning & Memory* 14, 548–553.
- Hallgató, E., Györi-Dani, D., Pekár, J., Janacsek, K., Nemeth, D., in press. The differential consolidation of perceptual and motor learning in skill acquisition. *Cortex*.
- Hawkins, J., George, D., Niemasik, J., 2009. Sequence memory for prediction, inference and behaviour. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences* 364, 1203–1209.
- Heerey, E.A., Velani, H., 2010. Implicit learning of social predictions. *Journal of Experimental Social Psychology* 46, 577–581.
- Henke, K., 2010. A model for memory systems based on processing modes rather than consciousness. *Nature Reviews. Neuroscience* 11, 523–532.
- Hikosaka, O., Nakahara, H., Rand, M.K., Sakai, K., Lu, X., Nakamura, K., Miyachi, S., Doya, K., 1999. Parallel neural networks for learning sequential procedures. *Trends in Neurology Science* 22, 464–471.
- Hikosaka, O., Nakamura, K., Sakai, K., Nakahara, H., 2002. Central mechanisms of motor skill learning. *Current Opinion in Neurobiology* 12, 217–222.
- Hobson, J.A., 2009. REM sleep and dreaming: towards a theory of protoconsciousness. *Nature Reviews. Neuroscience* 10, 803–813.
- Hoekema, A., Stegenga, B., Bakker, M., Brouwer, W.H., de Bont, L.G., Wijkstra, P.J., van der Hoeven, J.H., 2007. Simulated driving in obstructive sleep apnoea–hypopnoea; effects of oral appliances and continuous positive airway pressure. *Sleep & Breathing* 11, 129–138.
- Howard, D.V., Howard Jr., J.H., 1989. Age differences in learning serial patterns: direct versus indirect measures. *Psychology and Aging* 4, 357–364.
- Howard, D.V., Howard, J.H., 1992. Adult age differences in the rate of learning serial patterns: evidence from direct and indirect tests. *Psychology and Aging* 7, 232–241.
- Howard Jr., J.H., Howard, D.V., 1997. Age differences in implicit learning of higher-order dependencies in serial patterns. *Psychology and Aging* 12, 634–656.
- Howard, D.V., Howard Jr., J.H., Japikse, K., DiYanni, C., Thompson, A., Somberg, R., 2004. Implicit sequence learning: effects of level of structure, adult age, and extended practice. *Psychology and Aging* 19, 79–92.
- Howard Jr., J.H., Howard Jr., J.H., Howard, D.V., Japikse, K.C., Eden, G.F., 2006. Dyslexics are impaired on implicit higher-order sequence learning, but not on implicit spatial context learning. *Neuropsychologia* 44, 1131.
- Jimenez, L., Vázquez, G.A., 2011. Implicit sequence learning and contextual cueing do not compete for central cognitive resources. *Journal of Experimental Psychology: Human* 37, 222–235.
- Karni, A., 1994. Dependence on Rem-sleep of overnight improvement of a perceptual skill. *Science* 265, 679–682.
- Karni, A., Meyer, G., Jezard, P., Adams, M.M., Turner, R., Ungerleider, L.G., 1995. Functional MRI evidence for adult motor cortex plasticity during motor skill learning. *Nature* 377, 155–158.
- Kaufman, S.B., DeYoung, C.G., Gray, J.R., Jiménez, L., Brown, J., Mackintosh, N., 2010. Implicit learning as an ability. *Cognition* 116, 321–340.
- Keele, S.W., Ivry, R., Mayr, U., Hazeltine, E., Heuer, H., 2003. The cognitive and neural architecture of sequence representation. *Psychological Review* 110, 316–339.
- Kemény, F., Lukács, Á., 2009. The effects of feature analysis, transparency in probabilistic category learning in adults and children. *Learning Perception* 1, 199–213.
- Kemény, F., Lukács, Á., 2011. Perceptual effect on motor learning in the serial reaction-time task. *The Journal of General Psychology* 138, 110–126.
- Kincses, T.Z., Antal, A., Nitsche, M.A., Bártfai, O., Paulus, W., 2004. Facilitation of probabilistic classification learning by transcranial direct current stimulation of the prefrontal cortex in the human. *Neuropsychologia* 42, 113–117.
- Kincses, T., Johansen-Berg, H., Tomassini, V., Bosnell, R., Matthews, P., Beckmann, C., 2008. Model-free characterization of brain functional networks for motor sequence learning using fMRI. *NeuroImage* 39, 1950–1958.
- Korman, M., Raz, N., Flash, T., Karni, A., 2003. Multiple shifts in the representation of a motor sequence during the acquisition of skilled performance. *Proceedings of the National Academy of Sciences of the United States of America* 100, 12492–12497.
- Korman, M., Doyon, J., Doljansky, J., Carrier, J., Dagan, Y., Karni, A., 2007. Daytime sleep condenses the time course of motor memory consolidation. *Nature Neuroscience* 10, 1206–1213.
- Krakauer, J.W., Shadmehr, R., 2006. Consolidation of motor memory. *Trends in Neurosciences* 29, 58–64.
- Lieberman, M.D., 2000. Intuition: a social cognitive neuroscience approach. *Psychological Bulletin* 126, 109–137.
- Maquet, P., Laureys, S., Peigneux, P., Fuchs, S., Petiau, C., Phillips, C., Aerts, J., Del Fiore, G., Degueldre, C., Meulemans, T., Luxen, A., Franck, G., Van Der Linden, M., Smith, C., Cleeremans, A., 2000. Experience-dependent changes in cerebral activation during human REM sleep. *Nature Neuroscience* 3, 831–836.
- Mayr, U., 1996. Spatial attention and implicit sequence learning: evidence for independent learning of spatial and nonspatial sequences. *Journal of Experimental Psychology* 22, 350–364.
- Meier, B., Cock, J., 2010. Are correlated streams of information necessary for implicit sequence learning? *Acta Psychologica* 133, 17–27.
- Morin, A., Doyon, J., Dostie, V., Barakat, M., Hadj Tahar, A., Korman, M., Benali, H., Karni, A., Ungerleider, L.G., Carrier, J., 2008. Motor sequence learning increases sleep spindles and fast frequencies in post-training sleep. *Sleep* 31, 1149–1156.
- Muzur, A., Pace-Schott, E.F., Hobson, J.A., 2002. The prefrontal cortex in sleep. *Trends in Cognitive Sciences* 6, 475–481.
- Naegele, B., Launois, S.H., Mazza, S., Feuerstein, C., Pepin, J.L., Levy, P., 2006. Which memory processes are affected in patients with obstructive sleep apnea? An evaluation of 3 types of memory. *Sleep* 29, 533–544.
- Nemeth, D., Janacsek, K., 2011. The dynamics of implicit skill consolidation in young and elderly adults. *The Journals of Gerontology. Series B: Psychological Sciences* 66, 15–22.
- Nemeth, D., Hallgató, E., Janacsek, K., Sandor, T., Londe, Z., 2009. Perceptual and motor factors of implicit skill learning. *NeuroReport* 20, 1654–1658.
- Nemeth, D., Janacsek, K., Balogh, V., Londe, Z., Mingesz, R., Fazekas, M., Jambori, S., Danyi, I., Vetro, A., 2010a. Learning in autism: implicitly superb. *PLoS One* 5, e11731.
- Nemeth, D., Janacsek, K., Londe, Z., Ullman, M.T., Howard, D., Howard, J., 2010b. Sleep has no critical role in implicit motor sequence learning in young and old adults. *Experimental Brain Research* 201, 351–358.
- Nemeth, D., Janacsek, K., Csicsak, G., Szvoboda, G., Howard Jr., J.H., Howard, D.V., 2011. Interference between sentence processing and probabilistic implicit sequence learning. *PLoS ONE* 6, e17577.
- Nemeth, D., Csábi, E., Janacsek, K., Várszegi, M., Mari, Z., in press. Intact implicit probabilistic sequence learning in obstructive sleep apnea. *Journal of Sleep Research*. doi:10.1111/j.1365-2869.2011.00983.x.
- Nissen, M.J., Bullemer, P., 1987. Attentional requirements of learning: evidence from performance measures. *Cognitive Psychology* 19, 1–32.
- Peigneux, P., Laureys, S., Fuchs, S., Destrebecqz, A., Collette, F., Delbeuck, X., Phillips, C., Aerts, J., Del Fiore, G., Degueldre, C., Luxen, A., Cleeremans, A., Maquet, P., 2003. Learned material content and acquisition level modulate cerebral reactivation during posttraining rapid-eye-movements sleep. *NeuroImage* 20, 125–134.
- Pierobon, A., Giardini, A., Fanfulla, F., Callegari, S., Majani, G., 2008. A multidimensional assessment of obese patients with obstructive sleep apnoea syndrome (OSAS): a study of psychological, neuropsychological and clinical relationships in a disabling multifaceted disease. *Sleep Medicine* 9, 882–889.
- Poldrack, R.A., Rodriguez, P., 2004. How do memory systems interact? Evidence from human classification learning. *Neurobiology of Learning and Memory* 82, 324–332.
- Poldrack, R.A., Sabb, F.W., Foerster, K., Tom, S.M., Asanow, R.F., Bookheimer, S.Y., Knowlton, B.J., 2005. The neural correlates of motor skill automaticity. *The Journal of Neuroscience* 25, 5356–5364.
- Press, D.Z., Casement, M.D., Pascual-Leone, A., Robertson, E.M., 2005. The time course of off-line motor sequence learning. *Brain Research. Cognitive Brain Research* 25, 375–378.
- Raz, N., Lindenberger, U., Rodrigue, K.M., Kennedy, K.M., Head, D., Williamson, A., Dahle, C., Gerstorf, D., Acker, J.D., 2005. Regional brain changes in aging healthy adults: general trends, individual differences and modifiers. *Cerebral Cortex* 15, 1676–1689.
- Reber, A.S., 1989. Implicit learning and tacit knowledge. *Journal of Experimental Knowledge* 118, 219–235.
- Reber, A.R., 1993. *Implicit Learning and Tacit Knowledge: An Essay on the Cognitive Unconscious*. Oxford University Press, New York.
- Remillard, G., 2003. Pure perceptual-based sequence learning. *Journal of Experimental Psychology. Learning* 29, 581–597.
- Remillard, G., 2008. Implicit learning of second-, third-, and fourth-order adjacent and nonadjacent sequential dependencies. *The Quarterly Journal of Experimental Psychology* 61, 400–424.
- Remillard, G., 2009. Pure perceptual-based sequence learning: a role for visuospatial attention. *Journal of Experimental Psychology. Learning* 35, 528.
- Rieckmann, A., Bäckman, L., 2009. Implicit learning in aging: extant patterns and new directions. *Neuropsychologia Review* 19, 490–503.
- Rieckmann, A., Fischer, H., Bäckman, L., 2010. Activation in striatum and medial temporal lobe during sequence learning in younger and older adults: relations to performance. *NeuroImage* 50, 1303–1312.
- Robertson, E.M., 2009. From creation to consolidation: a novel framework for memory processing. *PLoS Biology* 7, e1000019.
- Robertson, E.M., Pascual-Leone, A., Press, D.Z., 2004. Awareness modifies the skill-learning benefits of sleep. *Current Biology* 14, 208–212.
- Robertson, E.M., Press, D.Z., Pascual-Leone, A., 2005. Off-line learning and the primary motor cortex. *The Journal of Neuroscience* 25, 6372–6378.
- Romano Bergstrom, J.C., Howard, J.H., Howard, D.V., in press. Enhanced Implicit Sequence Learning in College-age Video Game Players and Musicians. *Applied Cognitive Psychology*. doi:10.1002/acp.1800.
- Romano, J.C., Howard, J.H., Howard, D.V., 2010. One-year retention of general and sequence-specific skills in a probabilistic, serial reaction time task. *Memory* 18, 427–441.
- Römer, S., Schulz, A., Richter, S., Lass-Hennemann, J., Schälinger, H., 2011. Oral cortisol impairs implicit sequence learning. *Psychopharmacology* 215, 33–40.
- Russeler, J., Rosler, F., 2000. Implicit and explicit learning of event sequences: evidence for distinct coding of perceptual and motor representations. *Acta Psychologica* 104, 45–67.
- Schendan, H., Searl, M., Melrose, R., Stern, C., 2003. An fMRI study of the role of the medial temporal lobe in implicit and explicit sequence learning. *Neuron* 37, 1013–1025.
- Sefcsik, T., Nemeth, D., Janacsek, K., Hoffmann, I., Scialabba, J., Klivenyi, P., Ambrus, G.G., Haden, G., Vecsei, L., 2009. The role of the putamen in cognitive functions—a case study. *Learning Perception* 1, 215–227.

- Shadmehr, R., Brashers-Krug, T., 1997. Functional stages in the formation of human long-term motor memory. *The Journal of Neuroscience* 17, 409–419.
- Shadmehr, R., Holcomb, H.H., 1997. Neural correlates of motor memory consolidation. *Science* 277, 821–825.
- Siengsukon, C.F., Boyd, L.A., 2009a. Does sleep promote motor learning? Implications for physical rehabilitation. *Physical Therapy* 89, 370.
- Siengsukon, C.F., Boyd, L.A., 2009b. Sleep to learn after stroke: implicit and explicit offline motor learning. *Neuroscience Letters* 451, 1–5.
- Song, S., 2009. Consciousness and the consolidation of motor learning. *Behavioural Brain Research* 196, 180–186.
- Song, S., Howard Jr., J.H., Howard, D.V., 2007. Sleep does not benefit probabilistic motor sequence learning. *The Journal of Neuroscience* 27, 12475–12483.
- Song, S., Howard, J.H.J., Howard, D.V., 2008. Perceptual sequence learning in a serial reaction time task. *Experimental Brain Research* 189, 145–158.
- Spencer, R.M., Sunm, M., Ivry, R.B., 2006. Sleep-dependent consolidation of contextual learning. *Current Biology* 16, 1001–1005.
- Spencer, R.M., Gouw, A.M., Ivry, R.B., 2007. Age-related decline of sleep-dependent consolidation. *Learning & Memory* 14, 480–484.
- Squire, L.R., Zola-Morgan, S., Cave, S.B., Haist, F., Musen, G., Suzuki, W.A., 1990. Memory: organization of brain systems and cognition. *Cold Spring Harbor Symposia on Quantitative Biology*. Cold Spring Harbor Laboratory Press, pp. 1007–1023.
- Squire, L.R., Knowlton, B.J., Musen, G., 1993. The Structure and Organization of Memory. *Annual Review of Psychology* 44, 453–495.
- Stickgold, R., Walker, M.P., 2005. Sleep and Memory: the Ongoing Debate. *Sleep* 28, 1225–1227.
- Stickgold, R., Fosse, R., Walker, M.P., 2002. Linking brain and behavior in sleep-dependent learning and memory consolidation. *Proceedings of the National Academy of Sciences of the United States of America* 99, 16519–16521.
- Uddén, J., Folia, V., Petersson, K.M., 2010. The neuropharmacology of implicit learning. *Current Neuropharmacology* 8, 367–381.
- Ullman, M.T., 2004. Contributions of memory circuits to language: the declarative/procedural model. *Cognition* 92, 231–270.
- Walker, M.P., Stickgold, R., 2004. Sleep-dependent learning and memory consolidation. *Neuron* 44, 121–133.
- Walker, M.P., Brakefield, T., Morgan, A., Hobson, J.A., Stickgold, R., 2002. Practice with sleep makes perfect: sleep-dependent motor skill learning. *Neuron* 35, 205–211.
- Walker, M.P., Brakefield, T., Hobson, J.A., Stickgold, R., 2003a. Dissociable stages of human memory consolidation and reconsolidation. *Nature* 425, 616–620.
- Walker, M.P., Brakefield, T., Seidman, J., Morgan, A., Hobson, J.A., Stickgold, R., 2003b. Sleep and the time course of motor skill learning. *Learning & Memory* 10, 275–284.
- Weiermann, B., Cock, J., Meier, B., 2010. What matters in implicit task sequence learning: perceptual stimulus features, task sets, or correlated streams of information? *Journal of Experimental Psychology*. *Learning* 36, 1492.
- Willingham, D.B., 1997. Systems of memory in the human brain. *Neuron* 18, 5–8.
- Willingham, D.B., 1999. Implicit motor sequence learning is not purely perceptual. *Memory & Cognition* 27, 561–572.
- Ziessler, M., Nattkemper, D., 2001. Learning of event sequences is based on response-effect learning: further evidence from a serial reaction task. *Journal of Experimental Psychology*. *Learning* 27, 595–613.

Appendix II

Janacsek, K., Fiser J., & Nemeth, D. (in press).

The best time to acquire new skills: Age-related differences in implicit sequence learning across life span.

Developmental Science. DOI: 10.1111/j.1467-7687.2012.01150.x.

Impact factor: 3.53.

PAPER

The best time to acquire new skills: age-related differences in implicit sequence learning across the human lifespan

Karolina Janacsek,¹ József Fiser² and Dezso Nemeth¹

1. Institute of Psychology, University of Szeged, Hungary

2. Department of Cognitive Science, Central European University, Budapest, Hungary

Abstract

Implicit skill learning underlies obtaining not only motor, but also cognitive and social skills through the life of an individual. Yet, the ontogenetic changes in humans' implicit learning abilities have not yet been characterized, and, thus, their role in acquiring new knowledge efficiently during development is unknown. We investigated such learning across the lifespan, between 4 and 85 years of age with an implicit probabilistic sequence learning task, and we found that the difference in implicitly learning high- vs. low-probability events – measured by raw reaction time (RT) – exhibited a rapid decrement around age of 12. Accuracy and z-transformed data showed partially different developmental curves, suggesting a re-evaluation of analysis methods in developmental research. The decrement in raw RT differences supports an extension of the traditional two-stage lifespan skill acquisition model: in addition to a decline above the age 60 reported in earlier studies, sensitivity to raw probabilities and, therefore, acquiring new skills is significantly more effective until early adolescence than later in life. These results suggest that due to developmental changes in early adolescence, implicit skill learning processes undergo a marked shift in weighting raw probabilities vs. more complex interpretations of events, which, with appropriate timing, prove to be an optimal strategy for human skill learning.

Introduction

It is widely accepted that children should be introduced to sports, music or languages early in their life if they are to develop a high proficiency, because late learners seldom become true champions or elite musicians or gain command of a second language similar to that of a native speaker. These observations contradict traditional measures of the ability of factual learning of declarative memories, which showed that humans become increasingly better at many learning tasks up until their late twenties (Craik & Bialystok, 2006). However, an important component of developing new abilities is related to implicit unconscious statistical learning processes (Hikosaka, Nakamura, Sakai & Nakahara, 2002; Keele, Ivry, Mayr, Hazeltine & Heuer, 2003) that underlie the acquisition of not only motor but also cognitive and social skills (Doyon, Bellec, Amsel, Penhune, Monchi, Carrier, Lehericy & Benali, 2009; Hikosaka *et al.*, 2002; Lieberman, 2000; Poldrack, Sabb, Foerde, Tom, Asarnow, Bookheimer & Knowlton, 2005; Ullman, 2001). Thus, to understand complex skill acquisition, the characteristics of both explicit declarative and implicit learning, such as the differences in their efficiency across

the lifespan, must be clarified. In contrast to declarative memory (Tulving & Craik, 2000), the ontogenetic changes in humans' implicit learning abilities have not yet been comprehensively characterized, and, thus, their role in acquiring new knowledge efficiently during development is unknown. The main goal of our study was to examine age differences in implicit learning across the human lifespan using the same task for all groups.

The computational underpinnings and the neural substrates of these different kinds of learning mechanisms are also controversial (Henke, 2010). Explicit learning has been linked more closely to medial temporal lobes of the cortex (Dennis & Cabeza, 2011; Squire & Zola, 1996). By contrast, implicit skill learning often requires fine-tuning of the perceptual-motor system based on experience; therefore, most models of implicit skill learning emphasize the role of the basal ganglia and the cerebellum (Cohen, Pascual-Leone, Press & Robertson, 2005; Dennis & Cabeza, 2011; Doyon *et al.*, 2009; Hikosaka, Nakahara, Rand, Sakai, Lu, Nakamura, Miyachi & Doya, 1999; Hikosaka *et al.*, 2002), whereas the role of the hippocampus remains inconclusive (Albouy, Sterpenich, Balteau, Vandewalle, Desseilles, Dang-Vu, Darsaud, Ruby, Luppi, Degueldre, Peigneux,

Address for correspondence: Dezso Nemeth, Institute of Psychology, University of Szeged, Egyetem u. 2, 6722, Szeged, Hungary; e-mail: nemethd@edpsy.u-szeged.hu

The authors contributed equally to this work. The authors report no conflict of interest and have no financial disclosure.

Luxen & Maquet, 2008; Schendan, Searl, Melrose & Stern, 2003). However, these models focused mostly on motor skill-related learning with less emphasis on more complex skills that could involve learning abstract cognitive dependencies implicitly. The second goal of our study was to relate our behavioral results to the various computational models of explicit and implicit learning.

Two main approaches to implicit learning emerged in developmental neuroscience with a different assessment of how learning abilities change with age: (1) the developmental invariance model and (2) the age-related changes model. Studies supporting the developmental invariance model of implicit learning failed to find significant age-related differences in learning (Meulemans, Van der Linden & Perruchet, 1998; Vinter & Perruchet, 2000). In support of this view, infant studies have shown that adult-like implicit learning mechanisms exist even in very early infancy (Clohessy, Posner & Rothbart, 2001; Saffran, Aslin & Newport, 1996). Developmental invariance models explain this age-independence by linking implicit (or procedural) learning to evolutionarily primitive brain regions, such as the basal ganglia and the cerebellum. These regions are characterized as early-maturation regions and are relatively resistant to neurological impairments (Reber, 1993).

By contrast, the age-related changes models posit that considerable developmental differences can be observed in implicit learning. Several of these studies found that older children and young adults showed stronger learning effects compared to very young participants (Fletcher, Maybery & Bennett, 2000; Kirkham, Slemmer, Richardson & Johnson, 2007; Maybery, Taylor & O'Brien-Malone, 1995; Thomas, Hunt, Vizueta, Sommer, Durston, Yang & Worden, 2004). These models accept the fronto-striatal origin of such learning, but they focus on evidence of continued development of these regions that form the basis of the behavioral changes with age (e.g. Thomas *et al.*, 2004). We compared our empirical results using a new approach to the problem of multiple neural substrates of learning proposed by Daw, Niv and Dayan (2005).

Serial reaction time task and the development of implicit learning

In our study, we used a modified version of the Serial Reaction Time (SRT) Task, which is one of most commonly used methods for measuring implicit skill learning. Serial Reaction Time Task is a four-choice reaction time task containing a hidden repeating sequence that the subject comes to predict and learn implicitly (Nissen & Bullemer, 1987; Poldrack *et al.*, 2005). In an SRT study, Meulemans *et al.* (1998) found that 6- and 10-year-old children showed similar degrees of learning as young adults. In contrast, Thomas *et al.* (2004) found that the learning performance of young adults was better than 7- to 10-year-old children. Studies investigating implicit skill learning at older ages also revealed inconsistent results. For example, several studies have demonstrated

that, for simple repeating patterns (in the SRT task), the extent of implicit sequence learning in elderly adults was comparable to young adults (Frensch & Miner, 1994; D.V. Howard & Howard, 1989, 1992). Moreover, in a recent study, Gaillard, Destrebecqz, Michiels and Cleeremans (2009) found that young (22-year-old), middle-aged (45-year-old), and elderly (71-year-old) participants performed at the same level.

The studies mentioned above used fixed (deterministic) sequences, which can be easily learned, making it less possible to detect age-related differences in learning. Furthermore, they cannot purely determine the acquired sequence-specific knowledge because these tasks (finger-tapping, classical SRT) confound general improvements with sequence-specific learning. Here, we used a modified version of the SRT task, the Alternating Serial Reaction Time (ASRT) task (J.H. Howard & Howard, 1997), which enabled us to measure the 'pure' sequence-specific learning distinguished from general improvements. In the classical SRT task, the structure of a sequence is deterministic with the stimuli following a simple cyclically repeating pattern (e.g. 2134121341213412..., where numbers refer to distinct events within the repeating 21341 pattern). By contrast, in the ASRT task (J.H. Howard & Howard, 1997; Remillard, 2008), repeating events alternate with random ones. Thus, the location of every second stimulus on the screen was determined randomly. If, for instance, the sequence was 12341234..., where the numbers represent locations on the screen, the sequence of stimuli would be 1R2R3R4R1R2R3R4R... in the ASRT paradigm, with R representing a random element. Therefore, the location of every second stimulus on the screen was determined randomly. Because fixed, sequence-specific and random stimuli were alternating, some sequences of three events (called 'triplets') occurred more frequently than others. For example, in the above illustration 1x2, 2x3, 3x4 and 4x1 would occur often, whereas 1x3 or 4x2 would occur infrequently. Following previous studies, we referred to the former as *high-frequency* triplets and the latter as *low-frequency* triplets (Nemeth, Janacek, Londe, Ullman, Howard & Howard, 2010b; Song, Howard & Howard, 2007). Previous studies have shown that as people practice the ASRT task, they respond more quickly to the high- than low-frequency triplets, revealing probabilistic, sequence-specific learning (J.H. Howard & Howard, 1997; Song *et al.*, 2007). This learning is statistical in nature because it depends on the frequency of the event sequences. Thus, the RT difference between the high- and low-frequency triplets in this ASRT task is a measure of human sensitivity to the relative raw probabilities of events observed implicitly in their environment (Perruchet & Pacton, 2006). In addition, the participants are not generally aware of the alternating structure of the sequences, even after extended practice, or when sensitive recognition tests are used to assess explicit knowledge (D.V. Howard, Howard, Japikse, DiYanni, Thompson & Somberg, 2004; J.H.

Howard & Howard, 1997; Song *et al.*, 2007). Thus, the ASRT task is more implicit than the classical deterministic sequence learning tasks.

Using the ASRT task, recent studies have shown that, although elderly adults can also learn the higher-order structure of these complex sequences, they showed age-related deficits (D.V. Howard *et al.*, 2004; J.H. Howard & Howard, 1997; Nemeth *et al.*, 2010b). Both young and elderly adults were able to learn third-order dependencies (1RR2RR3RR1RR2RR3RR...) although the elderly participants performed at a lower level than the younger participants (Bennett, Howard & Howard, 2007). Whereas several studies investigated implicit learning in children using the ASRT task (Barnes, Howard, Howard, Gilotty, Kenworthy, Gaillard & Vaidya, 2008; Barnes, Howard, Howard, Kenealy & Vaidya, 2010; Nemeth, Janacek, Balogh, Londe, Mingesz, Fazekas, Jámboři, Danyi & Vetró, 2010a), no child–adult comparison of implicit skill learning performance has yet been reported.

In summary, previous studies have addressed the development and aging in implicit skill learning, but no studies have examined age-related differences from childhood to old age with identical methods. Furthermore, in contrast to general skill improvements, using a probabilistic sequence learning task (ASRT) can help us to reveal the age-related differences of the underlying mechanisms of complex skill learning by measuring explicitly the sensitivity to raw probabilities of high- and low-frequency events. Therefore, in this study, we compared the implicit probabilistic sequence learning across the age range of 4–85 years.

Method

Participants

There were 421 participants in the experiment between the ages of 4 and 85 that were clustered into nine age groups between 4–6, 7–8, 9–10, 11–12, 14–17, 18–29, 30–44, 45–59 and 60–85 years of age (Table 1). None of them suffered from any developmental, psychiatric or neurological disorders. All subjects gave signed informed consent (parental consent was obtained for children), and they received no financial compensation for partic-

ipation. All experimental procedures were approved by the Ethics Committee of the University of Szeged.

Implicit probabilistic sequence learning task

We used the ASRT task (J.H. Howard & Howard, 1997; Nemeth *et al.*, 2010b) where a stimulus (a dog's head) appeared in one of the four empty circles arranged in a line on a computer screen. The participants were instructed to respond to different stimulus events by pressing the corresponding response keys (Z, C, B or M) as fast and accurately as possible. The ASRT task consisted of 20 blocks with 85 key presses in each block. The first five responses of each stimulus block served for practice only, and then the eight-element alternating sequence (e.g. 1R2R3R4R) was repeated 10 times within a block. Stimuli were presented 120 ms after the response to the previous stimulus. Between blocks, the participants received feedback on the screen about their overall reaction time (RT) and accuracy. The computer program generated a different repeating ASRT sequence of the four locations for each participant using a permutation rule such that each of the six unique permutations of the four repeating events occurred with equal probability.

To determine the amount of explicit knowledge the subjects acquired about the task, a short questionnaire was administered after the experimental session (Song *et al.*, 2007). This questionnaire included increasingly specific questions, such as 'Have you noticed anything special regarding the task?', 'Have you noticed some regularity in the sequence of stimuli?' The experimenter rated subjects' answers on a 5-point scale where 1 denoted 'Nothing noticed' and 5 denoted 'Total awareness'. None of the participants, young or old, reported noticing the hidden repeating sequence.

Statistical properties of the ASRT task

As mentioned above, the ASRT allows a comparison between responses to high- and low-probability events. For example, if the eight-element sequence is 1R2R3R4R, 1x2, 2x3, 3x4, and 4x1 would occur often (high-frequency triplets) because two consecutive stimuli of the repeating sequence (e.g. 132 consisting of 1R2) as well

Table 1 Demographic data and mean RT and accuracy in the different groups. In all columns, numbers in parentheses show standard deviation

Group	Age	Sex	Education	Mean RT (ms)	Mean accuracy (%)
4–6-year-old (<i>n</i> = 30)	5.31 (0.98)	17 M / 13 F	–	960.06 (214.67)	90.09 (6.34)
7–8-year-old (<i>n</i> = 55)	7.09 (0.56)	31 M / 24 F	1.18 (0.39)	773.24 (159.29)	90.66 (7.03)
9–10-year-old (<i>n</i> = 35)	9.89 (0.58)	14 M / 21 F	3.2 (0.96)	602.84 (121.03)	93.44 (4.21)
11–12-year-old (<i>n</i> = 29)	11.5 (0.5)	21 M / 8 F	4.66 (0.67)	544.15 (95.00)	92.52 (4.23)
14–17-year-old (<i>n</i> = 62)	14.89 (1.06)	46 M / 15 F	8.23 (1.02)	452.52 (67.06)	95.44 (2.92)
18–29-year-old (<i>n</i> = 63)	23.09 (3.67)	40 M / 23 F	15.45 (2.6)	401.79 (50.85)	95.47 (2.45)
30–44-year-old (<i>n</i> = 59)	35 (4.24)	24 M / 35 F	16.64 (3.1)	419.85 (58.68)	95.85 (2.98)
45–59-year-old (<i>n</i> = 36)	50.8 (5.07)	12 M / 24 F	14.18 (3.58)	526.7 (112.99)	97.4 (3.45)
60–85-year-old (<i>n</i> = 52)	69.85 (6.16)	16 M / 36 F	13.39 (3.04)	634.37 (126.54)	96.92 (2.38)

as two consecutive random elements by chance (e.g. the same 132 consisting of R3R) could form these triplets. By contrast, 1x3 or 4x2 would occur less frequently (low-frequency triplets) because they could never be obtained consisting of two consecutive sequence elements. Of the 64 possible triplets, 16 triplets were high-frequency triplets, occurring 62.5% of the time, whereas the remaining 48 triplets were low-frequency triplets, occurring 37.5% of the time. Thus, each low-frequency triplet occurred in approximately 0.8% of the total number of trials, whereas each high-frequency triplet occurred about five times more often, in approximately 4% of the trials. For each keypress response, we defined whether it was in response to a high- or a low-frequency element, depending on whether the element was more or less predictable based on the previous two items in the sequence.

Following the method of previous studies (D.V. Howard *et al.*, 2004; Song *et al.*, 2007), two kinds of low-frequency triplets were excluded from our analyses: repetitions (e.g. 222, 333) and trills (e.g. 212, 343). Repetitions and trills were low frequency for all participants, and in previous studies, the participants often showed pre-existing response tendencies towards them (D.V. Howard *et al.*, 2004; Soetens, Melis & Notebaert, 2004). The elimination of these special triplets from the analyses ensured that the high- versus low-frequency differences found in the study were not confounded by pre-existing response tendencies. After this adjustment, previous studies have found that, following the practice, participants responded more quickly to the high- than to the low-frequency triplets, revealing a sequence learning effect (D.V. Howard *et al.*, 2004; J.H. Howard & Howard, 1997; Nemeth *et al.*, 2010b; Song *et al.*, 2007).

Statistical analysis

We calculated the mean accuracy of all trials and the median reaction time (RT) of correct responses separately for high- and low-frequency triplets. The accuracy and RT measures were analyzed by mixed-model ANOVA with TRIPLET TYPE (high vs. low frequency) as the within-subject factor, and AGE (9 groups) as the between-subject factor. All significant results are reported together with the η_p^2 effect size and Greenhouse-Geisser ϵ correction factors where applicable. Post-hoc analysis was conducted by Fisher's LSD pairwise comparisons.

Results

Overall RTs significantly differed among the age groups (main effect of AGE: $F(8, 412) = 107.11$, $p < .001$, $\eta_p^2 = 0.675$; Table 1). The RT decreased significantly between each group from 4–6 to 18–29 years of age (all $ps < .04$), they were similar between the age groups of 18–29 and 30–44 ($p > .38$) and significantly increased after 44 years of age ($ps < .001$) (Figure 1a). The accuracy monotonically increased over the years (main effect of AGE: $F(8, 412) = 16.94$, $p < .001$, $\eta_p^2 = 0.25$) (Figure 1b).

The comparison of RT to high- vs. low-probability triplets showed a surprising pattern of implicit sequence learning across the age groups. Even though there was significant learning at all ages because the RTs were faster for high-frequency than to low-frequency triplets (main effect of TRIPLET TYPE: $F(1, 412) = 333.7$, $p < .001$, $\eta_p^2 = 0.45$, all $ps < .03$) (Figure 1c), the magnitude of this difference was not uniform. Although the age groups differed significantly from each other in

Figure 1 Sequence learning in all groups. Reaction time (a) and accuracy (b) for high- and low-frequency triplets are plotted. Learning measures of RT (c) and accuracy (d) represents the RT/ACC difference between low- and high-frequency triplets. Error bars indicate SEM.

sequence learning (TRIPLET TYPE $\times$ AGE interaction: $F(8, 412) = 6.79, p < .001, \eta_p^2 = 0.12$), the post-hoc test revealed that learning was significantly higher in the 4- to 12-year-old groups than in any other group in the 14–85 range ($ps < .02$). There was no difference in learning between the 14–59 years of age ($ps > .37$), whereas the magnitude of learning decreased significantly in the 60–85-year-old group ($ps < .02$). Thus, learning high-probability events was uniformly effective until the age of 12 where it reduced significantly and remained at a lower level of sensitivity until the age of 60 (Figure 1c).

However, it is a long-standing issue in developmental and aging studies how to compare groups with different baseline speeds. A customary approach to this problem is to analyze the data using z -transformation. Therefore, we calculated the z -scores within each subject (thus, each participant's own mean and SD were used to transform that participant's data (see, for example, Christ, White, Mandernach & Keys, 2001) and conducted an ANOVA based on these z -scores (Figure 2). ANOVA revealed significant sequence-specific learning (main effect of TRIPLET TYPE: $F(1, 412) = 320.12, p < .001, \eta_p^2 = 0.44$), but the extent of learning differed across groups (TRIPLET TYPE $\times$ AGE interaction: $F(8, 412) = 8.91, p < .001, \eta_p^2 = 0.15$). We found that the participants from 9 years of age showed a similar extent of sequence

learning as the adult groups to 44 years of age (all $ps > .25$), but the learning in 4–8 years of age was smaller compared to these adult groups (all $ps < .014$). At older ages, there was a decline in the sequence learning, with both the 45–59 and 60–85-year-old group differing significantly from the groups between 11 and 44 years of age (all $ps < .025$).

The analysis of the response accuracy further enhanced the picture emerging from the results with RTs. We found a significantly greater accuracy for high- than low-frequency triplets (main effect of TRIPLET TYPE: $F(1, 412) = 217.14, p < .001, \eta_p^2 = 0.345$). Although all age groups older than 6 showed significant sequence learning (all $ps < .011$), the age groups differed significantly in the strength of the sequence learning (TRIPLET TYPE $\times$ AGE interaction: $F(8, 412) = 3.73, p < .001, \eta_p^2 = 0.07$). Whereas groups between 7 and 44 years of age showed similar degrees of learning, this was significantly higher than the youngest (4–6) and the two oldest (45–59 and 60–85) groups ($ps < .03$) (Figure 1d).

Figure 3 shows the individual data for sequence learning measured by raw RT, accuracy and z -transformed RT data. The pattern of these data is consistent with the ANOVA results: (1) children between 4 and 12 years of age showed greater sequence learning as measured by raw RT, whereas (2) in sequence-specific learning as measured by accuracy and z -transformed RTs, adults exhibited the highest performance.

Discussion

The goal of the present study was to investigate the differences in implicit skill learning across the human lifespan. This work extends previous studies (Bennett *et al.*, 2007; Gaillard *et al.*, 2009; J.H. Howard & Howard, 1997; Meulemans *et al.*, 1998; Thomas *et al.*, 2004) in two ways: (1) it examined a wide range of ages between 4 and 85 years, and (2) it used a probabilistic task, which enabled us to measure the 'pure' sequence-specific learning defined by the sensitivity to raw probabilities of high- and low-frequency events. We found that the 4- to

Figure 2 Sequence learning measured the z -transformed RT data in all groups.

Figure 3 Individual data for sequence learning measured by raw RT (a), accuracy (b), and z -scores (c) in all ages.

12-year-old age groups showed the strongest learning effect measured by the raw RT difference scores. Around the age of 12, we found a striking transition to less pronounced sequence-specific learning, as measured by smaller differences between the responses to high- and low-frequency triplets. Confirming earlier results (D.V. Howard *et al.*, 2004; J.H. Howard & Howard, 1997; Nemeth *et al.*, 2010b), we found that this learning capacity was significantly reduced in the oldest age group. Thus, in contrast to the developmental invariance (Reber, 1993) and the age-related changes approaches (Meulemans *et al.*, 1998; Vinter & Perruchet, 2000), our results demonstrate a gradual decline in learning across the lifespan.

Sequence learning scores based on the accuracy and raw reaction time showed different curves: the former is a bell-shaped curve, whereas the latter is a gradually declining curve (Figure 1c–d). Hence, these two types of learning scores can reflect different underlying mechanisms and brain systems. The accuracy learning score may be more related to attention, mainly voluntary attention, whereas the RT learning score may be related to involuntary attention and intuitive processes (Burgess, Gilbert & Dumontheil, 2007; Prinzmetal, McCool & Park, 2005). The relatively weaker accuracy learning effects in children and older groups may be due to the underdeveloped/deteriorating attentional brain circuits connected to the frontal lobe.

Our study raised a methodological issue that affects developmental studies in general. It is a long-standing issue in the literature how to compare groups with different baseline speeds. It could be argued that the youngest groups in our study have larger raw RT learning scores (greater difference between high- and low-frequency triplets) because they have more room to do so, given their longer RTs. However this argument does not seem to hold in our study, because the oldest group showed equally long RTs as 9–10-year-olds, yet they had a fourfold reduced learning score (Figure 1). Along a similar line of argument, the accuracy measures of the youngest group (4–6-year-olds) and the oldest groups (45–85-year-olds) showed a marked difference (~8%) in general accuracy, giving more room for the young group to produce larger differences between high- and low-frequency triplets, yet the actual sequence learning (i.e. high-low differences in accuracy) showed no difference. In contrast, the two smallest age groups had almost identical general accuracy yet there was a more than twofold increase in sequence learning in the second youngest group (7–8-year-olds). Thus, our data suggest no linear relationship between the general magnitude of reaction time/accuracy and the learning measures.

A second customary approach to the problem of comparing groups with different baseline speeds is to analyze the data using *z*-transformation. *Z*-scores of our results show a different picture from the raw RT data analysis: the learning performances from 9 to 44 years of age are similar with weaker performance in the younger

and older age groups. The *z*-transformation is often used to control general processing speed across different age groups in developmental studies. However, the main function of *z*-transformation is not to control the processing speed, but to normalize the distribution of responses. Thus, *z*-transformation has mathematical assumptions about the form of distributions and is therefore not theory independent (Yap, Balota, Sibley & Ratcliff, 2012). *Z*-scoring fully adjusts for processing speed only if all participants have the same type of distributions. Therefore using *z*-scores in developmental studies might be misleading. Furthermore, it is unclear how general processing speed and variability contribute to learning and performance in different ages. In recent years, several studies analyzed the variability and noise across a wide age range (Der & Deary, 2006; Hultsch, MacDonald, Hunter, Levy-Bencheton & Strauss, 2000; Rabbitt, Osman, Moore & Stollery, 2001). For example, Rabbitt *et al.* (2001) found that people's fastest RTs were relatively unaffected by age, but the number of unnecessarily slow responses was higher in older ages, and, thus, the increase in the mean RT was a result of increasing variability, which was an important component of cognitive aging. Moreover, several studies outline that the noise and the performance variability enables adaptive plasticity of motor skills (Slifkin & Newell, 1998, 1999; Turner & Brainard, 2007) and high variability can support effective learning and performance (Sanger, 2010). Thus, based on previous studies and on our analyses, we suggest that both the processing speed (mean reaction time) and variability are inherent aspects of development and aging. We think that the *z*-transformation eliminates these inherent aspects of learning, making the transformed results difficult to understand and explain.

Based on the reasoning above, we propose that the raw RT difference between the high- and low-frequency triplets in the ASRT task is a measure of human sensitivity to the relative raw probabilities of events observed implicitly in their environment. Thus, our results show a marked decrease in this sensitivity around the age of 12, which is in contrast to the traditional view of a steady improvement of cognitive learning abilities until late in adulthood (Craik & Bialystok, 2006). However, this discrepancy might be explained based on a shift in the structural development of implicit learning. Although the raw probabilities of the sensory environment are important for learning and both infants (Aslin, Saffran & Newport, 1998; Fiser & Aslin, 2002; Saffran *et al.*, 1996; Saffran, Johnson, Aslin & Newport, 1999) and adults (Fiser & Aslin, 2001; Hunt & Aslin, 2001) are highly sensitive to them, there is an ongoing debate on how using these simple probabilities can lead to highly complex knowledge of the world, such as sensory invariances and development of a language (Gomez & Gerken, 1999; Marcus, Vijayan, Rao & Vishton, 1999). Recent studies have proposed that using an internally stored structured model of the world that emerges based

on past experience together with probabilistic learning could help to address this issue and also provide evidence that humans might implement such a strategy during implicit learning (Orban, Fiser, Aslin & Lengyel, 2008; Tenenbaum, Kemp, Griffiths & Goodman, 2011). In this framework, as the internal model develops with experiences becoming more influential, internal interpretations of events become more elaborate and less directly related to their raw probabilities. A recent study argued that from a normative standpoint, existence of multiple learning mechanisms in the brain (cf. model-free vs. model-based learning) with an uncertainty-based arbitration between them would be computationally optimal (Daw *et al.*, 2005). Anchoring this hypothesis biologically, the presumed mechanisms related to these two types of learning were suggested to be related to the prefrontal areas and temporal lobe of the cortex, respectively (Daw *et al.*, 2005). Support for the separated

and complementary nature of the prefrontal- and medial temporal lobe (MTL)-dependent learning based on internal models vs. basal ganglia-dependent model-free learning comes from various studies investigating learning under specific conditions. These studies showed that obstructing the PFC and/or MTL by a demanding secondary task (Foerde, Knowlton & Poldrack, 2006) does not adversely affect implicit learning. Other studies found that inserting a task between the learning sessions (Brown & Robertson, 2007a, 2007b), performing a working memory and an implicit learning task simultaneously (Filoteo, Lauritzen & Maddox, 2010), or a neuropharmacological blockage (Frank, O'Reilly & Curran, 2006) even had a positive effect on performance in an implicit learning task. Importantly, it is known that the cortical areas connected to the internal models related to model-based learning become truly functional late in development, around the age of 12 (Blakemore &

Figure 4 Competition between model-based and model-free neurocognitive subsystems of skill learning across lifespan. (a) Before adolescence, underdeveloped internal models (dashed boundary) have little influence on interpretations of detected raw statistical probabilities of events in the environment (dashed arrows). Skill learning performance is determined by raw probabilities. (b) From adolescence to late adulthood, well-developed internal models (solid boundary) strongly modulate the interpretations of observed statistics of the input. This helps in extracting complex relations but relatively impairs measuring and utilizing raw probabilities in skill learning (dotted arrow). (c) In older ages, skill learning performance decreases. This decline could be caused by the combination of reduced sensitivity to raw statistical probabilities (dashed boundary), increasingly rigid internal models (dashed boundary) and/or weaker connection between these systems (dashed arrows).

Choudhury, 2006; Giedd, Blumenthal, Jeffries, Castellanos, Liu, Zijdenbos, Paus, Evans & Rapoport, 1999), which is about the age when we found the sudden decrement in sensitivity to the relative raw probabilities. We propose that this enhanced functionality signals the shift when the system adapts efficiently to more complex aspects of the world by relying more on internal model-based interpretations, while somewhat neglecting the raw probabilities of the sensory input (Figure 4a–b), and therefore, decreasing the ability to develop and stabilize fundamentally new basic competences. Thus the seemingly paradoxical result of gradually becoming less sensitive to basic statistics, if timed appropriately, could be the optimal strategy for human implicit learning in general.

Our results did not reveal any differences between the young adults and middle-aged groups. Salthouse's (1996) 'simultaneity mechanism' theory of cognitive aging predicts the age-related deficits in probabilistic sequence learning (Curran, 1997; J.H. Howard & Howard, 1997). Feeney, Howard and Howard (2002) found age-related deficits in pattern sensitivity in 'older' (mean age: 49.4) compared with 'younger' (mean age: 41.4) middle-aged groups. These different results could be related to the fact that Feeney used a smaller sample size, a longer version of the ASRT and a different method of analysis.

What are the underlying mechanisms of the decreased performance of the elderly group? In a recent fMRI study, Dennis and Cabeza (2011) showed that older adults recruited the MTL for implicit learning, and this activation was significantly greater, while striatal activity decreased in older people compared with young adults during implicit learning. In a recent study, Rieckmann, Fischer and Bäckman (2010) found similar results: in young adults during the learning session, the activation of the striatum increased, but that of the MTL decreased. By contrast, in older adults, sequence learning positively related to activation increases in both the striatum and MTL. Using multimodal imaging measures, Giorgio, Santelli, Tomassini, Bosnell, Smith, De Stefano and Johansen-Berg (2010) found extensive reductions in the gray matter volume in aging, but reductions were detected earlier in the frontal cortex. Furthermore, a recent diffusion tensor imaging aging study by Bennett, Madden, Vaidya, Howard and Howard (2011) found that the caudate-dorsolateral prefrontal cortex (DLPFC) and hippocampus-DLPFC tract integrity were related to ASRT sequence learning. The caudate-DLPFC tract integrity decreased in the older ages, mediating age-related differences in sequence learning. Within the computational framework proposed by Daw and colleagues (2005), these findings can be interpreted as a deterioration in three mechanisms that contribute to the age-related decline in skill learning: (1) reduced detection of probabilities, (2) rigidity of internal models and/or (3) more restricted connections between internal models and probability detection (Figure 4c). Thus, not only

model-free but also model-based learning might be limited in older ages. Future studies are needed to systematically examine the underlying neural mechanisms of age-related differences in skill learning.

In summary, based on our raw RT results we suggest that acquiring fundamentally new skills that cannot be derived from skills already possessed is most effective before adolescence, and it might be largely based on the fronto-striatal circuitry, such as the basal ganglia and cerebellum, in agreement with earlier skill learning models (Doyon *et al.*, 2009; Hikosaka *et al.*, 1999; Hikosaka *et al.*, 2002) and computational learning models (Daw *et al.*, 2005). Our findings are in good agreement with everyday life experience showing that an early (~ before 12 years) start in learning some sports, musical instruments, second language, etc. often leads to higher levels of competence. These results may have implications for the development of learning and memory, facilitating new skill training and pedagogic methods (e.g. for teaching languages) and may also contribute to the understanding of neurodevelopmental and age-related disorders (e.g. autism, SLI, dyslexia and dementia) and lead to relevant treatment options.

Acknowledgements

Thanks to Darlene V. Howard, James H. Howard, Jr. and Michael Ullman for numerous illuminating discussions. Thanks also to Robert Mingesz, Attila Krajcsi, István Winkler, Gábor Orosz, Ágnes Lukács, Ágnes Kovács and Zsuzsa Londe for their comments on the previous version of the manuscript. This research was supported by the Bolyai Scholarship Program (to DN), by OTKA K 82068 (to DN), and by NIH NEI R01 EY018196 (to JF).

References

- Albouy, G., Sterpenich, V., Baletau, E., Vandewalle, G., Desseilles, M., Dang-Vu, T., Darsaud, A., Ruby, P., Luppi, P.H., Degueldre, C., Peigneux, P., Luxen, A., & Maquet, P. (2008). Both the hippocampus and striatum are involved in consolidation of motor sequence memory. *Neuron*, **58** (2), 261–272.
- Aslin, R.N., Saffran, J.R., & Newport, E.L. (1998). Computation of conditional probability statistics by 8-month-old infants. *Psychological Science*, **9** (4), 321–324.
- Barnes, K.A., Howard, J.H. Jr, Howard, D.V., Gilotty, L., Kenworthy, L., Gaillard, W.D., & Vaidya, C.J. (2008). Intact implicit learning of spatial context and temporal sequences in childhood autism spectrum disorder. *Neuropsychology*, **22** (5), 563–570.
- Barnes, K.A., Howard, J.H. Jr, Howard, D.V., Kenealy, L., & Vaidya, C.J. (2010). Two forms of implicit learning in childhood ADHD. *Developmental Neuropsychology*, **35** (5), 494–505.
- Bennett, I.J., Howard, J.H. Jr, & Howard, D.V. (2007). Age-related differences in implicit learning of subtle third-order sequential structure. *Journal of Gerontology: Psychological Sciences*, **62B** (2), 98–103.

- Bennett, I.J., Madden, D., Vaidya, C., Howard, J., & Howard, D. (2011). White matter integrity correlates of implicit sequence learning in healthy aging. *Neurobiology of Aging*, **32** (12), 2317.e1–2317.e12. doi: 10.1016/j.neurobiolaging.2010.03.017
- Blakemore, S.J., & Choudhury, S. (2006). Development of the adolescent brain: implications for executive function and social cognition. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, **47** (3–4), 296–312.
- Brown, R.M., & Robertson, E.M. (2007a). Inducing motor skill improvements with a declarative task. *Nature Neuroscience*, **10** (2), 148–149.
- Brown, R.M., & Robertson, E.M. (2007b). Off-line processing: reciprocal interactions between declarative and procedural memories. *Journal of Neuroscience*, **27** (39), 10468–10475.
- Burgess, P.W., Gilbert, S.J., & Dumontheil, I. (2007). Function and localization within rostral prefrontal cortex (area 10). *Philosophical Transactions of the Royal Society B: Biological Sciences*, **362** (1481), 887–899.
- Christ, S.E., White, D.A., Mandernach, T., & Keys, B.A. (2001). Inhibitory control across the life span. *Developmental Neuropsychology*, **20** (3), 653–669.
- Clohessy, A.B., Posner, M.I., & Rothbart, M.K. (2001). Development of the functional visual field. *Acta Psychologica*, **106** (1), 51–68.
- Cohen, D., Pascual-Leone, A., Press, D., & Robertson, E. (2005). Off-line learning of motor skill memory: a double dissociation of goal and movement. *Proceedings of the National Academy of Sciences, USA*, **102** (50), 18237–18241.
- Craik, F.I., & Bialystok, E. (2006). Cognition through the lifespan: mechanisms of change. *Trends in Cognitive Sciences*, **10** (3), 131–138.
- Curran, T. (1997). Effects of aging on implicit sequence learning: accounting for sequence structure and explicit knowledge. *Psychological Research*, **60** (1–2), 24–41.
- Daw, N.D., Niv, Y., & Dayan, P. (2005). Uncertainty-based competition between prefrontal and dorsolateral striatal systems for behavioral control. *Nature Neuroscience*, **8** (12), 1704–1711.
- Dennis, N., & Cabeza, R. (2011). Age-related dedifferentiation of learning systems: an fMRI study of implicit and explicit learning. *Neurobiology of Aging*, **32** (12), 2318.e17–2318.e30. doi: 10.1016/j.neurobiolaging.2010.04.004
- Der, G., & Deary, I.J. (2006). Age and sex differences in reaction time in adulthood: results from the United Kingdom Health and Lifestyle Survey. *Psychology and Aging*, **21** (1), 62–73.
- Doyon, J., Bellec, P., Amsel, R., Penhune, V., Monchi, O., Carrier, J., Lehericy, S., & Benali, H. (2009). Contributions of the basal ganglia and functionally related brain structures to motor learning. *Behavioral Brain Research*, **199** (1), 61–75.
- Feeney, J.J., Howard, J.H., & Howard, D.V. (2002). Implicit learning of higher order sequences in middle age. *Psychology and Aging*, **17** (2), 351–355.
- Filoteo, J.V., Lauritzen, S., & Maddox, W.T. (2010). Removing the frontal lobes. *Psychological Science*, **21** (3), 415–423.
- Fiser, J., & Aslin, R.N. (2001). Unsupervised statistical learning of higher-order spatial structures from visual scenes. *Psychological Science*, **12** (6), 499–504.
- Fiser, J., & Aslin, R.N. (2002). Statistical learning of new visual feature combinations by infants. *Proceedings of the National Academy of Sciences, USA*, **99** (24), 15822–15826. doi: 10.1073/pnas.232472899
- Fletcher, J., Maybery, M.T., & Bennett, S. (2000). Implicit learning differences: a question of developmental level? *Journal of Experimental Psychology: Learning, Memory, and Cognition*, **26** (1), 246–252.
- Foerde, K., Knowlton, B.J., & Poldrack, R.A. (2006). Modulation of competing memory systems by distraction. *Proceedings of the National Academy of Sciences, USA*, **103** (31), 11778–11783.
- Frank, M.J., O'Reilly, R.C., & Curran, T. (2006). When memory fails, intuition reigns: midazolam enhances implicit inference in humans. *Psychological Science*, **17** (8), 700–707.
- Frensch, P.A., & Miner, C.S. (1994). Effects of presentation rate and individual differences in short-term memory capacity on an indirect measure of serial learning. *Memory and Cognition*, **22** (1), 95–110.
- Gaillard, V., Destrebecqz, A., Michiels, S., & Cleeremans, A. (2009). Effects of age and practice in sequence learning: a graded account of ageing, learning, and control. *European Journal of Cognitive Psychology*, **21** (2), 255–282.
- Giedd, J.N., Blumenthal, J., Jeffries, N.O., Castellanos, F.X., Liu, H., Zijdenbos, A., Paus, T., Evans, A.C., & Rapoport, J.L. (1999). Brain development during childhood and adolescence: a longitudinal MRI study. *Nature Neuroscience*, **2** (10), 861–863.
- Giorgio, A., Santelli, L., Tomassini, V., Bosnell, R., Smith, S., De Stefano, N., & Johansen-Berg, H. (2010). Age-related changes in grey and white matter structure throughout adulthood. *NeuroImage*, **51** (3), 943–951.
- Gomez, R.L., & Gerken, L. (1999). Artificial grammar learning by 1-year-olds leads to specific and abstract knowledge. *Cognition*, **70** (2), 109–135.
- Henke, K. (2010). A model for memory systems based on processing modes rather than consciousness. *Nature Reviews Neuroscience*, **11**, 523–532.
- Hikosaka, O., Nakahara, H., Rand, M.K., Sakai, K., Lu, X., Nakamura, K., Miyachi, S., & Doya, K. (1999). Parallel neural networks for learning sequential procedures. *Trends in Neurosciences*, **22** (10), 464–471.
- Hikosaka, O., Nakamura, K., Sakai, K., & Nakahara, H. (2002). Central mechanisms of motor skill learning. *Current Opinion in Neurobiology*, **12** (2), 217–222.
- Howard, D.V., & Howard, J.H. Jr (1989). Age differences in learning serial patterns: direct versus indirect measures. *Psychology and Aging*, **4** (3), 357–364.
- Howard, D.V., & Howard, J.H. (1992). Adult age differences in the rate of learning serial patterns: evidence from direct and indirect tests. *Psychology and Aging*, **7** (2), 232–241.
- Howard, D.V., Howard, J.H. Jr, Japikse, K., DiYanni, C., Thompson, A., & Somberg, R. (2004). Implicit sequence learning: effects of level of structure, adult age, and extended practice. *Psychology and Aging*, **19** (1), 79–92.
- Howard, J.H. Jr, & Howard, D.V. (1997). Age differences in implicit learning of higher-order dependencies in serial patterns. *Psychology and Aging*, **12** (4), 634–656.
- Hultsch, D.F., MacDonald, S.W.S., Hunter, M.A., Levy-Bencheton, J., & Strauss, E. (2000). Intraindividual variability in cognitive performance in older adults: comparison of adults with mild dementia, adults with arthritis, and healthy adults. *Neuropsychology*, **14** (4), 588–598.
- Hunt, R.H., & Aslin, R.N. (2001). Statistical learning in a serial reaction time task: access to separable statistical cues by individual learners. *Journal of Experimental Psychology: General*, **130** (4), 658–680.

- Keele, S.W., Ivry, R., Mayr, U., Hazeltine, E., & Heuer, H. (2003). The cognitive and neural architecture of sequence representation. *Psychological Review*, **110** (2), 316–339.
- Kirkham, N., Slemmer, J., Richardson, D., & Johnson, S. (2007). Location, location, location: development of spatio-temporal sequence learning in infancy. *Child Development*, **78** (5), 1559–1571.
- Lieberman, M.D. (2000). Intuition: a social cognitive neuroscience approach. *Psychological Bulletin*, **126** (1), 109–137.
- Marcus, G.F., Vijayan, S., Rao, S.B., & Vishton, P.M. (1999). Rule learning by seven-month-old infants. *Science*, **283**, 77–80.
- Maybery, M., Taylor, M., & O'Brien-Malone, A. (1995). Implicit learning: sensitive to age but not IQ. *Australian Journal of Psychology*, **47** (1), 8–17.
- Meulemans, T., Van der Linden, M., & Perruchet, P. (1998). Implicit sequence learning in children. *Journal of Experimental Child Psychology*, **69** (3), 199–221.
- Nemeth, D., Janacsek, K., Balogh, V., Londe, Z., Mingesz, R., Fazekas, M., Jámboři, S., Danyi, I., & Vetró, A. (2010a). Learning in autism: implicitly superb. *PLoS ONE*, **5** (7), e11731.
- Nemeth, D., Janacsek, K., Londe, Z., Ullman, M.T., Howard, D., & Howard, J. (2010b). Sleep has no critical role in implicit motor sequence learning in young and old adults. *Experimental Brain Research*, **201** (2), 351–358. doi: 10.1007/s00221-009-2024-x
- Nissen, M.J., & Bullemer, P. (1987). Attentional requirements of learning: evidence from performance measures. *Cognitive Psychology*, **19**, 1–32.
- Orban, G., Fiser, J., Aslin, R.N., & Lengyel, M. (2008). Bayesian learning of visual chunks by human observers. *Proceedings of the National Academy of Sciences, USA*, **105** (7), 2745–2750.
- Perruchet, P., & Pacton, S. (2006). Implicit learning and statistical learning: one phenomenon, two approaches. *Trends in Cognitive Sciences*, **10** (5), 233–238.
- Poldrack, R.A., Sabb, F.W., Foerde, K., Tom, S.M., Asarnow, R.F., Bookheimer, S.Y., & Knowlton, B.J. (2005). The neural correlates of motor skill automaticity. *Journal of Neuroscience*, **25**, 5356–5364.
- Prinzmetal, W., McCool, C., & Park, S. (2005). Attention: reaction time and accuracy reveal different mechanisms. *Journal of Experimental Psychology: General*, **134** (1), 73–92.
- Rabbitt, P., Osman, P., Moore, B., & Stollery, B. (2001). There are stable individual differences in performance variability, both from moment to moment and from day to day. *Quarterly Journal of Experimental Psychology Section A*, **54** (4), 981–1003.
- Reber, A.R. (1993). *Implicit learning and tacit knowledge: An essay on the cognitive unconscious (Vol. 19)*. New York: Oxford University Press.
- Remillard, G. (2008). Implicit learning of second-, third-, and fourth-order adjacent and nonadjacent sequential dependencies. *Quarterly Journal of Experimental Psychology*, **61** (3), 400–424.
- Rieckmann, A., Fischer, H., & Bäckman, L. (2010). Activation in striatum and medial temporal lobe during sequence learning in younger and older adults: relations to performance. *NeuroImage*, **50** (3), 1303–1312.
- Saffran, J.R., Aslin, R.N., & Newport, E.L. (1996). Statistical learning by 8-month-old infants. *Science*, **274**, 1926–1928.
- Saffran, J.R., Johnson, E.K., Aslin, R.N., & Newport, E.L. (1999). Statistical learning of tone sequences by human infants and adults. *Cognition*, **70** (1), 27–52. doi: S0010-0277(98)00075-4
- Salthouse, T.A. (1996). The processing-speed theory of adult age differences in cognition. *Psychological Review*, **103** (3), 403–428.
- Sanger, T.D. (2010). Controlling variability. *Journal of Motor Behavior*, **42** (6), 401–407.
- Schendan, H., Searl, M., Melrose, R., & Stern, C. (2003). An fMRI study of the role of the medial temporal lobe in implicit and explicit sequence learning. *Neuron*, **37** (6), 1013–1025.
- Slifkin, A.B., & Newell, K.M. (1998). Is variability in human performance a reflection of system noise? *Current Directions in Psychological Science*, **7** (6), 170–177.
- Slifkin, A.B., & Newell, K.M. (1999). Noise, information transmission, and force variability. *Journal of Experimental Psychology: Human Perception and Performance*, **25** (3), 837–851.
- Soetens, E., Melis, A., & Notebaert, W. (2004). Sequence learning and sequential effects. *Psychological Research*, **69** (1), 124–137.
- Song, S., Howard, J.H. Jr, & Howard, D.V. (2007). Sleep does not benefit probabilistic motor sequence learning. *Journal of Neuroscience*, **27** (46), 12475–12483.
- Squire, L.R., & Zola, S.M. (1996). Structure and function of declarative and nondeclarative memory systems. *Proceedings of the National Academy of Sciences, USA*, **93**, 13515–13522.
- Tenenbaum, J.B., Kemp, C., Griffiths, T.L., & Goodman, N.D. (2011). How to grow a mind: statistics, structure, and abstraction. *Science*, **331** (6022), 1279–1285.
- Thomas, K.M., Hunt, R.H., Vizueta, N., Sommer, T., Durston, S., Yang, Y., & Worden, M.S. (2004). Evidence of developmental differences in implicit sequence learning: an fMRI study of children and adults. *Journal of Cognitive Neuroscience*, **16** (8), 1339–1351.
- Tulving, E., & Craik, F.I.M. (Eds.) (2000). *The Oxford handbook of memory*. Oxford: Oxford University Press.
- Turner, E.C., & Brainard, M.S. (2007). Performance variability enables adaptive plasticity of crystallized adult birdsong. *Nature*, **450** (7173), 1240–1244.
- Ullman, M.T. (2001). A neurocognitive perspective on language: the declarative/procedural model. *Nature Reviews Neuroscience*, **2**, 717–726.
- Vinter, A., & Perruchet, P. (2000). Implicit learning in children is not related to age: evidence from drawing behavior. *Child Development*, **71** (5), 1223–1240.
- Yap, M.J., Balota, D.A., Sibley, D.E., & Ratcliff, R. (2012). Individual differences in visual word recognition: insights from the English Lexicon Project. *Journal of Experimental Psychology: Human Perception and Performance*, **38** (1), 53–79. doi: 10.1037/a0024177

Received: 26 July 2010

Accepted: 13 January 2012

Appendix III

Nemeth, D., Janacsek, K., Polner, B., & Kovacs, Z. A. (in press).

Boosting human learning by hypnosis.

Cerebral Cortex. DOI: 10.1093/cercor/bhs068.

Impact factor: 6.844.

Boosting Human Learning by Hypnosis

Dezso Nemeth¹, Karolina Janacsek¹, Bertalan Polner¹ and Zoltan Ambrus Kovacs^{1,2}

¹Institute of Psychology and ²Department of Psychiatry, School of Medicine, University of Szeged, Szeged 6722, Hungary

Address correspondence to Dr Dezso Nemeth, Institute of Psychology, University of Szeged, Hungary, Egyetem u. 2., Szeged 6722, Hungary.
Email: nemethd@edpsy.u-szeged.hu.

Human learning and memory depend on multiple cognitive systems related to dissociable brain structures. These systems interact not only in cooperative but also sometimes competitive ways in optimizing performance. Previous studies showed that manipulations reducing the engagement of frontal lobe-mediated explicit attentional processes could lead to improved performance in striatum-related procedural learning. In our study, hypnosis was used as a tool to reduce the competition between these 2 systems. We compared learning in hypnosis and in the alert state and found that hypnosis boosted striatum-dependent sequence learning. Since frontal lobe-dependent processes are primarily affected by hypnosis, this finding could be attributed to the disruption of the explicit attentional processes. Our result sheds light not only on the competitive nature of brain systems in cognitive processes but also could have important implications for training and rehabilitation programs, especially for developing new methods to improve human learning and memory performance.

Keywords: functional connectivity, hypnosis, memory systems, prefrontal cortex, sequence learning, striatum

Introduction

Human learning and memory rely upon multiple cognitive systems related to separable brain structures. These systems interact in cooperative and sometimes competitive ways in optimizing memory and information processing performance (Poldrack et al. 2001; Poldrack and Packard 2003; Brown and Robertson 2007). Support for the competitive nature of memory systems comes from studies showing interactions between explicit/hypothesis-testing and implicit/procedural systems (Ashby et al. 1998; Poldrack and Packard 2003; Daw et al. 2005; Seger and Cincotta 2005; Filoteo et al. 2010). The former is often characterized by voluntary mechanisms relying more on attentional resources, and thought to be mediated by frontal and medial temporal lobe (MTL) structures, while the latter relies more on automatic, non-conscious processes mediated primarily by striatum. Manipulations reducing the engagement of the explicit, hypothesis-testing system, such as a demanding secondary task (Foerdes et al. 2006; Fu and Anderson 2008; Filoteo et al. 2010), a distractor task inserted between the learning sessions (Brown and Robertson 2007), or neuropharmacological blockage (Frank et al. 2006) had no effect or even led to performance improvements in striatum-dependent learning tasks. In a recent study, Galea et al. (2010) also found improvements in procedural learning after the disruption of the dorsolateral prefrontal cortex (PFC) using theta burst stimulation. In contrast, strengthening the reliance on explicit hypothesis-testing processes resulted in impaired procedural learning (Howard DV and Howard JH 2001; Fletcher et al.

2005) with greater PFC activity during the acquisition (Fletcher et al. 2005).

As rapid and reversible changes of cognitive processing are encountered in hypnosis, this phenomenon is an excellent tool of research in the cognitive neurosciences (Raz and Shapiro 2002; Egner et al. 2005). Regarding the neural background of hypnosis, studies demonstrated that people (especially with high susceptibility) show decreased performance on some frontal lobe-related tasks in hypnosis (Kaiser et al. 1997; Kallio et al. 2001; Wagstaff et al. 2007). More recent studies suggest reduced functional brain connectivity between cortical areas in hypnosis, and this is especially typical for frontal areas (Fingelkurts et al. 2007; Oakley and Halligan 2009). Hypnosis temporarily disconnects certain frontal areas from the anterior cingulate cortex and other brain areas, disturbing the frontal attentional control and executive system (Kaiser et al. 1997; Egner et al. 2005; Gruzelier 2006).

In our experiment, we used hypnosis as a tool to reduce the competition between frontal lobe-related explicit hypothesis testing and striatum-related procedural-based systems. In order to measure procedural learning in the hypnotic and alert states, we administered a motor sequence acquisition test, which is based on statistical learning mechanisms (Howard JH Jr and Howard DV 1997; Janacsek et al. Forthcoming; Perruchet and Pacton 2006). This fundamental learning mechanism underlies not only motor but also cognitive and social skills (Lieberman 2000; Ullman 2004; Kaufman et al. 2010; Nemeth et al. 2011; Romano Bergstrom et al. 2011); it is therefore an important aspect of life from infancy to old age. Sequence learning is essential for learning languages and operating appliances, such as, for example, computers and musical instruments (Howard et al. 2004; Romano et al. 2010). Social skills appear in compound behaviors realized in proper sequences activated under appropriate circumstances. Most models and empirical studies of sequence learning highlight the role of the basal ganglia (Hikosaka et al. 1999, 2002; Keele et al. 2003; Kincses et al. 2008; Dennis and Cabeza 2010; Rieckmann et al. 2010).

The main question of the study was how the disruption of frontal lobe functions by hypnosis affects performance in procedural-based sequence learning. Sequence learning was measured by the Alternating Serial Reaction Time (ASRT) task (Howard JH Jr and Howard DV 1997) on highly hypnotizable young adults. Participants performed the ASRT task both in waking alert and hypnotic state. In addition, executive functions were assessed by the Wisconsin Card Sorting Test (WCST) (Heaton et al. 1993; Anokhin et al. 2010) and Verbal Fluency Task (Spreen and Strauss 1991) in order to investigate the possible interactions between frontal lobe functions and the effect of hypnosis on sequence learning.

Materials and Methods

Participants

Fourteen healthy right-handed students from the University of Szeged participated in the experiment (12 females; mean age: 22.70, SD: 1.70; mean years of education: 15.50, SD: 1.58). All participants provided signed informed consent. They received no financial compensation for participating in the study. The study was approved by the Psychology Ethics Committee at the University of Szeged, Institute of Psychology.

Participants were selected from a pool of pretested subjects, on the basis of their hypnotic susceptibility. Hypnotizability was measured using the Hungarian version of the Harvard Group Scale of Hypnotic Susceptibility: Form A (HGSHS:A, Shor et al. 1962). Scoring procedure was based on the original English version (scores ranging from 0 to 12). Similarly to previous studies (Kallio et al. 2001; Halsband 2006), we defined high hypnotizability as having 8 or higher score on the HGSHS:A. The mean hypnotizability score of the participants was 9.07 (SD = 0.997; range from 8 to 12).

Tasks

Sequence Learning Task

The ASRT task is a widely used paradigm measuring implicit sequence learning in cognitive neuroscience (Song et al. 2007; Nemeth and Janacsek 2011). In this task, stimuli appear in 1 of 4 empty circles on the screen and participants are required to press the corresponding key (Y, C, B, or M on Hungarian keyboard) as quickly and accurately as possible. Participants are told that stimuli appear randomly, but, in fact, stimuli delivered in a random order (low predictability) alternate with the same stimulus items, which, however, follow a regular presentation order (high predictability). RTs to the high predictability stimuli become shorter than to the low predictability items as participants (implicitly) learn the hidden sequence.

The current ASRT task consisted of 1 practice block with random stimuli and 15 blocks with the alternating pattern described above. The latter blocks consisted of 85 key presses—the first 5 button pressings were random for practice purposes, then an 8-element alternating sequence (e.g., 2r1r3r4r, where numbers represent specific stimuli and r represents a random stimulus) was repeated 10 times. The response to stimulus interval was 120 ms. Participants were given different sequences in the 2 conditions (see below), in order to eliminate intersession learning effects.

Fluency Task

In this task, participants are instructed to produce as many words belonging to the same category (animals and supermarket) as possible in 60 s, without repetitions, synonyms, or generated forms of the same word (Spreen and Strauss 1991). The average number of correct words was used as the performance score. Higher score reflects better frontal lobe functions (Baldo et al. 2006).

The WCST

This task is one of the most specific tests of prefrontal functions (Heaton et al. 1993; Anokhin et al. 2010). Participants are required to derive a correct card-sorting rule based on a trial-by-trial feedback. As the rule changes without warning, the participant has to modify the previously learned response strategy on the basis of the feedback information. A key indicator of cognitive flexibility is the number of perseverative errors that occur when the participant persists in using the old strategy despite the negative feedback. A lower number of perseverative errors indicate better frontal lobe functions.

Design and Procedure

All tests were conducted on an individual basis. Participants performed the ASRT task in both the alert waking and the hypnotic state, with the same standard instructions. The order of the 2 conditions was counterbalanced across participants. The delay between the 2 sessions was 30 days. Fluency and WCST task was administered once, in a third session in alert state. Two participants did not take part in the third session due to time schedule problems.

A skilled hypnotist therapist (the author, Z.A.K.), who has extensive experience with hypnosis, tape recorded the induction, instructions, and dehypnotizing phases (similar to the study of Szendi et al. 2009). This recording was played to each participant. The type of hypnosis induction, similarly to that of the hypnotizability scale, was essentially relaxational.

The induction took approximately 14 min. After the induction, the hypnosis session began. When participants had completed half of the ASRT task, we played an approximately 30-s long induction in order to maintain their hypnotic state (in the wake condition, subjects had a 30-s rest). In the hypnosis condition, after the task had ended, the dehypnosis instruction was played.

Statistical Analysis

As there is a fixed sequence in the ASRT with alternating random elements (e.g., 2r3r1r4r), some triplets or runs of 3 events occur more frequently than others. For example, following the illustration above, triplets such as 2_3, 3_1, 1_4, 4_2 occur more often because the third element (bold numbers) could be both derived from the sequence as well as from a random element. In contrast, triplets such as 4_1, 4_4 would occur infrequently because, in this case, the third element could only come from the random stimuli. Following previous studies (e.g., Howard et al. 2004; Song et al. 2007), we refer to the former as high-frequency triplets and the latter as low-frequency triplets. Because of this difference in the occurrence frequencies of certain triplets, after observing two stimuli, a certain third stimulus can be expected with 62.5% of probability (for example, 223 is five times more probable than 221 or 222 or 224). In our analysis, we determined for each stimulus whether it was a more or less probable continuation for the previous 2 stimuli. Participants gave faster responses to the more probable than to the less probable stimuli, thus revealing sequence learning in the ASRT paradigm (Howard et al. 2004; Song et al. 2007). In addition, general skill learning can be observed in the ASRT task in the overall increase of the response speed, irrespective of the triplet type. Thus, we were able to separately measure sequence-specific and general skill learning in the ASRT task.

Similar to previous studies (e.g., Howard et al. 2004; Song et al. 2007; Nemeth et al. 2010), 2 kinds of low-frequency triplets were eliminated; repetitions (e.g., 222, 333) and trills (e.g., 212, 343). Repetitions and trills were low frequency for all participants, and participants often show preexisting response tendencies to them (Howard et al. 2004; Soetens et al. 2004). By eliminating these triplets, we could ascertain that any high- versus low-frequency differences were due to learning and not to preexisting tendencies.

Since the participants' accuracy remained very high throughout the test (as is typical, the average was more than 92% for both conditions; Howard JH Jr and Howard DV 1997; Nemeth et al. 2010), we focused on RT for the analyses reported. We calculated medians for correct responses only, separately for high- and low-frequency triplets, and for each participant and each third of the stimulus blocks (1–5, 6–10, and 11–15). To compare sequence learning between hypnosis and alert condition, and between groups with high- and low-executive functions, we conducted repeated measures and mixed design analyses of variance (ANOVAs) with Fisher's Least Significant Difference Test (LSD) post hoc tests. We reported the relevant effect sizes: η_p^2 for main effects and interactions, and Cohen's d measures for post hoc tests.

Results and Discussion

A repeated measures ANOVA was conducted with TRIPLET (2: high vs. low), BLOCK (3: 1–5, 6–10, and 11–15), and CONDITION (2: alert and hypnosis) as within-subjects factors. In this ANOVA, a significant main effect of TRIPLET reflects sequence-specific learning, which can increase with practice (TRIPLET $\times$ BLOCK interaction), while a significant TRIPLET $\times$ CONDITION interaction indicate differences in sequence-specific learning between the hypnosis and alert conditions.

ANOVA revealed significant sequence-specific learning (main effect of TRIPLET: $F_{1,13} = 16.21$, $P = 0.001$, $\eta_p^2 = 0.55$),

Figure 1. Sequence learning across blocks is plotted for waking alert (A) and hypnotic state (B), separately. Sequence learning performance (measured by the reaction time differences between high- and low-predictability events) was higher in the hypnotic state compared with the waking alert condition (C). Error bars represent standard error of mean.

Figure 2. Relationship between high versus low executive functions and the effect of hypnosis on sequence learning. Sequence learning across blocks for the high executive function group in waking alert (A) and hypnotic state (B), as well as sequence learning across blocks for the low executive function group in waking alert (D) and hypnotic state (E) is plotted. Participants with high executive functions showed smaller sequence-specific learning (measured by reaction time differences between high and low probability events) in the alert state compared with the hypnosis condition (C), while participants with lower executive functions showed similar extent of sequence learning in the waking alert and hypnotic state (F). Error bars represent standard error of mean.

which increased with practice (TRIPLET $\times$ BLOCK interaction: $F_{2,26} = 9.36$, $P = 0.001$, $\eta_p^2 = 0.42$). The 2 states differed significantly from each other (TRIPLET $\times$ CONDITION interaction: $F_{1,13} = 7.08$, $P = 0.02$, $\eta_p^2 = 0.35$): sequence learning was 2.5-times higher under hypnosis than in the waking alert state (Fig. 1). Independently from sequence learning, general RT decreased with practice (main effect of BLOCK: $F_{2,26} = 4.93$, $P = 0.034$, $\eta_p^2 = 0.27$). Other main effects and interactions were not significant (all P s > 0.34), thus the general RT was similar in the waking alert and hypnotic state (main effect of CONDITION: $F_{1,13} = 0.12$, $P = 0.73$, $\eta_p^2 = 0.009$). In addition, the rate of sequence learning process was also similar between the 2 conditions (TRIPLET $\times$ BLOCK $\times$ CONDITION: $F_{2,26} = 0.42$,

$P = 0.63$, $\eta_p^2 = 0.03$). Thus, hypnosis affected only sequence-specific learning (the difference between RTs for low- and high-probability events) and not the general RT. As we used a within-subject design with 2 learning sessions, a further analysis was conducted to test the possible effect of whether hypnosis was in the first or in the second session, and ANOVA revealed no order effect on sequence learning.

To calculate a composite score for executive function, first, we transformed measures of fluency task and WCST into z -scores. Then, we averaged these 2 transformed data into a composite score. Based on the median of this composite measure, we assigned half of the participants to the higher and other half to the lower executive function group. To compare

sequence learning between the high and the low executive function groups, in the alert and the hypnosis conditions (see Fig. 2), a mixed design ANOVA was conducted with TRIPLET (2: high vs. low), BLOCK (3: 1-5, 6-10, and 11-15), and CONDITION (2: alert and hypnosis) as within-subjects factors and GROUP (2: high vs. low executive function) as a between-subject factor.

The general RT was similar in the 2 groups (main effect of GROUP: $F_{1,10} = 1.5$, $P = 0.25$, $\eta_p^2 = 0.13$), and it was not affected differently by the 2 conditions (CONDITION $\times$ GROUP interaction: $F_{1,10} = 0.71$, $P = 0.42$, $\eta_p^2 = 0.07$). The TRIPLET $\times$ CONDITION interaction almost reached the significance ($F_{1,10} = 4.5$, $P = 0.06$, $\eta_p^2 = 0.31$) replicating that participants exhibited greater sequence learning under the hypnosis condition compared with the waking alert condition. The overall sequence learning was similar in the high and low executive function groups (TRIPLET $\times$ GROUP: $F_{1,10} = 3.07$, $P = 0.11$, $\eta_p^2 = 0.24$). Although the TRIPLET $\times$ CONDITION $\times$ GROUP interaction did not reach significance ($F_{1,10} = 1.72$, $P = 0.219$, $\eta_p^2 = 0.15$), the LSD post hoc tests revealed that participants with higher executive functions showed smaller sequence learning in the waking alert state compared with the hypnotic condition (Fig. 2C; $P = 0.03$, $d = 0.94$), while participants with lower executive functions showed similar extent of sequence learning (Fig. 2E; $P = 0.58$, $d = 0.25$). In addition, the learning performance of the high executive function group was significantly smaller compared with the low executive function group in the waking alert state ($P = 0.04$, $d = 1.31$), while it was similar in the hypnotic state ($P = 0.51$, $d = 0.396$).

Taken together, we found enhanced sequence learning performance in hypnosis. Our results provide support for the idea that learning and memory processes may not only involve the engagement of specific neuroplastic mechanisms but may also rely upon the disengagement of interacting systems (Brown and Robertson 2007, p. 149). Our finding is in line with previous studies demonstrating that manipulations reducing the reliance on frontal lobe-dependent processes improved procedural based learning performance (e.g., Filoteo et al. 2010; Galea et al. 2010).

The improved sequence learning in hypnosis could be attributed to the disruption of attentional control and executive system (Kaiser et al. 1997; Kallio et al. 2001; Wagstaff et al. 2007), by weakening the engagement of the frontal lobe and/or the interconnectivity between related brain areas (Egner et al. 2005; Gruzeliér 2006; Fingelkurts et al. 2007; Oakley and Halligan 2009). This could diminish the competition between 2 fundamentally incompatible modes of learning: 1) PFC/MTL-mediated hypothesis-testing attention-dependent processes versus 2) basal ganglia-dependent procedural learning (Ashby et al. 1998; Poldrack et al. 2001; Filoteo et al. 2010; Henke 2010). Reducing the reliance on executive hypothesis-testing processes could have improved sequence learning capacity by heightening the sensitivity to statistical probabilities, which is essential for automatic procedural mechanisms (Daw et al. 2005; Janacsek et al. Forthcoming). This interpretation is consistent with the result that participants with better executive functions showed decreased sequence learning in the waking alert condition compared with the participants with lower executive functions, suggesting that in the alert state relying more on attentional processes prevented the learning of statistical contingencies to a greater

extent (Fletcher et al. 2005). In the hypnotic state, participants with higher executive functions shifted from relying on frontal lobe-related attentional processes to automatic procedural-based mechanisms, resulting in enhanced sequence learning. However, future neuroimaging studies need to corroborate these results, providing direct evidence for the underlying brain systems.

In sum, we found that hypnosis substantially boosted procedural-based sequence learning. This result sheds light not only on the competitive nature of brain systems in cognitive processes but also could have important implications for training and rehabilitation programs, especially for developing new methods to improve human skill learning.

Funding

Hungarian Science Foundation (OTKA MB08A 84743).

Notes

Thanks to Gabor Orosz, Zsuzsa Londe, Istvan Winkler for helpful comments on the manuscript. Thanks to Nora Fallah and Szandra Laszlo for their valuable assistance during data collection. *Conflict of interest*: None declared.

References

- Anokhin AP, Golosheykin S, Grant JD, Heath AC. 2010. Developmental and genetic influences on prefrontal function in adolescents: a longitudinal twin study of WCST performance. *Neurosci Lett*. 472:119-122.
- Ashby FG, Alfonso-Reese LA, Turken AU, Waldron EM. 1998. A neuropsychological theory of multiple systems in category learning. *Psychol Rev*. 105:442-481.
- Baldo JV, Schwartz S, Wilkins D, Dronkers NF. 2006. Role of frontal versus temporal cortex in verbal fluency as revealed by voxel-based lesion symptom mapping. *J Int Neuropsychol Soc*. 12:896-900.
- Brown RM, Robertson EM. 2007. Inducing motor skill improvements with a declarative task. *Nat Neurosci*. 10:148-149.
- Daw ND, Niv Y, Dayan P. 2005. Uncertainty-based competition between prefrontal and dorsolateral striatal systems for behavioral control. *Nat Neurosci*. 8:1704-1711.
- Dennis N, Cabeza R. 2010. Age-related dedifferentiation of learning systems: an fMRI study of implicit and explicit learning. *Neurobiol Aging*. 32:2318.e17-2318.e30.
- Egner T, Jamieson G, Gruzeliér J. 2005. Hypnosis decouples cognitive control from conflict monitoring processes of the frontal lobe. *Neuroimage*. 27:969-978.
- Filoteo JV, Lauritzen S, Maddox WT. 2010. Removing the frontal lobes. *Psychol Sci*. 21:415-423.
- Fingelkurts AA, Kallio S, Revonsuo A. 2007. Cortex functional connectivity as a neurophysiological correlate of hypnosis: an EEG case study. *Neuropsychologia*. 45:1452-1462.
- Fletcher PC, Zafiris O, Frith CD, Honey RAE, Corlett PR, Zilles K, Fink GR. 2005. On the benefits of not trying: brain activity and connectivity reflecting the interactions of explicit and implicit sequence learning. *Cereb Cortex*. 15:1002-1015.
- Foerde K, Knowlton BJ, Poldrack RA. 2006. Modulation of competing memory systems by distraction. *Proc Natl Acad Sci U S A*. 103:11778-11783.
- Frank MJ, O'Reilly RC, Curran T. 2006. When memory fails, intuition reigns: midazolam enhances implicit inference in humans. *Psychol Sci*. 17:700-707.
- Fu W-T, Anderson JR. 2008. Solving the credit assignment problem: explicit and implicit learning of action sequences with probabilistic outcomes. *Psychol Res*. 72:321-330.
- Galea JM, Albert NB, Ditye T, Miall RC. 2010. Disruption of the dorsolateral prefrontal cortex facilitates the consolidation of procedural skills. *J Cogn Neurosci*. 22:1158-1164.

- Gruzelier JH. 2006. Frontal functions, connectivity and neural efficiency underpinning hypnosis and hypnotic susceptibility. *Contemp Hypn*. 23:15–32.
- Halsband U. 2006. Learning in trance: functional brain imaging studies and neuropsychology. *J Physiol Paris*. 99:470–482.
- Heaton R, Chelune G, Talley J, Kay G, Curtiss G. 1993. Wisconsin card sorting test manual. Odessa (FL): Psychological Assessment Resources.
- Henke K. 2010. A model for memory systems based on processing modes rather than consciousness. *Nat Rev Neurosci*. 11:523–532.
- Hikosaka O, Nakahara H, Rand MK, Sakai K, Lu X, Nakamura K, Miyachi S, Doya K. 1999. Parallel neural networks for learning sequential procedures. *Trends Neurosci*. 22:464–471.
- Hikosaka O, Nakamura K, Sakai K, Nakahara H. 2002. Central mechanisms of motor skill learning. *Curr Opin Neurobiol*. 12:217–222.
- Howard DV, Howard JH. 2001. When it does hurt to try: adult age differences in the effects of instructions on implicit pattern learning. *Psychon Bull Rev*. 8:798–805.
- Howard DV, Howard JH Jr, Japikse K, DiYanni C, Thompson A, Somborg R. 2004. Implicit sequence learning: effects of level of structure, adult age, and extended practice. *Psychol Aging*. 19:79–92.
- Howard JH Jr, Howard DV. 1997. Age differences in implicit learning of higher-order dependencies in serial patterns. *Psychol Aging*. 12:634–656.
- Janacek K, Fiser J, Nemeth D. Forthcoming. The best time to acquire new skills: age-related differences in implicit sequence learning across human life span. *Dev Sci*.
- Kaiser J, Barker R, Haenschel C, Baldeweg T, Gruzelier JH. 1997. Hypnosis and event-related potential correlates of error processing in a stroop-type paradigm: a test of the frontal hypothesis. *Int J Psychophysiol*. 27:215–222.
- Kallio S, Revonsuo A, Hamalainen H, Markela J, Gruzelier J. 2001. Anterior brain functions and hypnosis: a test of the frontal hypothesis. *Int J Clin Exp Hypn*. 49:95–108.
- Kaufman SB, DeYoung CG, Gray JR, Jiménez L, Brown J, Mackintosh N. 2010. Implicit learning as an ability. *Cognition*. 116:321–340.
- Keele SW, Ivry R, Mayr U, Hazeltine E, Heuer H. 2003. The cognitive and neural architecture of sequence representation. *Psychol Rev*. 110:316–339.
- Kincses T, Johansen-Berg H, Tomassini V, Bosnell R, Matthews P, Beckmann C. 2008. Model-free characterization of brain functional networks for motor sequence learning using fMRI. *Neuroimage*. 39:1950–1958.
- Lieberman MD. 2000. Intuition: a social cognitive neuroscience approach. *Psychol Bull*. 126:109–137.
- Nemeth D, Janacek K. 2011. The dynamics of implicit skill consolidation in young and elderly adults. *J Gerontol B Psychol Sci Soc Sci*. 66:15–22.
- Nemeth D, Janacek K, Csifcsak G, Szvoboda G, Howard JH Jr, Howard DV. 2011. Interference between sentence processing and probabilistic implicit sequence learning. *PLoS One*. 6:e17577.
- Nemeth D, Janacek K, Londe Z, Ullman MT, Howard D, Howard J. 2010. Sleep has no critical role in implicit motor sequence learning in young and old adults. *Exp Brain Res*. 201:351–358.
- Oakley DA, Halligan PW. 2009. Hypnotic suggestion and cognitive neuroscience. *Trends Cogn Sci*. 13:264–270.
- Perruchet P, Pacton S. 2006. Implicit learning and statistical learning: one phenomenon, two approaches. *Trends Cogn Sci*. 10:233–238.
- Poldrack RA, Clark J, Pare-Blagoev EJ, Shohamy D, Creso Moyano J, Myers C, Gluck MA. 2001. Interactive memory systems in the human brain. *Nature*. 414:546–550.
- Poldrack RA, Packard MG. 2003. Competition among multiple memory systems: converging evidence from animal and human brain studies. *Neuropsychologia*. 41:245–251.
- Raz A, Shapiro T. 2002. Hypnosis and neuroscience: a cross talk between clinical and cognitive research. *Arch Gen Psychiatry*. 59:85–90.
- Rieckmann A, Fischer H, Bäckman L. 2010. Activation in striatum and medial temporal lobe during sequence learning in younger and older adults: relations to performance. *Neuroimage*. 50:1303–1312.
- Romano JC, Howard JH, Howard DV. 2010. One-year retention of general and sequence-specific skills in a probabilistic, serial reaction time task. *Memory*. 18:427–441.
- Romano Bergstrom JC, Howard JH Jr, Howard DV. 2011. Enhanced implicit sequence learning in collage-age video game players and musicians. *Appl Cogn Psychol*. 26:91–96.
- Seger CA, Cincotta CM. 2005. Dynamics of frontal, striatal, and hippocampal systems during rule learning. *Cereb Cortex*. 16:1546–1555.
- Shor RE, Orne EC, Press CP. 1962. Harvard group scale of hypnotic susceptibility. Palo Alto, CA: Consulting Psychologists Press.
- Soetens E, Melis A, Notebaert W. 2004. Sequence learning and sequential effects. *Psychol Res*. 69:124–137.
- Song S, Howard JH Jr, Howard DV. 2007. Sleep does not benefit probabilistic motor sequence learning. *J Neurosci*. 27:12475–12483.
- Spreen O, Strauss E. 1991. Language tests. A compendium of neuropsychological tests. p. 268–275.
- Szendi I, Kovacs ZA, Szekeres G, Galsi G, Boda K, Boncz I, Janka Z. 2009. Effects of a hypnotically altered state of consciousness on intensification of semantic processing. *Int J Clin Exp Hypn*. 57:382–401.
- Ullman MT. 2004. Contributions of memory circuits to language: the declarative/procedural model. *Cognition*. 92:231–270.
- Wagstaff GF, Cole JC, Brunas-Wagstaff J. 2007. Effects of hypnotic induction and hypnotic depth on phonemic fluency: a test of the frontal inhibition account of hypnosis. *Int J Psychol Psychol Ther*. 7:27–40.

Appendix IV

Nemeth, D., & Janacsek, K. (2011).

The dynamics of implicit skill consolidation in young and elderly adults.

Journal of Gerontology Series B: Psychological Sciences, 66B(1), 15-22.

DOI: 10.1093/geronb/gbq063.

Impact factor: 2.094.

The Dynamics of Implicit Skill Consolidation in Young and Elderly Adults

Dezso Nemeth* and Karolina Janacsek*

Institute of Psychology, University of Szeged, Szeged, Hungary.

Objectives. Implicit skill learning underlies not only motor but also cognitive and social skills; it is therefore an important aspect of life from infancy to old age. We studied aging effects on the time course of implicit skill consolidation.

Methods. Young and elderly adults performed a probabilistic implicit sequence-learning task before and after a 12-, a 24-hr, and a 1-week interval. The task enabled us to separate the components of skill learning and consolidation: (a) general skill and (b) sequence-specific learning (SSL).

Results. We found improvement of general skill for the young adults in all delay conditions. The elderly adults also showed enhancement after the 12-hr period, revealing brain plasticity similar to young adults. This improvement disappeared in the 24-hr and the 1-week delay conditions. Regarding SSL, no improvement was found in either age group and at either consolidation intervals. In contrast, sequences-specific knowledge decreased in the elderly group independently of the delay.

Discussion. These results draw attention to the fact that consolidation is not a single process, rather there are multiple mechanisms that are differentially affected by time course and by aging.

Key Words: Aging—ASRT—Implicit sequence learning—Memory consolidation—Skill learning.

INTRODUCTION

Skill learning can be differentiated by phases (rapid and slower), modalities, and whether or not it is conscious (implicit and explicit) (Doyon, Bellec, et al., 2009). Implicit skill learning occurs when information is acquired from an environment of complex stimuli without conscious access either to what was learned or to the fact that learning occurred (Reber, 1993). In everyday life, this learning mechanism is crucial for adapting to the environment and evaluate events unconsciously. Implicit skill learning underlies not only motor but also cognitive and social skills (Lieberman, 2000; Ullman, 2004); it is therefore an important aspect of life from infancy to old age. Implicit skills remain essential to healthy functioning with the advancement of age in various contexts, such as social interactions, everyday habits, or reading skills. Social skills appear in compound behaviors realized in proper sequences and under appropriate circumstances. These skills (e.g., communication of emotions) are needed for normal social functioning in various sorts of situations: in the workplace, in the family, in the neighborhood, during recreation, shopping, or in the context of medical and mental care. Furthermore, these skills are crucial for an effective participation in educational, training, and rehabilitation programs. Most models of skill learning (Doyon, Bellec, et al., 2009; Hikosaka et al., 1999; Hikosaka, Nakamura, Sakai, & Nakahara, 2002; Keele, Ivry, Mayr, Hazeltine, & Heuer, 2003; Kincses et al., 2008) emphasize the role of the basal ganglia and the cerebellum. Skill learning does not occur only during practice, in the

so-called “online” periods, but also between practice periods, during the so-called “off-line” periods. The process that occurs during the off-line periods is referred to as consolidation, which means stabilization of a memory trace after the initial acquisition; it can result increased resistance to interference or even improvement in performance following an off-line period (Krakauer & Shadmehr, 2006; Nemeth et al., 2010; Robertson, 2009; Song, 2009). Understanding the time course of skill consolidation can help us reveal the nature of aging memory and age-related changes in brain plasticity.

Recent reviews conclude that off-line changes vary with factors such as the phase and awareness of learning, the formation of contextual associations, the type of information to be learned, as well as the age of the participants (Brashers-Krug, Shadmehr, & Bizzi, 1996; Deroost & Soetens, 2006; Doyon, Korman, et al., 2009; Jiménez, Vaquero, & Lupiáñez, 2006; Keele et al., 2003; Robertson, Pascual-Leone, & Press, 2004; Siengsukon & Boyd, 2008; Song, 2009; Song, Howard, & Howard, 2007). Studies on the time course of skill consolidation indicate that there is a “critical period” after the learning phase, which is necessary for the stabilization of memory traces. This time period depends on the task demand, and it varies from 1 to 2 hr (Robertson, Press, & Pascual-Leone, 2005), to 5 hr (Shadmehr & Brashers-Krug, 1997; Shadmehr & Holcomb, 1997), or 6 hr (Walker, Brakefield, Hobson, & Stickgold, 2003). Using the serial reaction time (SRT) task, which is a widely known sequence-learning paradigm, one study found that the off-line

enhancement increased with the length of delay (Press, Casement, Pascual-Leone, & Robertson, 2005). In this SRT study, no enhancement was found 1 hr after the learning phase, but significant enhancement was observed after 4 hr, which further increased after 12 hr. These results suggest that off-line learning may be a dynamic process. However, this study examined only a shorter stretch of time, so the question can be raised, what happens in skill consolidation after more than 12 hr.

The modified version of SRT is the alternating serial reaction time (ASRT) task (Howard & Howard, 1997), which enables us to separate general skill learning and sequence-specific learning (SSL). In the ASRT task, repeating events alternate with random ones in an eight-element sequence so that the location of every second stimulus in the stream is determined randomly (e.g., 1R2R3R4R, where the numbers represent the repeating events, and *R* represents random stimulus events). This sequence structure has been termed “probabilistic second-order dependency” (Remillard, 2008), because to predict stimulus “*n*,” we need only to remember stimulus *n* – 2 in the sequence, regardless of stimulus *n* – 1. The repeating sequence in the ASRT task is better hidden than in the classical SRT task (e.g., Press et al., 2005), so that the task relies more on implicit mechanisms of learning (Howard & Howard, 1997; Song et al., 2007). Why are the properties of the ASRT interesting in everyday life in aging? The key factor is the implicit nature of the task. In everyday life, we use explicit, conscious, and implicit, nonconscious processes at the same time to do a task (e.g., typing on the computer or learning a foreign language). These processes interact in cooperative and sometimes competitive ways (Brown & Robertson, 2007; Poldrack & Packard, 2003; Song, Marks, Howard, & Howard, 2009) in order to optimize the memory performance (Ullman, 2004). It has long been known that there is age-related decline in explicit memory and executive functions (Cowan, Naveh-Benjamin, Kilb, & Saults, 2006; Craik & Bialystok, 2006; Craik & Salthouse, 2000; Einstein & McDaniel, 1990). It is therefore possible that the efficiency of acquiring new skills decreases in older ages because of these weaker explicit processes and not because of the impairment of pure implicit skill learning. In the ASRT task, explicit memory processes and conscious awareness on sequence learning are almost totally eliminated. That is why it can model implicit learning better than other tasks. So, finding intact learning in the ASRT task could suggest that everyday difficulties regarding skill learning are mainly due to weak explicit processes.

Several studies using the ASRT task demonstrated that older adults show age-related deficits in online skill learning (Curran, 1997; Howard et al., 2004; Howard & Howard, 1997). However, little is known about the effects of aging on skill consolidation and its time course (Siengsukon & Boyd, 2009; Spencer, Gouw, & Ivry, 2007).

Previous studies using the ASRT task focused primarily on whether sleep affects skill consolidation and have concluded that these processes are not influenced by sleep (Nemeth et al., 2010; Song et al., 2007). For example, in a recent ASRT study, Nemeth and colleagues investigated implicit skill learning after 12-hr off-line period. The novelty of this research compared with previous studies of consolidation in older adults (Siengsukon & Boyd, 2009; Spencer et al., 2007) was that (a) it used probabilistic second-order sequences and (b) it dealt separately with general skill and SSL (relatively faster responses to repeated patterns). They showed that general skill learning, as assessed via overall reaction time (RT), improved off-line in both the young and older groups, with the young group improving more than the old group. However, the improvement was not sleep dependent, in that it was not relevant whether the 12-hr off-line period included sleep or not. They did not find sequence-specific off-line improvement in either age group, and similarly to general skill learning, it was not influenced by sleep. However, this study did not examine the consolidation of skills after more than 12 hr in healthy aging.

The current experiment was designed to study the effects of aging on the off-line time course of implicit skill learning. The present study goes beyond Nemeth and colleagues (2010) and other previous studies (Press et al., 2005; Siengsukon & Boyd, 2009; Spencer et al., 2007) by comparing the performance after 12-, 24-hr, and 1-week delays from the initial learning session. We focused on the consolidation of implicit SSL and, separately, general skill learning between young and elderly adults. Despite the results of previous studies that found greater improvement after longer off-line periods (more after 12 hr compared with 4 hr, see Press et al.), it is less plausible that this is true for 12-, 24-hr, and 1-week delays as well. Therefore, we aim to determine a time point in a longer stretch of time at which improvement can still be observed in skill consolidation and analyze its possible age-related differences.

METHODS

Participants

Seventy-one young and 58 elderly right-handed adults participated in the experiment. They were randomly assigned to the 12-, 24-hr, or 1-week delay group (70% of the 12-hr groups were also presented in the study of Nemeth et al., 2010). The age and education of participants were controlled (for detailed data, see Table 1). Participants did not suffer from any developmental, psychiatric, or neurological disorders, did not have sleeping disorders, and all reported having 7–8 hr of sleep a day. Participants were recruited from Szeged and Békéscsaba (Hungary). They gave informed written consent after the aims and procedures of the experiment were explained to them and received no financial compensation for participation.

Table 1. Mean Age, Education, and Sex of Young and Elderly Groups

Group		Age (years)	Education (years)	Sex
Young	12 hr (<i>n</i> = 23)	20.83 (1.11)	13.48 (1.27)	8 men/15 women
	24 hr (<i>n</i> = 31)	21.74 (4.15)	14.77 (2.13)	12 men/ 19 women
	1 week (<i>n</i> = 17)	19.88 (1.27)	13.65 (1.32)	3 men/14 women
Elderly	12 hr (<i>n</i> = 23)	66.43 (6.18)	12.90 (3.26)	7 men/14 women
	24 hr (<i>n</i> = 22)	67.36 (5.30)	13.84 (2.44)	5 men/17 women
	1 week (<i>n</i> = 13)	65.15 (4.14)	14.23 (2.00)	6 men/7 women

Procedure

There were two sessions in the experiment to examine the off-line changes of implicit skill learning: a learning phase (Session 1) and a testing phase (Session 2) separated by a 12-, 24-hr, or 1-week interval off-line period (see Figure 1). Previous studies with similar tasks and experimental designs showed no time of day effect either on general RTs or on learning measures (Nemeth et al., 2010; Press et al., 2005; Robertson et al., 2004; Song et al., 2007); the time of testing was however controlled in our study.

ASRT Task

We used a modification of the original ASRT task (Howard & Howard, 1997) in which a stimulus (a dog head) appeared in one of the four empty circles shown all the time on the screen (Nemeth et al., 2010). Participants were instructed to press one of four possible response keys on an IBM PC keyboard as fast as they could to indicate the location of the stimulus. Each response key (Y, C, B, or M on Hungarian keyboard) was assigned to one of the four stimulus locations on the screen.

During Session 1 (learning phase), the ASRT task consisted of 25 blocks, with 85 stimuli in each block. For practice purposes, the locations of the first five stimuli of each stimulus block were always random. These were followed by the eight-element sequence (e.g., 1R2R3R4R) repeating

10 times. Following the design of Howard and Howard (1997), stimuli were presented 120 ms after the response to the previous stimulus. Between stimulus blocks, the subjects received feedback about their overall RT and accuracy presented on the screen, and then they had a rest period of between 10 and 20 s before starting the next stimulus block. Session 2 (testing phase) consisted of only five stimulus blocks of the same type as in Session 1.

The computer program selected a different ASRT sequence for each subject based on a permutation rule such that each of the six unique permutations of the four repeating events occurred with equal probability (Howard & Howard, 1997; Nemeth et al., 2010). The repeating sequence was identical between Session 1 and Session 2 for each participant.

To explore how much explicit knowledge subjects acquired about the task, we administered a short questionnaire (the same as Song et al., 2007 and Nemeth et al., 2010) after the second session. This questionnaire included increasingly specific questions such as “Have you noticed anything special regarding the task? Have you noticed some regularity in the sequence of stimuli?” The experimenter rated the participants’ answers on a 5-point scale, where 1 meant *Nothing noticed* and 5 meant *Total awareness*. None of the participants reported noticing the repeating sequence of the stimulus locations.

Statistical Analysis

Because in ASRT a repeating sequence alternates with random events, some runs of three consecutive events (termed triplets) occur more frequently than others. For example, with the 1R2R3R4R sequence, 1x2, 2x3, 3x4, and 4x1 (where “x” denotes any location) would occur more often than e.g., 1x3 or 4x2. We refer to the former as “high-frequency” triplets and the latter as “low-frequency” triplets. The triplets including two consecutive repeating-sequence stimuli were always of high-frequency triplets, whereas one-fourth of the triplets that included two consecutive random stimulus events were high-frequency by chance, the rest being low frequency. Of the 64 possible triplets, the 16 high-frequency triplets occurred 62.5% of the time and the 48 low-frequency triplets occurred 37.5% of the time. Note that the final event of high-frequency triplets is therefore more predictable from the initial event compared with the low-frequency triplets. Previous results showed that as participants practice the ASRT task, they come to respond more quickly to the high- than to low-frequency triplets, thus revealing SSL (Howard et al., 2004; Howard & Howard, 1997; Song et al., 2007). Therefore, SSL is reflected in the RT difference between low- and high-frequency triplets (Song, Howard, & Howard, 2008; Song et al., 2007). The larger this SSL score, the greater the SSL is. In addition, general skill learning is revealed in the ASRT task in the improving overall response speed, irrespective of the triplet

Figure 1. The design of the experiment.

types. To facilitate data processing, stimulus blocks were organized into larger clusters (called epochs); where the first epoch contained blocks 1–5, the second epoch blocks 6–10, etc. (Barnes et al., 2008; Bennett, Howard, & Howard, 2007). Consequently, Session 1 consisted of 5 epochs, whereas Session 2 consisted of 1 epoch.

The accuracy of responses remained very high throughout the test (average over 97% for all groups), as is typical (e.g., Howard and Howard, 1997 and Nemeth et al., 2010). Therefore, we analyzed the median RT for correct responses only, calculated separately for high- and low-frequency triplets and for each epoch.

RESULTS

Online Learning—Session 1

To be able to investigate the off-line changes, the learning in Session 1 must be similar in the groups. From this point of view, the end of Session 1 is crucial (Nemeth et al., 2010; Press et al., 2005; Song et al., 2007). Therefore, SSL score was computed for RTs on low- minus high-frequency triplets in Epoch 5 (Song et al., 2007, 2008). The larger this difference, the greater the SSL was. This score was submitted to a univariate analysis of variance (ANOVA) with age group (young versus older) and delay (12, 24 hr, or 1 week) as between-subject factors. The ANOVA revealed significant SSL ($F(1,123) = 136.69$, mean square error (MSE) = 157.95, $\eta_p^2 = .53$, $p < .00001$). Neither the main effects of age group and delay nor the Age group $\times$ Delay interaction was significant (p 's $> .19$); thus, there were no differences in SSL among age groups or delays at the end of Session 1 (see Figure 2A,B).

Regarding the general skill, a univariate ANOVA was conducted on overall RTs of Epoch 5 (collapsed across triplet types) with age group (young versus older) and delay (12, 24 hr, or 1 week) as between-subject factors. The only significant effect was the main effect of age group ($F(1,123) = 409.89$, MSE = 1784.08, $\eta_p^2 = .77$, $p < .001$) reflecting longer RTs for the elderly group (537 ms) compared with the younger participants (381 ms). The main effect of delay and Age group $\times$ Delay interaction did not approach significance (all p 's $> .41$). Thus, participants responded with similar overall RTs in all delay conditions (see Figure 2C,D).

Off-line SSL

To define the index of off-line SSL, we calculated the SSL score at the first epoch of Session 2 similarly to the learning score in Session 1 (Epoch 5). The greater this score at the beginning of the Session 2 compared with the end of Session 1, the larger the off-line SSL was. These two SSL scores were submitted to a mixed-design ANOVA with session (end of Session 1 versus beginning of Session 2) as within-subject factor and age group (young versus elder) and delay (12, 24 hr, or 1 week) as between-subject factors.

Thus, any off-line change of SSL would be revealed by main effects and/or interactions with session.

The main effect of SESSION was not significant ($F(1,123) = 2.28$, MSE = 178.4, $\eta_p^2 = .02$, $p = .13$). However, the ANOVA revealed significant main effect of age group ($F(1,123) = 15.01$, MSE = 194.89, $\eta_p^2 = .11$, $p < .001$) and significant Session $\times$ Age group interaction ($F(1,123) = 5.35$, MSE = 178.4, $\eta_p^2 = .04$, $p = .02$) reflecting that age groups differed in off-line SSL. Sequence-specific knowledge decreased in the elderly groups (–6 ms) compared with the young, who retained the previously acquired skill (1 ms). There was no off-line improvement of sequence-specific knowledge in either group.

Regarding the time course, there were no differences among delay conditions (neither the main effect nor the interactions with delay was significant, p 's $> .38$); thus, the consolidation of sequence-specific knowledge was not affected by the elapsed time between the learning and the testing session (see Figure 2E).

Off-line General Skill Learning

Off-line general skill learning was tested by comparing the overall RTs (collapsed across triplet types) between the last epoch of Session 1 and the first epoch of Session 2 (see Figure 2F). The greater the decrease from Session 1 to Session 2, the larger the off-line general skill learning was. These two variables were submitted to a mixed-design ANOVA with session (end of Session 1 versus beginning of Session 2) as within-subject factor and age group (young versus older) and delay (12, 24 hr, or 1 week) as between-subject factors. Similarly to previous analysis, any off-line change of general skill would be revealed by main effects and/or interactions with session.

This ANOVA revealed significant off-line general skill improvement (indicated by the main effect of session: $F(1,123) = 27.88$, MSE = 399.98, $\eta_p^2 = .19$, $p < .00001$) participants responding faster at the beginning of Session 2 than at the end of Session 1. This off-line improvement was larger for the young group than for the elderly participants (shown by the Session $\times$ Age group interaction: $F(1,123) = 11.45$, MSE = 399.98, $\eta_p^2 = .09$, $p = .001$). In addition, the elapsed time between the two sessions influenced the improvement of general skill improvements as well (indicated by the Session $\times$ Delay interaction: $F(2,123) = 5.29$, MSE = 399.98, $\eta_p^2 = .08$, $p = .006$). Thus, participants' response speed improved more after the 12-hr than after the 24-hr (least significant difference post hoc test: $p = .007$) or 1-week delay ($p = .006$), whereas there was no difference between the 24-hr and 1-week delay conditions ($p = .64$).

The subsequent paired-samples t -tests conducted separately for all age and delay groups revealed that the off-line improvement of general skill was significant in all young groups (all p 's $< .047$), whereas in the elderly groups only the 12-hr delay led to off-line enhancement ($p = .032$).

Figure 2. Sequence-specific learning (SSL) (reaction time [RT] on low- minus high-frequency triplets) at the end of Session 1 and at the beginning of Session 2 for young (A) and elderly groups (B). All groups showed significant SSL by the end of Session 1. Overall RTs are also plotted at the end of Session 1 and at the beginning of Session 2 for young (C) and elderly groups (D). Young participants were generally faster than the old ones. Regarding the off-line SSL (measured by the difference of SSL score at the end of Session 1 and the beginning of Session 2), young groups retained the previously acquired sequence-specific skill, whereas the elderly groups showed decrement in this skill compared with the young group. There were no differences across the 12-, 24-hr, and 1-week conditions (E). Off-line general skill learning (measured by the overall RT changes between the end of Session 1 and the beginning of Session 2) was obtained in all young groups and decreased gradually across delay conditions. Older adults showed off-line improvement only after a 12-hr interval (F). These groups responded significantly faster at the beginning of Session 2 compared with the end of Session 1, whereas elderly participants did not speed up after 24-hr or 1-week delay. Error bars indicate standard errors of mean (SEM).

We also compared the performance of young and elderly groups in all conditions separately. Young adults showed larger off-line improvement than the elderly participants in the 12-hr ($t(44) = 2.46, p = .019$) as well as in the 24-hr delay period ($t(51) = 2.96, p = .0060$), whereas there was no significant difference between groups in the 1-week condition ($t(28) = 0.71, p = 0.44$).

DISCUSSION

We studied the time course of implicit skill consolidation in young and elderly adults with probabilistic second-order regularity sequences (the ASRT task), which enabled us to separate general skill and SSL. In the young adults, we found off-line improvement of the general skill (overall RT)

after the 12-, 24-hr, and 1-week delay as well, with gradual decline among delays. The elderly adults showed off-line improvement of general skill only after the 12-hr off-line period, and this improvement was weaker than that in the young group. Although the pattern in age groups is similar, these results suggest that the off-line course of general skill learning may be affected by aging because we did not find improvement either after 24-hr or 1-week delay in the elderly group. No off-line improvement was found in SSL in either age group with either the 12-, 24-hr, or 1-week consolidation interval. SSL did not decrease significantly between sessions for young participants suggesting that sequence-specific knowledge was well consolidated in this group, whereas the older group showed weaker consolidation in all delay conditions compared with the young. So,

according our results, off-line general skill learning is influenced both by the time course and aging, whereas the off-line sequence learning is affected only by aging.

The significant off-line general skill improvement after the 12-hr delay period is compatible with the results of Song and colleagues (2007) and Nemeth and colleagues (2010). It is also possible that the improvement in overall RTs after the delay period reflects a release from fatigue rather than consolidation per se. However, studies that have included that a fatigue control group (Spencer, Sunm, & Ivry, 2006; Walker, Brakefield, Morgan, Hobson, & Stickgold, 2002) make this interpretation unlikely. The current results confirm those of Song and colleagues and Nemeth and colleagues in finding no evidence of any off-line improvement of sequence-specific performance and extend them to the 24-hr and 1-week consolidation period.

The off-line general skill improvement after 12 hr in the elderly adults contrasts the findings of both Spencer and colleagues (2007) and Siengsukon and Boyd (2009) who obtained no off-line improvement in elderly adults. Due to the different paradigms used in the current and the two previous studies, it is difficult to identify the source of differences. We refer to the point that these previous studies did not distinguish between general skill and SSL, and therefore the signal-to-noise ratio might be reduced making it difficult to detect the off-line improvements for elderly participants.

The differences among the 12-, 24-hr, and 1-week off-line intervals suggest that the consolidation of general skill learning is time dependent. In addition, older participants are more sensitive for this off-line time course in that they showed no off-line improvement even after 24-hr delay. These results are congruent with recent theories of motor skill consolidation (Press et al., 2005; Robertson et al., 2005; Shadmehr & Brashers-Krug, 1997; Walker et al., 2003) that claim that memory stabilization occurs during the first 5–6 hr after learning. The observed strong off-line improvement after 12 hr may reflect this first stabilization process of memory traces, including the previously mentioned critical time period. The differences among 12-, 24-hr, and 1-week consolidation intervals suggest that during new skill acquisition, it could be important to place the training sessions closer to each other for optimal performance, with shorter intervals for elderly participants.

The current results as well as previous findings (Doyon, Korman et al., 2009; Robertson et al., 2004) are compatible with the notion that skill consolidation processes may be different and they could be profoundly dependent on the nature of task demand, such as on the relative proportion of general skill and SSL requirements of the task. Given that these different components of learning are usually not separated in classical sequence learning tasks, off-line improvements in such studies could be falsely attributed to SSL itself. Nevertheless, in the current study consolidation

of sequence-specific information was similar in the 12-, 24-hr, and 1-week off-line periods, with a decline for elderly compared with the young, independently of time course. These results suggest that stabilization of sequence-specific memory is a faster process, whereas off-line changes of general skill are more influenced by a longer stretch of time.

On the functional level, there are at least three mechanisms that may underlie the age-related decline in the consolidation of skill learning: (a) cognitive slowing may hinder elderly adults from having multiple representations simultaneously activated (see simultaneity theory of Salthouse, 1996); (b) associative binding deficits may cause impairment in making associations between multiple stimuli or stimulus features and binding these associations into long-term memory traces (Bennett et al., 2007; Harrison, Duggins, & Friston, 2006); and (c) increased sensitivity to interference also can result weaker stabilization of representations (Park, Smith, Dudley, & Lafronza, 1989). On the neuronal level, age-related decrement was observed both structurally and functionally in the basal ganglia (Chen et al., 2005; Dennis & Cabeza, 2010; Erixon-Lindroth et al., 2005; Raz et al., 2005), that is involved in skill learning. Future studies are still needed to systematically examine the background mechanisms of age-related differences in skill consolidation.

Our findings are compatible with skill learning and consolidation models (Cohen, Pascual-Leone, Press, & Robertson, 2005; Doyon, Bellec et al., 2009; Hikosaka et al., 1999, 2002; Song, 2009; Walker et al., 2003) and draw attention to the fact that the consolidation is not a single process; instead there are multiple mechanisms in off-line learning (general skill, sequence-specific processes), which are differently influenced by time course and by aging. Based on these results, therapists can design more effective educational, training, and rehabilitation programs for age-related disorders.

FUNDING

Bolyai Scholarship Program (to D.N.) and Hungarian National Research Fund (OTKA) K 82068.

ACKNOWLEDGMENTS

Thanks to our mentors: Darlene V. Howard, James H. Howard, Jr., and Michael Ullman from Georgetown University. Thanks to Ildikó Vízi, Maria Tarnai, Krisztina Turay, and Anna Rácz for their valuable assistance during data collection and Gábor Csifcsák for his help. István Winkler and Agnes Szokolszky helped us in the final version of the manuscript.

CORRESPONDENCE

Address correspondence to Dezso Nemeth, PhD, Institute of Psychology, University of Szeged, Egyetem u. 2., 6722 Szeged, Hungary. Email: nemethd@edpsy.u-szeged.hu.

*The authors contributed equally to this work.

REFERENCES

- Barnes, K. A., Howard, J. H., Jr., Howard, D. V., Gilotty, L., Kenworthy, L., Gaillard, W. D., & Vaidya, C. J. (2008). Intact implicit learning of spatial context and temporal sequences in childhood autism spectrum disorder. *Neuropsychology*, 22, 563–570.

- Bennett, I. J., Howard, H. J., & Howard, D. V. (2007). Age-related differences in implicit learning of subtle third-order sequential structure. *Journal of Gerontology: Psychological Sciences*, 62B, 98–103.
- Brashers-Krug, T., Shadmehr, R., & Bizzi, E. (1996). Consolidation in human motor memory. *Nature*, 382, 252–255.
- Brown, R. M., & Robertson, E. M. (2007). Off-line processing: Reciprocal interactions between declarative and procedural memories. *Journal of Neuroscience*, 27, 10468–10475.
- Chen, P. S., Yang, Y. K., Lee, Y. S., Yeh, T. L., Lee, I. H., Chiu, N. T., & Chu, C. L. (2005). Correlation between different memory systems and striatal dopamine D2/D3 receptor density: A single photon emission computed tomography study. *Psychological Medicine*, 35, 197–204.
- Cohen, D., Pascual-Leone, A., Press, D., & Robertson, E. (2005). Off-line learning of motor skill memory: A double dissociation of goal and movement. *Proceedings of the National Academy of Sciences*, 102, 18237–18241.
- Cowan, N., Naveh-Benjamin, M., Kilb, A., & Sauls, J. (2006). Life-span development of visual working memory: When is feature binding difficult? *Developmental Psychology*, 42, 1089–1102.
- Craik, F., & Salthouse, T. (2000). *The handbook of aging and cognition*. Mahwah, NJ: Lawrence Erlbaum Associates.
- Craik, F. I., & Bialystok, E. (2006). Cognition through the lifespan: Mechanisms of change. *Trends in Cognitive Sciences*, 10, 131–138.
- Curran, T. (1997). Effects of aging on implicit sequence learning: Accounting for sequence structure and explicit knowledge. *Psychological Research*, 60, 24–41.
- Dennis, N., & Cabeza, R. (2010). Age-related dedifferentiation of learning systems: An fMRI study of implicit and explicit learning. *Neurobiology of Aging* doi: 10.1016/j.neurobiolaging.2010.04.004.
- Deroost, N., & Soetens, E. (2006). The role of response selection in sequence learning. *The Quarterly Journal of Experimental Psychology*, 59, 449–456.
- Doyon, J., Bellec, P., Amsel, R., Penhune, V., Monchi, O., Carrier, J., Lehericy, S., & Benali, H. (2009). Contributions of the basal ganglia and functionally related brain structures to motor learning. *Behavioural Brain Research*, 199, 61–75.
- Doyon, J., Korman, M., Morin, A., Dostie, V., Hadj Tahar, A., Benali, H., Karni, A., Ungerleider, L. G., & Carrier, J. (2009). Contribution of night and day sleep vs. simple passage of time to the consolidation of motor sequence and visuomotor adaptation learning. *Experimental Brain Research*, 195, 15–26.
- Einstein, G., & McDaniel, M. (1990). Normal aging and prospective memory. *Journal of experimental Psychology: Learning, Memory, and Cognition*, 16, 717–726.
- Erixon-Lindroth, N., Farde, L., Wahlin, T. B., Sovago, J., Halldin, C., & Backman, L. (2005). The role of the striatal dopamine transporter in cognitive aging. *Psychiatry Research*, 138, 1–12.
- Harrison, L., Duggins, A., & Friston, K. (2006). Encoding uncertainty in the hippocampus. *Neural Networks*, 19, 535–546.
- Hikosaka, O., Nakahara, H., Rand, M. K., Sakai, K., Lu, X., Nakamura, K., Miyachi, S., & Doya, K. (1999). Parallel neural networks for learning sequential procedures. *Trends in Neurosciences*, 22, 464–471.
- Hikosaka, O., Nakamura, K., Sakai, K., & Nakahara, H. (2002). Central mechanisms of motor skill learning. *Current Opinion in Neurobiology*, 12, 217–222.
- Howard, D. V., Howard, J. H., Jr., Japikse, K. C., DiYani, C., Thompson, A., & Somberg, R. (2004). Implicit sequence learning: Effects of level of structure, adult age, and extended practice. *Psychology and Aging*, 19(1), 79–92.
- Howard, J. H., Jr., & Howard, D. V. (1997). Age differences in implicit learning of higher-order dependencies in serial patterns. *Psychology and Aging*, 12, 634–656.
- Jiménez, L., Vaquero, J., & Lupiáñez, J. (2006). Qualitative differences between implicit and explicit sequence learning. *Journal of Experimental Psychology. Learning, Memory and Cognition*, 32, 475–489.
- Keele, S. W., Ivry, R., Mayr, U., Hazeltine, E., & Heuer, H. (2003). The cognitive and neural architecture of sequence representation. *Psychological Review*, 110, 316–339.
- Kinches, T., Johansen-Berg, H., Tomassini, V., Bosnell, R., Matthews, P., & Beckmann, C. (2008). Model-free characterization of brain functional networks for motor sequence learning using fMRI. *Neuroimage*, 39, 1950–1958.
- Krakauer, J. W., & Shadmehr, R. (2006). Consolidation of motor memory. *Trends in Neuroscience*, 29, 58–64.
- Lieberman, M. D. (2000). Intuition: A social cognitive neuroscience approach. *Psychological Bulletin*, 126, 109–137.
- Nemeth, D., Janacek, K., Londe, Z., Ullman, M. T., Howard, D., & Howard, J. (2010). Sleep has no critical role in implicit motor sequence learning in young and old adults. *Experimental Brain Research*, 201, 351–358.
- Park, D. C., Smith, A. D., Dudley, W. N., & Lafronza, V. N. (1989). Effects of age and a divided attention task presented during encoding and retrieval on memory. *Journal of Experimental Psychology. Learning, Memory and Cognition*, 15, 1185–1191.
- Poldrack, R. A., & Packard, M. G. (2003). Competition among multiple memory systems: Converging evidence from animal and human brain studies. *Neuropsychologia*, 41, 245–251.
- Press, D. Z., Casement, M. D., Pascual-Leone, A., & Robertson, E. M. (2005). The time course of off-line motor sequence learning. *Brain Research. Cognitive Brain Research*, 25, 375–378.
- Raz, N., Lindenberger, U., Rodrigue, K. M., Kennedy, K. M., Head, D., Williamson, A., Dahle, C., Gerstorf, D., & Acker, J. D. (2005). Regional brain changes in aging healthy adults: general trends, individual differences and modifiers. *Cerebral Cortex*, 15, 1676–1689.
- Reber, A. R. (1993). *Implicit learning and tacit knowledge: An essay on the cognitive unconscious* (Vol. 19). New York: Oxford University Press.
- Remillard, G. (2008). Implicit learning of second-, third-, and fourth-order adjacent and nonadjacent sequential dependencies. *Quarterly Journal of Experimental Psychology (Colchester)*, 61, 400–424.
- Robertson, E. M. (2009). From creation to consolidation: A novel framework for memory processing. *PLoS Biology*, 7, e1000019.
- Robertson, E. M., Pascual-Leone, A., & Press, D. Z. (2004). Awareness modifies the skill-learning benefits of sleep. *Current Biology*, 14, 208–212.
- Robertson, E. M., Press, D. Z., & Pascual-Leone, A. (2005). Off-line learning and the primary motor cortex. *Journal of Neuroscience*, 25, 6372–6378.
- Salthouse, T. A. (1996). The processing-speed theory of adult age differences in cognition. *Psychological Review*, 103, 403–428.
- Shadmehr, R., & Brashers-Krug, T. (1997). Functional stages in the formation of human long-term motor memory. *Journal of Neuroscience*, 17, 409–419.
- Shadmehr, R., & Holcomb, H. H. (1997). Neural correlates of motor memory consolidation. *Science*, 277, 821–825.
- Siengsukon, C. F., & Boyd, L. A. (2008). Sleep enhances implicit motor skill learning in individuals poststroke. *Topics in Stroke Rehabilitation*, 15, 1.
- Siengsukon, C. F., & Boyd, L. A. (2009). Sleep to learn after stroke: implicit and explicit off-line motor learning. *Neuroscience Letters*, 451, 1–5.
- Song, S. (2009). Consciousness and the consolidation of motor learning. *Behavioural Brain Research*, 196, 180–186.
- Song, S., Howard, J. H., Jr., & Howard, D. V. (2007). Sleep does not benefit probabilistic motor sequence learning. *Journal of Neuroscience*, 27, 12475–12483.
- Song, S., Howard, J. H., & Howard, D. V. (2008). Perceptual sequence learning in a serial reaction time task. *Experimental Brain Research*, 189, 145–158.
- Song, S., Marks, B., Howard, J. H., Jr., & Howard, D. V. (2009). Evidence for parallel explicit and implicit sequence learning systems in older adults. *Behavioural Brain Research*, 196, 328.

- Spencer, R. M., Gouw, A. M., & Ivry, R. B. (2007). Age-related decline of sleep-dependent consolidation. *Learning and Memory*, 14, 480–484.
- Spencer, R. M., Sunm, M., & Ivry, R. B. (2006). Sleep-dependent consolidation of contextual learning. *Current Biology*, 16, 1001–1005.
- Ullman, M. T. (2004). Contributions of memory circuits to language: The declarative/procedural model. *Cognition*, 92, 231–270.
- Walker, M. P., Brakefield, T., Hobson, J. A., & Stickgold, R. (2003). Dissociable stages of human memory consolidation and reconsolidation. *Nature*, 425, 616–620.
- Walker, M. P., Brakefield, T., Morgan, A., Hobson, J. A., & Stickgold, R. (2002). Practice with sleep makes perfect: sleep-dependent motor skill learning. *Neuron*, 35, 205–211.