Application of the Hybrid Approach to Chronic Total Coronary Occlusion Percutaneous Coronary Intervention

Ph.D. Thesis

PÉTER TAJTI M.D.

Supervisors: Imre Ungi M.D., Ph.D., Ferenc Tamás Nagy M.D., Ph.D.

Division of Invasive Cardiology

Second Department of Internal Medicine and Cardiology Center

Faculty of Medicine

University of Szeged

Szeged

2019.

TABLE OF CONTENTS

LIST OF PUBLICATIONS.....	3
LIST OF ABBREVIATIONS	11
SUMMARY	12
INTRODUCTION.....	13
CLINICAL BENEFITS OF CTO PCI – RANDOMIZED STUDIES	13
CLINICAL BENEFITS OF CTO PCI – OBSERVATIONAL STUDIES	15
CTO PCI SUCCESS RATES	17
THE HYBRID APPROACH	19
<i>Dual catheter angiography</i>	19
<i>Antegrade Wiring</i>	20
<i>Antegrade Dissection and Re-Entry</i>	20
<i>Retrograde</i>	20
<i>Alteration of crossing strategies</i>	21
AIMS	22
MATERIAL AND METHODS.....	22
DEFINITIONS	22
STATISTICAL ANALYSIS	23
RESULTS	23
CLINICAL CHARACTERISTICS	24
ANGIOGRAPHIC CHARACTERISTICS.....	25
PROCEDURAL OUTCOMES OF THE HYBRID APPROACH	26
DISCUSSION	37
CONCLUSIONS	44
ACKNOWLEDGEMENTS.....	45
REFERENCES	46

LIST OF PUBLICATIONS

This doctoral thesis is based on the following publications:

- I. **Tajti P**, Karmaliotis D, Alaswad K, Jaffer FA, Yeh RW, Patel M, Mahmud E, Choi JW, Burke MN, Doing AH, Dattilo P, Toma C, Smith AJC, Uretsky B, Holper E, Wyman RM, Kandzari DE, Garcia S, Krestyaninov O, Khelinskii D, Koutouzis M, Tsiafoutis I, Moses JW, Lembo NJ, Parikh M, Kirtane AJ, Ali ZA, Doshi D, Rangan BV, Ungi I, Banerjee S, Brilakis ES. The Hybrid Approach of Percutaneous Coronary Interventions for Chronic Total Occlusions: Update from the PROGRESS-CTO (PROspective Global REgiStry for the Study of Chronic Total Occlusion Intervention) International Registry. JACC Cardiovasc Interv. 2018 Jul 23;11(14):1325-1335. *IF: 9.881*
- II. **Tajti P**, Ungi I. Az anterográd technikák szerepe a krónikus teljes koszorúér-elzáródás perkután revaszkularizációjában. Cardiologia Hungarica 2018 November; 48(5):330-339.
- III. **Tajti P**, Abu-Fanne R, Ungi I, Katona A, Sasi V, Nagy FT. Kettős lumenű mikrokatóterek alkalmazása krónikus teljes koronária okklúzió katéteres revaszkularizációjában - egy komplex beavatkozás tanulságai. Cardiologia Hungarica 2019 September; [In Press].

Total impact factor of original papers directly related to the thesis: 9.881

Publications not directly related to the thesis:

- I. **Tajti P**, Alaswad K, Karmaliotis D, Jaffer FA, Yeh RW, Patel M, Mahmud E, Choi JW, Burke MN, Doing AH, Dattilo P, Toma C, Smith AJC, Uretsky BF, Holper E, Potluri S, Wyman RM, Kandzari DE, Garcia S, Krestyaninov O, Khelinskii D, Koutouzis M, Tsiafoutis I, Khatri JJ, Jaber W, Samady H, Jefferson B, Patel T, Abdullah S, Moses JW, Lembo NJ, Parikh M, Kirtane AJ, Ali ZA, Doshi D, Xenogiannis ¹, Stanberry LI, Rangan BV, Ungi I, Banerjee S, Brilakis ES. Procedural Outcomes of Percutaneous Coronary Interventions for Chronic Total Occlusions via the Radial Approach: Insight from an International CTO Registry. JACC Cardiovasc Interv. 2019 Feb 25;12(4):346-358. *IF: 9.881*

- II. **Tajti P**, Karpaliotis D, Alaswad K, Jaffer FA, Yeh RW, Patel M, Mahmud E, Choi JW, Burke MN, Doing AH, Dattilo P, Toma C, Smith AJC, Uretsky BF, Holper E, Potluri S, Wyman RM, Kandzari DE, Garcia S, Krestyaninov O, Khelimskii D, Koutouzis M, Tsiafoutis I, Khatri JJ, Jaber W, Samady H, Moses JW, Lembo NJ, Parikh M, Kirtane AJ, Ali ZA, Doshi D, Xenogiannis I, Stanberry LI, Rangan BV, Ungi I, Banerjee S, Brilakis ES. In-hospital Outcomes of Chronic Total Occlusion Percutaneous Coronary Interventions in Patients with Prior Coronary Artery Bypass Graft Surgery: Insights from an International Multicenter CTO Registry. *Circ Cardiovasc Interv.* 2019 Mar;12(3):e007338. *IF: 6.504*
- III. **Tajti P**, Xenogiannis I, Chavez I, Gössl M, Mooney M, Poulouse A, Sorajja P, Traverse J, Wang Y, Burke MN, Brilakis ES. Expecting the unexpected: preventing and managing the consequences of coronary perforations. *Expert Rev Cardiovasc Ther.* 2018 Nov;16(11):805-814. *IF: 1.67*
- IV. **Tajti P**, Xenogiannis I, Karpaliotis D, Alaswad K, Jaffer FA, Burke MN, Ungi I, Brilakis ES. Chronic total occlusions: update on current tips and tricks. *Curr Cardiol Rep.* 2018 Oct 22;20(12):141. *IF: 1.958*
- V. **Tajti P**, Burke MN, Karpaliotis D, Alaswad K, Jaffer FA, Yeh RW, Patel M, Mahmud E, Choi JW, Doing AH, Dattilo P, Toma C, MD, Smith AJC, Uretsky B, Holper E, Garcia S, Krestyaninov O, Khelimskii D, Koutouzis M, Tsiafoutis I, Moses JW, Lembo NJ, Parikh M, Kirtane AJ, Ali ZA, Doshi D, Jaber W, Samady H, Rangan BV, Xenogiannis I, Ungi I, Banerjee S, Brilakis ES. Prevalence and Outcomes of Percutaneous Coronary Interventions for Ostial Chronic Total Occlusions: Insights from a Multicenter CTO Registry. *Can J Cardiol.* 2018 Oct;34(10):1264-1274. *IF: 4.524*
- VI. **Tajti P**, Brilakis ES. Medical simulation in interventional cardiology: "More research is needed". *Catheter Cardiovasc Interv.* 2018;91:1060-1061.
- VII. **Tajti P**, Karatasakis A, Danek BA, Alaswad K, Karpaliotis D, Jaffer FA, Choi JW, Yeh RW, Patel MP, Mahmud E, Burke MN, Krestyaninov O, Khelimskii D, Toma C, Doing AH, Uretsky B, Koutouzis M, Tsiafoutis I, Wyman RM, Garcia S, Holper E, Xenogiannis I, Rangan BV, Banerjee S, Ungi I, Brilakis ES. In-Hospital Outcomes of Chronic Total

- Occlusion Percutaneous Coronary Intervention in Patients with Chronic Kidney Disease. *J Invasive Cardiol.* 2018 Nov;30(11):E113-E121. *IF: 1.537*
- VIII. **Tajti P**, Alaswad K, Karpaliotis D, Jaffer FA, Yeh RW, Patel MP, Mahmud E, Choi JW, Burke MN, Doing AH, Toma C, Uretsky B, Holper E, Wyman, MR, Kandzari DE, Garcia S, Krestyaninov O, Khelmiskii D, Koutouzis M, Tsiafoutis I, Jaber W, Samady H, Moses JW, Lembo NJ, Parikh M, Kirtane AJ, Ali ZA, Doshi D, Xenogiannis I, Rangan BV, Ungi I, Banerjee S, Brilakis ES. In-hospital Outcomes of Attempting More Than One Chronic Total Coronary Occlusion Via Percutaneous Intervention During the Same Procedure. *Am J Cardiol.* 2018 Aug 1;122(3):381-387. *IF: 3.171*
- IX. **Tajti P**, Burke MN, Karpaliotis D, Alaswad K, Werner GS, Azzalini L, Carlino M, Patel M, Mashayekhi K, Egred M, Krestyaninov O, Khelinskii D, Nicholson WJ, Ungi I, Galassi AR, Banerjee S, Brilakis ES. Update in the Percutaneous Management of Coronary Chronic Total Occlusions. *JACC Cardiovasc Interv.* 2018 Apr 9;11(7):615-625. *IF: 9.881*
- X. **Tajti P**, Brilakis ES. Chronic Total Occlusion Percutaneous Coronary Intervention: Evidence and Controversies. *J Am Heart Assoc.* 2018 Jan 12; 7(2). *IF: 4.45*
- XI. **Tajti P**, Karatasakis A, Karpaliotis D, Alaswad K, Jaffer FA, Yeh RW, Patel M, Mahmud E, Choi JW, Doing AH, Toma C, Uretsky B, Garcia S, Moses JW, Parikh M, Kirtane A, Ali ZA, Hatem R, Karacsonyi J, Danek BA, Rangan BV, Banerjee S, Ungi I, Brilakis ES. Retrograde CTO-PCI of Native Coronary Arteries Via Left Internal Mammary Artery Grafts: Insights From a Multicenter U.S. Registry. *J Invasive Cardiol.* 2018 Mar;30(3):89-96. *IF: 1.537*
- XII. **Tajti P**, Karpaliotis D, Alaswad K, Toma C, Choi JW, Jaffer FA, Doing AH, Patel M, Mahmud E, Uretsky B, Karatasakis A, Karacsonyi J, Danek BA, Rangan BV, Banerjee S, Ungi I, Brilakis ES. Prevalence, Presentation and Treatment of 'Balloon Undilatable' Chronic Total Occlusions: Insights from a Multicenter US Registry. *Catheter Cardiovasc Interv.* 2018 Mar 1; 91(4):657-666. *IF: 2.602*
- XIII. **Tajti P**, Doshi D, Karpaliotis D, Brilakis ES. The "double stingray technique" for recanalizing chronic total occlusions with bifurcation at the distal cap. *Catheter Cardiovasc Interv.* 2018 May 1;91(6):1079-1083. *IF: 2.602*

- XIV. **Tajti P**, Sandoval Y, Brilakis ES. "Around the world" - How to reach native coronary artery lesions through long and tortuous aortocoronary bypass grafts. *Hellenic J Cardiol*. 2018 Jan 3. pii: S1109-9666(17)30607-3. [Epub ahead of print] *IF: 1.89*
- XV. **Tajti P**, Brilakis ES. Percutaneous Treatment of Coronary Perforation in Acutely Occluded Right Coronary Artery after Reimplantation in the Aortic Root. *Hellenic J Cardiol* 2018 May 25. pii: S1109-9666(18)30130-1. [Epub ahead of print] *IF: 1.89*
- XVI. **Tajti P**, Brilakis ES. Does the hybrid algorithm have real impact on long-term outcomes or should only be used as a valuable approach for CTO crossing? *J Thorac Dis*. 2018;10:1320-1324.
- XVII. **Tajti P**, Xenogiannis I, Hall A, Burke MN, Chavez I, Garcia S, Gössl M, Mooney M, Poulose A, Sorajja P, Wang Y, Vemmou E, Nikolakopoulos I, Morley P, Rangan BV, Ungi I, Brilakis ES. Use of the DyeVert System in Chronic Total Occlusion Percutaneous Coronary Intervention. *J Invasive Cardiol* 2019 [In Press] *I.537*
- XVIII. **Tajti P**, Brilakis ES. Retrograde via epicardial collaterals: With power comes responsibility. *J Interv Cardiol* 2018;31:31-32.
- XIX. **Tajti P**, Xenogiannis I, Najjar H, Brilakis ES. Management of Balloon Undilatable CTOs. *Cardiac Interventions Today* May/June 2018.
- XX. Sandoval Y, **Tajti P**, Karatasakis A, Burke MN, Danek BA, Karpaliotis D, Alaswad K, Jaffer FA, Yeh RW, Patel M, Mahmud E, Krestyaninov O, Khelinskii D, Choi JW, Doing AH, Toma C, Wyman RM, Uretsky B, Garcia S, Koutouzis M, Tsiafoutis I, Holper E, Moses JW, Lembo NJ, Parikh M, Kirtane AJ, Ali ZA, Doshi D, Kandzari DE, Karacsonyi J, Rangan BV, Thompson C, Banerjee S, Brilakis ES. Frequency and Outcomes of Ad Hoc Versus Planned Chronic Total Occlusion Percutaneous Coronary Intervention: Multicenter Experience. *J Invasive Cardiol*. 2019 May;31(5):133-139. *IF: 1.537*
- XXI. Xenogiannis I, **Tajti P**, Hall A, Alaswad K, Rinfret S, Nicholson W, Karpaliotis D, Mashayekhi K, Furkalo S, Cavalcante J, Burke MN, Brilakis ES. Update on Cardiac Catheterization in Patients With Prior Coronary Artery Bypass Graft Surgery. *JACC Cardiovasc Interv*. 2019 [In Press]. *IF: 9.881*

- XXII. Xenogiannis I, **Tajti P**, Burke MN, Chavez I, Gössl M, Mooney M, Poulouse A, Sorajja P, Traverse J, Wang Y, Brilakis ES. Coronary Revascularization and Use of Hemodynamic Support in Acute Coronary Syndromes. *Hellenic J Cardiol*. 2019 Jan 21. pii: S1109-9666(18)30490-1. doi: 10.1016/j.hjc.2019.01.010. [Epub ahead of print] *IF: 1.89*
- XXIII. Xenogiannis I, Karpaliotis D, Alaswad K, Jaffer FA, Yeh RW, Patel M, Mahmud E, Choi JW, Burke MN, Doing AH, Dattilo P, Toma C, Smith AJC, Uretsky B, Krestyaninov O, Khelimskii D, Holper E, Potluri S, Wyman RM, Kandzari DE, Garcia S, Koutouzis M, Tsiafoutis I, Jaber W, Samady H, Moses JW, Lembo NJ, Parikh M, Kirtane AJ, Ali ZA, Doshi D, **Tajti P**, Rangan BV, Abdullah S, Banerjee S, Brilakis ES. Comparison Between Traditional and Guide-Catheter Extension Reverse Controlled Antegrade Dissection and Retrograde Tracking: Insights From the PROGRESS-CTO Registry. *J Invasive Cardiol*. 2019 Jan;31(1):27-34. *IF: 1.537*
- XXIV. Xenogiannis I, Karpaliotis D, Alaswad K, Jaffer FA, Yeh RW, Patel M, Mahmud E, Choi JW, Burke MN, Doing AH, Dattilo P, Toma C, Smith AJC, Uretsky B, Krestyaninov O, Khelimskii D, Holper E, Potluri S, Wyman RM, Kandzari DE, Garcia S, Koutouzis M, Tsiafoutis I, Khatri JJ, Jaber W, Samady H, Jefferson BK, Patel T, Moses JW, Lembo NJ, Parikh M, Kirtane AJ, Ali ZA, Doshi D, **Tajti P**, Rangan BV, Abdullah S, Banerjee S, Brilakis ES. Usefulness of Atherectomy in Chronic Total Occlusion Interventions (from the PROGRESS-CTO Registry). *Am J Cardiol*. 2019 May 1;123(9):1422-1428. *IF: 3.171*
- XXV. Xenogiannis I, **Tajti P**, Nicholas Burke M, Brilakis ES. An alternative treatment strategy for large vessel coronary perforations. *Catheter Cardiovasc Interv*. 2019 Mar 1;93(4):635-638. *IF: 2.602*
- XXVI. Xenogiannis I, **Tajti P**, Burke MN, Brilakis ES. Staged revascularization in patients with acute coronary syndromes due to saphenous vein graft failure and chronic total occlusion of the native vessel: A novel concept. *Catheter Cardiovasc Interv*. 2019 Feb 15;93(3):440-444. *IF: 2.602*
- XXVII. Karacsonyi J, **Tajti P**, Rangan BV, Halligan SC, Allen RH, Nicholson WJ, Harvey JE, Spaedy AJ, Jaffer FA, Grantham JA, Salisbury A, Hart AJ, Safley DM, Lombardi WL, Hira R, Don C, McCabe JM, Burke MN, Alaswad K, Koenig GC, Sanghvi KA, Ice D, Kovach RC, Varghese V, Murad B, Baran KW, Resendes E, Martinez-Parachini JR,

- Karatasakis A, Danek BA, Iwnetu R, Roesle M, Khalili H, Banerjee S, Brilakis ES. Randomized Comparison of a CrossBoss First Versus Standard Wire Escalation Strategy for Crossing Coronary Chronic Total Occlusions: The CrossBoss First Trial. JACC Cardiovasc Interv. 2018 Feb 12; 11(3):225-233. *IF: 9.881*
- XXVIII. Xenogiannis I, **Tajti P**, Karpaliotis D, Garbo R, Gagnor A, Burke MN, Brilakis ES. Intravascular imaging for chronic total occlusion intervention. Curr Cardiovasc Imaging Rep (2018) 11: 31.
- XXIX. Xenogiannis I, Karpaliotis D, Alaswad K, Jaffer FA, Yeh RW, Patel M, Mahmud E, Choi JW, Burke MN, Doing AH, Dattilo P, Toma C, Uretsky B, Krestyaninov O, Khelimsii D, Holper E, Potluri S, Wyman RM, Kandzari DE, Garcia S, Koutouzis M, Tsiafoutis I, Khatri JJ, Jaber W, Samady H, Jefferson BK, Patel T, Moses JW, Lembo NJ, Parikh M, Kirtane AJ, Ali ZA, Gkargkoulas F, **Tajti P**, Hall AB, Rangan BV, Abdullah S, Banerjee S, Brilakis ES. Impact of concomitant treatment of non-chronic total occlusion lesions at the time of chronic total occlusion intervention. Int J Cardiol. 2019 Jul 3. [Epub ahead of print] *IF: 4.034*
- XXX. Xenogiannis I, Karpaliotis D, Alaswad K, Basir MB, Yeh RW, Tamez H, Patel M, Mahmud E, Choi JW, Burke MN, Doing AH, Dattilo P, Khatri JJ, Sheikh AM, Malik BA, Greene ME, Abi Rafeh N, Maalouf A, Abou Jaoudeh F, Moses JW, Lembo NJ, Parikh M, Kirtane AJ, Ali ZA, Gkargkoulas F, Russo J, Hakemi E, **Tajti P**, Hall AB, Vemmou E, Nikolakopoulos I, Rangan BV, Abdullah S, Banerjee S, Brilakis ES. Left Main Chronic Total Occlusion Percutaneous Coronary Intervention: A Case Series. J Invasive Cardiol. 2019 Jul;31(7):E220-E225. *IF: 1.537*
- XXXI. Shaukat A, Hryniewicz-Czeneszew K, Sun B, Mudy K, Wilson K, **Tajti P**, Stanberry L, Garberich R, Sandoval Y, Burke MN, Chavez I, Gössl M, Henry T, Lips D, Mooney M, Poulouse A, Sorajja P, Traverse J, Wang Y, Bradley S, Brilakis ES. Outcomes of Extracorporeal Membrane Oxygenation Support for Complex High-Risk Elective Percutaneous Coronary Interventions: A Single-Center Experience and Review of the Literature. J Invasive Cardiol. 2018 Dec;30(12):456-460. *IF: 1.537*
- XXXII. Shaukat A, **Tajti P**, Sandoval Y, Stanberry L, Garberich R, Nicholas Burke M, Gössl M, Henry T, Mooney M, Sorajja P, Traverse J, Bradley SM, Brilakis ES. Incidence, predictors,

management and outcomes of coronary perforations. Catheter Cardiovasc Interv. 2019 Jan 1;93(1):48-56. *IF: 2.602*

- XXXIII. Ungi I, **Tajti P**, Hausinger I, Katona A, Sasi V, Thury A. The retrograde approach to coronary chronic total occlusion interventions. Journal of Continuing Education in Cardiology 2015, 1:10-18.
- XXXIV. Danek BA, Karatasakis A, **Tajti P**, Sandoval Y, Karmpaliotis D, Alaswad K, Jaffer F, Yeh RW, Kandzari DE, Lembo NJ, Patel MP, Mahmud E, Choi JW, Doing AH, Lombardi WL, Wyman RM, Toma C, Garcia S, Moses JW, Kirtane AJ, Hatem R, Ali ZA, Parikh M, Karacsonyi J, Rangan BV, Khalili H, Burke MN, Banerjee S, Brilakis ES. Incidence, Treatment, and Outcomes of Coronary Perforation During Chronic Total Occlusion Percutaneous Coronary Intervention. Am J Cardiol. 2017 Oct 15; 120(8):1285-1292. *IF: 3.171*
- XXXV. Azzalini L, Karatasakis A, Spratt JC, **Tajti P**, Riley RF, Ybarra LF, Schumacher SP, Benincasa S, Bellini B, Candilio L, Mitomo S, Henriksen P, Hidalgo F, Timmers L, Kraaijeveld AO, Agostoni P, Roy J, Ramsay DR, Weaver JC, Knaapen P, Nap A, Starcevic B, Ojeda S, Pan M, Alaswad K, Lombardi WL, Carlino M, Brilakis ES, Colombo A, Rinfret S, Mashayekhi K. Subadventitial stenting around occluded stents: A bailout technique to recanalize in-stent chronic total occlusions. Catheter Cardiovasc Interv. 2018 Sep 1;92(3):466-476. *IF: 2.602*
- XXXVI. Sandoval Y, Lobo AS, Somers VK, Rosenfield K, Bradley SM, Sorajja P, **Tajti P**, Brilakis ES. Sleep deprivation in interventional cardiology: Implications for patient care and physician-health. Catheter Cardiovasc Interv. 2018 Apr 1;91(5):905-910. *IF: 2.602*
- XXXVII. Sandoval Y, Lobo AS, **Tajti P**, Brilakis ES. Laser-assisted orbital or rotational atherectomy: a hybrid treatment strategy for balloon-uncrossable lesions. Hellenic J Cardiol. 2018 Aug 6. [Epub ahead of print] *IF: 1.89*
- XXXVIII. Megaly M, Saad M, **Tajti P**, Burke MN, Chavez I, Gössl M, Lips D, Mooney M, Poulouse A, Sorajja P, Traverse J, Wang Y, Kohl LP, Bradley SM, Brilakis ES. Meta-analysis of the impact of successful chronic total occlusion percutaneous coronary intervention on left

ventricular systolic function and reverse remodeling. J Interv Cardiol. 2018 Oct;31(5):562-571. *IF: 1.728*

- XXXIX. Iverson A, Stanberry LI, **Tajti P**, Garberich R, Antos A, Burke MN, Chavez I, Gössl M, Henry TD, Lips D, Mooney M, Poulouse A, Sorajja P, Traverse J, Wang Y, Bradley S, Brilakis ES. Prevalence, trends, and outcomes of higher risk percutaneous coronary interventions among patients without acute coronary syndromes. Cardiovasc Revasc Med. 2019 Apr;20(4):289-292. *IF: 0.15*
- XL. Najjar H, **Tajti P**, Xenogiannis L, Brilakis E. CTO complexity begets complexity. Cath Lab Digest. 2018 May; 26(5):1, 18-22.
- XLI. Vemmou E, Nikolakopoulos I, Xenogiannis I, **Tajti P**, Hall A, Brilakis ES. The Gordian Knot-If You Cannot Solve it, Cut it: Treating Guidewire Bending While Obtaining Arterial Access. JACC Cardiovasc Interv. 2019 May 13;12(9):892-893.

Total impact factor of original papers not related to the thesis: 110.588

LIST OF ABBREVIATIONS

ADR – antegrade dissection and re-entry

AK – air kerma

AWE – antegrade wire escalation

BMI – body mass index

CABG – coronary artery bypass graft

CAD – coronary artery disease

CART – controlled antegrade and retrograde subintimal tracking

CCS – Canadian Cardiovascular Society Angina Classification

CI – confidence interval

cMRI – cardiac magnetic resonance imaging

CPX – cardiopulmonary exercise

CTO – chronic total occlusion

CVD – cerebrovascular disease

DES – drug eluting stent

DV – donor vessel

HF – heart failure

J-CTO – Japanese Chronic Total Occlusion

LAD – left anterior descending artery

LCX –left circumflex artery

LM – left main

LVEF – left ventricular ejection fraction

MACE – major adverse cardiac event

MI – myocardial infarction

MIB – myocardial ischemic burden

OMT – optimal medical treatment

OR – odds ratio

PVD –peripheral arterial disease

PCI – percutaneous coronary intervention

QoL – quality of life

RCA –right coronary artery

SAQ – Seattle Angina Questionnaire

SWT – segmental wall thickening

TVR – target vessel revascularization

SUMMARY

Percutaneous coronary intervention (PCI) for chronic total occlusion (CTO) continues to evolve constantly with developing equipment and techniques. The hybrid approach to CTO PCI advocates dual coronary injection, careful and structured review of the angiogram, and flexibility. Use of all crossing strategies [antegrade wire escalation, antegrade dissection re-entry and retrograde approach] is encouraged, with initial and subsequent choices influenced by the CTO anatomic characteristics and the outcomes of the originally selected approach. Application of the hybrid approach to CTO PCI has been associated with good outcomes in US and European registries, although CTO PCI outcomes in non-selected populations have been less optimal with approximately 60% success rate.

We analyzed the clinical, angiographic, and procedural characteristics of 3,122 CTO PCIs performed in 3,055 patients to determine the techniques and outcomes of hybrid CTO PCI in a diverse group of patients and operators in two continents (at eighteen US, one European, and one Russian centers) enrolled in the PROGRESS-CTO (**PRO**spective **G**lobal **RE**giStry for the Study of **Ch**ronic **T**otal **O**ccclusion Intervention, NCT02061436) registry between January 2012 and November 2017.

Technical success rate was 87% and the risk for in-hospital major complications was 3%, providing important benchmarks to use when discussing with patients and providers the risk/benefit ratio of CTO PCI. The final successful crossing strategy was antegrade wire escalation in 52%, retrograde in 27%, and antegrade dissection reentry in 21%; more than 1 crossing strategies were required in 41% of the cases.

CTO PCI is currently being performed with high success and acceptable complication rates among various experienced centers in the US and Europe. Bridging the gap between what is currently achieved and what can be achieved in chronic total occlusion intervention should be a major focus of upcoming research and education efforts.

INTRODUCTION

Coronary chronic total occlusions (CTOs) are defined as 100% occlusions with TIMI 0 flow with at least 3 month duration (1). Treatment options for patients with coronary CTOs include lifestyle changes, medications (as is appropriate for all patients with coronary artery disease) and coronary revascularization with either percutaneous coronary intervention (PCI) or coronary bypass graft surgery (CABG). Chronic total occlusion (CTO) percutaneous coronary intervention (PCI) is a rapidly evolving area of interventional cardiology that has undergone a remarkable technical improvement over the past two decades.

CTOs are found in 15-25% of patients undergoing coronary angiography who are found to have coronary artery disease (2-4) in various large cohorts. In a more recent study from SCAAR (Swedish Coronary Angiography and Angioplasty Registry), the prevalence of CTO among patients with at least one 50% luminal coronary stenosis was 16.1% (14441 of 89872 patients) (5). In a Canadian single center registry the prevalence of a CTO was 20%, however PCI was only performed in 9% of these patients whereas majority of them had CABG (34%) or were treated with medical therapy alone (57%) (6). Treatment of CTO with PCI currently remains less preferred in clinical practice possibly due to the limited operator expertise, high procedural cost, limited cardiac catheterization laboratory time availability, high risk for procedural complications and the ongoing controversy regarding its clinical benefits.

In the previous version of the appropriateness use criteria for coronary revascularization (7) revascularization recommendations were different for patients with and without a coronary CTO. This is no longer the case in the current (2016 and 2017) versions (8,9). The novel ESC/EACTS 2018 guidelines recommends Class II a (level of evidence B) for percutaneous CTO revascularization: “*CTO PCI should be considered in patients with angina resistant to medical therapy or with a large area of documented ischemia in the territory of the occluded vessel*”(10).

Clinical benefits of CTO PCI – randomized studies

Currently, symptom improvement is considered the main benefit of CTO PCI, despite criticisms that there is limited supportive prospective randomized-controlled clinical trial data. Indeed, only five randomized-controlled trials have been published (11-15).

The Evaluating Xience and Left Ventricular Function in Percutaneous Coronary Intervention on OccLusiOns After ST-Elevation Myocardial Infarction trial (**EXPLORE**;

NTR1108) enrolled 304 patients who underwent primary PCI for acute ST-segment elevation acute myocardial infarction and had a coexisting non-infarct related artery CTO. Patients were randomized to CTO PCI vs. medical therapy alone. CTO PCI success was 73%. Cardiac magnetic resonance imaging (cMRI) performed after 4 months showed similar left ventricular ejection fraction and left ventricular end-diastolic volume in the two study groups (11).

The **R**andomiz**E**d Trial to Assess Regional Left Ventricular Function After Stent Implantation in Chronic Total Occlusion trial (**REVASC**; NCT01924962) sought to investigate the impact of successful CTO PCI on left ventricular function. The study randomized 205 patients to CTO PCI or non-CTO PCI arms (OMT-only) with a primary endpoint of segmental wall thickening (SWT), change in left ventricular volume (LVV) and ejection fraction (LVEF) assessed with cMRI performed at baseline and at 6 months. The trial presented similar change in SWT, regional and global left ventricular function both in CTO and non-CTO PCI groups. Subgroup analysis, however, showed a significant improvement in SWT amongst patients with a CTO without major non-CTO lesion. In hospital major adverse cardiac event (MACE) rate was also significantly higher in the OMT group (16.3% vs. 5.9%, $p=0.02$) (14). The limitation of the study includes the relatively high baseline LVEF in both study arms (PCI group: 59.6% [IQR 45.8 to 64.3] 54.7% [IQR 42.9 to 65.1], $p=0.48$) and it was underpowered to identify hard clinical endpoints such as cardiac death or MI.

The **D**rug-**E**luting stent Implantation versus optimal Medi**C**al Treatment In patients with Chron**I**c Total Occlu**S**ION (**DECISION-CTO**; NCT01078051) trial randomized 834 patients with coronary CTOs (many of whom also had multivessel disease) to OMT (optimal medical therapy) alone vs. OMT + CTO-PCI. Patients in the OMT and the OMT + CTO PCI group had similar clinical outcomes during a median follow-up of 3.1 years (primary study endpoint was the composite of death, myocardial infarction, stroke or any revascularization). The study has several limitations, such as suboptimal primary endpoint selection, high rate non-CTO PCI (73% of the study patients had multivessel disease in both groups), early termination before achievement of target enrollment, high cross-over rates (18% in the OMT alone group underwent CTO PCI), and mild baseline symptoms in both study groups (12).

The Randomized Multicenter Trial to **E**valuate the Utilization of **R**evascularization or **O**ptimal Medical Therapy for the Treatment of **C**hronic **T**otal Coronary **O**clusions trial (**EUROCTO**; NCT01760083) aimed to randomize 1,200 patients to either CTO PCI versus OMT-

only groups with a primary endpoint of health status change assessed by the Seattle Angina Questionnaire (SAQ) at 12 months of follow-up. Due to slow enrollment the study ended prematurely after randomizing 407 patients instead of the 1,200 population. In contrast to DECISION-CTO, non-CTO lesions were treated before enrollment in the study. Compared with patients randomized to medical therapy only, patients randomized to CTO PCI had more improvement in angina frequency (5.23 OR, CI 95% 1.75 to 8.71, $p=0.003$), and quality of life (QoL) (6.62 OR, CI 95% 1.78 to 11.46, $p=0.007$) (13).

The **Impact on Inducible Myocardial Ischemia of Percutaneous Coronary Intervention versus Optimal Medical Therapy in Patients with Right Coronary Artery Chronic Total Occlusion (IMPACTOR-CTO)** randomized patients with exclusively right coronary artery (RCA) CTOs to PCI ($n=39$) or OMT-only ($n=33$) groups with endpoints of change in myocardial ischemic burden (MIB) (primary endpoint, assessed with cMRI), and 6 minute walking test (6MWT), quality of life (QoL) and MACE (secondary endpoints) at 12 months of follow-up. In the PCI group, myocardial ischemic burden showed significant decrease ($27.7\pm 8.5\%$ to $16.1\pm 8.6\%$, $p<0.01$), in comparison with the OMT group ($28.4\pm 6.1\%$ vs. 27.0 ± 8.0 , $p=0.83$) without significant change. Functional status and QoL also significantly improved in PCI group but not the OMT group.

Although randomized-controlled clinical trials are the gold standard for determining the efficacy and safety of an intervention, only 15% of the current American College of Cardiology/American Heart Association guideline recommendations are based on level A evidence (16). Therefore, in the majority of cases, clinical decision making relies on less robust clinical evidence, such as retrospective and observational studies.

Clinical benefits of CTO PCI – observational studies

Several observational, uncontrolled studies have suggested clinical benefit with CTO PCI, by improving angina, dyspnea, depression, exercise capacity, and risk for arrhythmias.

The OPEN-CTO (Outcomes, Patient Health Status, and Efficiency in Chronic Total Occlusion Hybrid Procedures) registry analyzed 1,000 consecutive patients undergoing CTO PCI with the hybrid approach using standardized questionnaires. A 10.8 (95% confidence interval [CI]: 6.3 to 15.3) point improvement in the QoL domain of the Seattle Angina Questionnaire (SAQ) was observed among successful vs. unsuccessful procedures ($p<0.001$) (17).

Similar results have been shown in multiple prior studies and meta-analyses, that have also reported lower mortality among successful vs. failed CTO PCIs (18). Several studies have assessed the long-term outcomes of CTO PCI as compared with medical therapy, reporting lower incidence of major adverse cardiac events with CTO PCI (19,20), even among patients with well-developed collateral circulation (21). Several other retrospective studies have compared the outcomes of CTO PCI with medical therapy. In an Italian multicenter registry of 1,777 patients with CTOs treatment was as follows: PCI (43.7%), MT (medical therapy, 46.5%), or surgery (9.8%). At 1-year follow-up, cardiac death (1.4% vs. 4.7% and 6.3%, $p<0.001$) and major adverse cardiac events (MACE, 2.6% vs. 8.2% and 6.9%, $p<0.001$) were significantly lower in the PCI group, than in MT or CABG group (19). In propensity-matched analysis MT was associated with higher MACE rate, death, and rehospitalization (19).

Patients with CTOs often have depression that improves after successful CTO PCI (22). CTO PCI also increases exercise capacity with longer six minute walking distance (417 ± 126 vs. 463 ± 103 m, $p=0.002$) in one study, likely due to lower angina frequency (39% vs. 8% $p<0.001$) and less ischemia, especially in patients with larger baseline ischemic burden (23). Two studies performed cardiopulmonary testing before and after CTO PCI, showing increased peak oxygen consumption and anaerobic threshold (24,25). CTOs may be associated with higher arrhythmic risk: patients with prior myocardial infarction and a CTO had greater area of scar tissue (34 cm^2 vs. 19 cm^2 , $P = 0.001$) and higher frequency of recurrent ventricular tachycardia after ablation during a median follow up of 19 months (47% vs. 16%, $p=0.003$) (26).

A recent meta-analysis focused on comparing CTO PCI to OMT including only the available randomized controlled trials (RCT) and propensity score matched (PSM) observational studies to date. A pooled analysis of eight studies (4 RCT and 4 PSM studies) with a total number of 3,971 patients were included (PCI arm [$n=2,050$] and OMT arm [$n=1,921$]). CTO PCI was associated with more favorable outcomes in terms of cardiac death (0.52 OR, CI 95% 0.01 to 0.81, $p<0.01$) – mostly driven by the PSM studies, however, there were no difference in terms of overall MACE, re-PCI or acute MI (27).

CTO PCI improves patient symptoms, whereas there is limited, retrospective data on whether it can impact the subsequent incidence of death, myocardial infarction and arrhythmias. Accordingly, the key indication for offering and performing CTO PCI should be the alleviation of symptoms.

CTO PCI Success Rates

Achieving clinical benefit with CTO PCI requires the procedure to be successful. With contemporary equipment and techniques (such as the hybrid algorithm (28)), high success rates (85-90%) are achieved at experienced centers (29-35). However, success rates in unselected populations remain low: 61.3% in the New York State PCI Registry (36) and 59% in the National Cardiovascular Data Registry in the US (vs. 96% in non-occlusive lesions, $p < 0.001$) (37). Therefore, there is a gap between what is achieved at dedicated CTO PCI centers and the outcomes at less experienced centers.

Patient-specific estimation of the likelihood of success can be facilitated by using various scoring systems. The first CTO PCI score was the J-CTO score (Multicenter CTO Registry of Japan) that was developed to estimate the likelihood of successful guidewire crossing within 30 minutes based on 5 criteria (intralesional $>45^\circ$ bend, length >20 mm, calcification, blunt stump, and previously failed attempt) (38). The J-CTO score has been validated in other CTO PCI cohorts (39) and is also associated with 1-year and 5-year clinical outcomes (40,41).

Other scores include the PROGRESS-CTO (**PRO**spective **G**lobal **RE**giStrY for the **S**tudy of **C**hronic **T**otal **O**clusion Intervention) score (42), the RECHARGE (**RE**gistry of **C**rossboss and **H**ybrid Procedures in **Fr**Ance, the **Ne**the**R**lands, **Be**l**G**ium and **Unit**Ed Kingdom) registry score (43), the CL-score (**C**linical and **L**esion related score) (44), EuroCTO/CASTLE score (**C**ABG history, **A**ge, **S**tump anatomy, **T**ortuosity degree, **L**ength of occlusion, **E**xtent of calcification) (45), and the ORA (**o**stial location, collateral filling of **R**entrop <2 , **a**ge over 75) score (46). Ellis et al. used a novel approach for developing a scoring system for predicting CTO PCI technical success by stratifying lesions according to proximal cap ambiguity. The Ellis-score is currently the only one that also takes into account on operators' experience with CTO PCI (47). One study showed that various scores had similar predictive capacity for technical success and that they performed better in antegrade-only cases (48).

In general, CTO PCI should be performed when the anticipated benefits (which depend on the patient's baseline clinical condition and the likelihood of success) exceed the potential short- and long-term risks (**Figure 1**) (1,49).

Figure 1 Overview of the Potential Risks and Benefits of CTO PCI

Parameters that can help determine the risks and benefits of chronic total occlusion percutaneous coronary intervention.

CABG, coronary artery bypass grafting; **CTO**, chronic total occlusion; **MI**, myocardial infarction; **PCI**, percutaneous coronary intervention.

Reproduced from reference (49).

The Hybrid Approach

The hybrid approach to chronic total occlusion (CTO) percutaneous coronary intervention (PCI) (**Figure 2**) advocates dual coronary injection, careful and structured review of the angiogram, and flexibility (50). Use of all crossing strategies (**Figure 3** [antegrade wire escalation (51), antegrade dissection reentry (52-55) and the retrograde approach (56-59)]) is encouraged (50), with initial and subsequent choices influenced by the CTO anatomic characteristics and the outcomes of the originally selected approach (29,30,34,50,60-64).

Figure 2 Hybrid algorithm for CTO PCI.

Dual angiography is performed and 4 parameters are assessed: based on them the initial crossing strategy is selected, followed by early change in case of failure to achieve progress with any other strategy.

CART: controlled antegrade and retrograde subintimal tracking.

Reproduced from reference (49).

Dual catheter angiography

The first and one of the most important steps of CTO PCI is to perform dual catheter angiography (bilateral angiography or dual coronary injection). Dual injection provides the most appropriate visualization of the proximal, distal portion of the target vessel and the collateral circulation, optimizing the selection of the most suitable initial crossing strategy. Dual injection can also assist in clarifying the current location of the guidewire(s) during crossing attempts, however it may increase the contrast utilization throughout CTO PCI (microcatheter tip injections may replace it). Based upon the diagnostic angiography, pre-procedural meticulous review should be done for at least 10 to 15 minutes focusing on four key anatomic parameters ('the anatomy

dictates' (28)): (a) proximal cap; (b) lesion length; (c) quality of the distal vessel; and (d) the presence of collaterals amenable for retrograde approach ('interventional collaterals') (1). Dual angiography optimally starts with donor vessel injection followed by injection of the occluded vessel 2 to 3 seconds later (or even triple injections in prior CABG patients), without magnification of the image or panning, using long acquisitions or even various views. Microcatheter tip injections can facilitate distal visualization if only ipsilateral epicardial collateral connection presents as can provide more optimal view than dual angiography.

Antegrade Wiring

Antegrade wire escalation (AWE) indicates the use of guidewires of increasing stiffness to cross a CTO. This technique is the most efficient in lesions with shorter occlusion length (<20mm), less tortuous vessel segments, and in occlusions with microchannels. Antegrade wire escalation technique is the most basic and widely used CTO PCI crossing approach, hence familiarity and confidence with antegrade wire escalation serves as the foundation of all other advanced CTO PCI techniques (65). In current practice AWE has four key steps: (a) approaching the CTO; (b) crossing the proximal cap; (c) navigation within the occluded segment; and (d) distal cap penetration (1).

Antegrade Dissection and Re-Entry

Antegrade dissection and re-entry are usually applied for long lesions that are approached in antegrade fashion. The subintimal space should be approached by creating a proximal dissection with advancing a "knuckle" formed at the tip of a polymer jacketed guidewire or by using CrossBoss microcatheter. Knuckle wiring minimizes the risk for perforation and also allows atraumatic subintimal crossing following the course of the vessel. The CrossBoss system is mostly recommended for in-stent CTOs due to safety purposes (preventing inadvertent wiring outside the stent struts) and cost effectiveness (66). Re-entry should be performed with the Stingray system (1,67).

Retrograde

The retrograde approach can be used upfront (primary) or after failed antegrade attempt. It has four key technical steps including (a) advancement of a guidewire and microcatheter through a collateral vessel or a bypass graft distal to the occlusion, (b) crossing the occlusion, (c) wire externalization and (d) balloon angioplasty with stent implantation. Upfront retrograde crossing in

CTOs are commonly used with proximal cap ambiguity, poor distal target, bifurcation at distal cap as well as after failure of antegrade crossing. In case of short lesions (<20 mm) retrograde true lumen puncture is recommended, whereas longer lesions (≥ 20 mm) should be approached with the reverse controlled antegrade and retrograde subintimal tracking (reverse CART) technique (1).

Figure 3 Summary of major crossing techniques in CTO PCI.

The four key crossing techniques in CTO PCI practice (* Retrograde indicates both retrograde wire escalation and retrograde dissection re-entry techniques).

ADR, antegrade dissection and re-entry; AWE, antegrade wire escalation.

Alteration of crossing strategies

Switching between different CTO PCI techniques is considered as the ‘heart’ of the hybrid algorithm. If one technique fails, another approach should be attempted. The general concept is to spend no more than 5-10 minutes in a stagnant state that may increase the risk for procedural failure due to excessive utilization of limited resources such as radiation, contrast, or time. Minor (wire tip re-shaping, change of guidewire with different crossing features), or major (change from an antegrade to a retrograde approach or vice versa) technical adjustments should be made throughout CTO PCI avoiding the stationary state, hence becoming a hybrid operator requires high level of expertise and comfortability with all CTO crossing strategies and devices (1).

AIMS

- (i) We sought to examine a contemporary, multicenter CTO PCI registry to determine the techniques and outcomes of CTO PCI.
- (ii) Our goal was to validate the hybrid algorithm in a diverse group of patients and operators in two continents.
- (iii) We aimed to identify areas in need for improvement in the field of CTO PCI.

MATERIAL AND METHODS

We analyzed the clinical, angiographic, and procedural characteristics of 3,122 CTO PCIs performed in 3,055 patients enrolled in the PROGRESS-CTO (Prospective Global Registry for the Study of Chronic Total Occlusion Intervention, NCT02061436) registry between January 2012 and November 2017 at eighteen US, one European, and one Russian centers. The study was approved by the institutional review board (IRB) of each center.

Definitions

Coronary CTOs were defined as coronary lesions with Thrombolysis In Myocardial Infarction (TIMI) grade 0 flow of at least 3 months duration. Estimation of the duration of occlusion was clinical, based on the first onset of angina, prior history of myocardial infarction in the target vessel territory, or comparison with a prior angiogram. Calcification was assessed by angiography as mild (spots), moderate (involving $\leq 50\%$ of the reference lesion diameter) and severe (involving $> 50\%$ of the reference lesion diameter). Moderate proximal vessel tortuosity was defined as the presence of at least 2 bends $> 70^\circ$ or 1 bend $> 90^\circ$ and severe tortuosity as 2 bends $> 90^\circ$ or 1 bend $> 120^\circ$ in the CTO vessel. Blunt or no stump was defined as lack of tapering or lack of a funnel shape at the proximal cap. Interventional collaterals were defined as collaterals considered amenable to crossing by a guidewire and a microcatheter by the operator. A procedure was defined as “retrograde” if an attempt was made to cross the lesion through a collateral vessel or bypass graft supplying the target vessel distal to the lesion; if not, the procedure was classified as “antegrade-only”. Antegrade dissection/re-entry was defined as antegrade PCI during which a guidewire was intentionally introduced into the subintimal space proximal to the lesion, or re-entry into the distal true lumen was attempted following intentional or inadvertent subintimal guidewire crossing.

Technical success was defined as successful CTO revascularization with achievement of <30% residual diameter stenosis within the treated segment and restoration of TIMI grade 3 antegrade flow. Procedural success was defined as the achievement of technical success without any in-hospital complications. In-hospital major adverse cardiac events (MACE) included any of the following adverse events prior to hospital discharge: death, myocardial infarction, recurrent symptoms requiring urgent repeat target vessel revascularization with PCI or coronary artery bypass graft surgery (CABG), tamponade requiring either pericardiocentesis or surgery, and stroke. Myocardial infarction (MI) was defined using the Third Universal Definition of Myocardial Infarction (type 4a MI) (68). Major bleeding was defined as bleeding causing reduction in hemoglobin >3 g/dl or bleeding requiring transfusion or surgical intervention. The J-CTO score was calculated as described by Morino et al (69), the PROGRESS-CTO score as described by Christopoulos et al (42), and the PROGRESS-CTO Complications score as described by Danek et al (70).

Statistical Analysis

Categorical variables were expressed as percentages and were compared using Pearson's chi-square test or Fisher's exact test. Continuous variables were presented as mean $\pm$ standard deviation or median [interquartile range, IQR] unless otherwise specified and were compared using the t-test and 1-way analysis of variance (ANOVA) for normally distributed variables; the Wilcoxon rank-sum test, and the Kruskal-Wallis test were applied for non-parametric continuous variables, as appropriate. Multivariable logistic regression with stepwise backward elimination was performed to examine the independent association between annual CTO PCI volume and procedural outcomes (procedural success and in-hospital MACE). Variables with univariable association in the present study ($p < 0.05$) were entered into the model, as well as variables that have been previously linked with procedural outcomes of CTO PCI: such variables included age, smoking, peripheral arterial disease, chronic lung disease, history of MI, stroke, PCI or CABG, left ventricular ejection fraction, CTO target vessel, multiple CTO vessel treated during the same procedure, and CTO PCI only. All statistical analyses were performed with JMP 13.0 (SAS Institute, Cary, North Carolina). A two-sided p value of 0.05 was considered statistically significant.

RESULTS

Clinical characteristics

The baseline clinical features of the study population are summarized in **Table 1**. As compared with patients in whom CTO PCI failed, patient in whom CTO PCI was successful were younger and less likely to be men, and to have hypertension. They were also less likely to have had a myocardial infarction, congestive heart failure, prior coronary artery bypass graft surgery, and prior PCI along with higher left ventricular ejection fraction. Most patients (89%) were symptomatic having at least Class 2 Canadian Cardiovascular Society (CCS) Angina Classification level (CCS 2: 25%; CCS 3: 54%; CCS 4: 10%), and mostly had stable (64%) or unstable (18%) angina.

Table 1. Clinical characteristics of the study population classified according to technical success.

Clinical characteristics	Overall (n=3055)	Technical success (n=2657)	Technical failure (n=398)	p value
Age (years) *	64.80 ± 10.09	64.6 ± 10.15	66.01 ± 9.63	0.0141
Male gender	85.25%	84.69%	88.95%	0.0378
BMI (kg/m ²) *	30.60 ± 6.14	30.50 ± 6.15	31.20 ± 6.02	0.0666
Smoking (current)	26.01%	25.37%	30.27%	0.0561
Diabetes	43.02%	43.53%	39.65%	0.1758
Dyslipidemia	92.20%	92.11%	92.75%	0.6781
Hypertension	90.26%	89.61%	94.49%	0.0044
Family history of CAD	33.35%	33.44%	32.80%	0.8423
CCS Angina Classification				0.4771
Class ≤1	11.44%	11.64%	10.15%	
Class 2≤	88.56%	88.36%	89.85%	
Myocardial viability performed	24.99%	24.28%	29.32%	0.0783
Prior MI	46.00%	44.82%	53.75%	0.0023
Heart failure	30.56%	29.71%	36.25%	0.0159
Prior valve surgery or procedure	3.17%	3.06%	3.89%	0.4210
Prior PCI	65.29%	64.49%	70.62%	0.0180
Prior CABG	32.49%	31.28%	40.68%	0.0003

Baseline creatinine (mg/dL) †	1.01 (0.89, 1.22)	1.01 (0.89, 1.21)	1.07 (0.90, 1.27)	0.1301
Currently on dialysis	2.67%	2.50%	3.80%	0.1633
Prior CVD	11.70%	11.51%	12.90%	0.4567
Prior PAD	15.02%	14.53%	18.29%	0.0709
Chronic lung disease	14.20%	13.80%	16.81%	0.1386
Left ventricular EF (%) †	54 (42, 60)	55 (44, 60)	50 (40, 60)	0.0357

*mean ± standard deviation; † median (interquartile range)

BMI, body mass index; **CABG**, coronary artery bypass graft; **CAD**, coronary artery disease; **CVD**, cerebrovascular disease; **EF**, ejection fraction; **MI**, myocardial infarction; **PCI**, percutaneous coronary intervention; **PVD**, peripheral arterial disease.

Angiographic characteristics

The angiographic characteristics of the study lesions are presented in **Table 2**. The CTO target lesions were located in the right coronary artery (55%), left anterior descending artery (24%) and left circumflex artery (20%). Failed CTO PCI was associated with longer lesion length (37.8 ± 24.0 mm vs. 33.4 ± 24.1 mm, $p=0.0030$), proximal cap ambiguity (54% vs. 32%, $p<0.0001$), moderate to severe calcification (73% vs. 52%, $p<0.0001$) and tortuosity (45% vs. 33%, $p<0.0001$), or previously failed recanalization attempt (27% vs. 19%, $p=0.0005$). Failed CTO PCI cases also had higher J-CTO score (3.07 ± 1.13 vs. 2.34 ± 1.29 , $p<0.0001$), PROGRESS-CTO score (1.77 ± 1.13 vs. 1.25 ± 1.01 , $p<0.0001$), and PROGRESS-CTO complications score (3.54 ± 1.97 vs. 3.00 ± 1.91 , $p<0.0001$).

Table 2. Angiographic characteristics of study lesions, classified according to technical success.

Angiographic characteristics	Overall (3122)	Technical success (n=2711)	Technical failure (n=411)	p value
Target vessel				0.0640
RCA	55.22%	54.93%	57.14%	
LAD	23.81%	24.57%	18.80%	
LCX	19.91%	19.47%	22.81%	
Other	1.06%	1.03%	1.25%	

CTO length (mm)*	33.99±24.16	33.43±24.14	37.80±23.99	0.0030
Vessel diameter (mm)*	2.85±0.51	2.86±0.51	2.81±0.47	0.1383
Proximal cap ambiguity	35.06%	31.98%	53.97%	<0.0001
Side branch at proximal cap	49.91%	47.56%	64.14%	<0.0001
Blunt stump/no stump	53.69%	50.61%	72.55%	<0.0001
Interventional collaterals	56.72%	58.80%	44.19%	<0.0001
Moderate/severe calcification	54.23%	52.30%	67.02%	<0.0001
Moderate/severe tortuosity	34.96%	33.43%	45.21%	<0.0001
In-stent restenosis	16.61%	16.13%	19.68%	0.0878
Previously failed CTO PCI	20.20%	19.21%	26.70%	0.0005
J-CTO score *	2.43±1.30	2.34±1.29	3.07±1.13	<0.0001
PROGRESS-CTO score *	1.32±1.03	1.25±1.01	1.77±1.01	<0.0001
PROGRESS-CTO complication score *	3.07±1.93	3.00±1.91	3.54±1.97	<0.0001

*mean ± standard deviation

CTO, chronic total occlusion; **J**, Japan; **LAD**, left anterior descending artery; **LCX**, left circumflex artery; **LM**, left main segment; **PCI**, percutaneous coronary intervention; **PROGRESS-CTO**, Prospective Global Registry of Chronic Total Occlusion Interventions; **RCA**, right coronary artery.

Procedural outcomes of the hybrid approach

Overall technical and procedural success was 87% and 85% respectively and the in-hospital major complications rate was 3.0%. The baseline technical and procedural characteristics are presented in **Tables 3, 4** and **5**. Antegrade wire escalation was the most commonly used initial approach (in 75%), especially in lower complexity CTOs (J-CTO score: 2.28±1.29, PROGRESS CTO score: 1.35±1.05), while antegrade dissection/reentry (8%; J-CTO score: 2.86±1.16, PROGRESS CTO score: 1.50±1.07) and the retrograde approach (16%; J-CTO score: 3.12±1.07, PROGRESS CTO score: 1.33±0.96) were used in more complex lesions ($p<0.0001$). The initial approach was successful in 55% of the cases, while 41% of the cases underwent further attempts that were technically successful in 79% (**Figure 4**).

Table 3. Techniques used for CTO PCI in the study patients.

Technical characteristics	Overall (3122)	Technical success (n=2711)	Technical failure (n=411)	p value
Crossing strategies used				
AWE	81.77%	81.15%	85.89%	0.0204
ADR	31.68%	29.69%	44.77%	<0.0001
Retrograde	38.57%	35.96%	55.72%	<0.0001
First crossing strategy				0.0124
AWE	75.36%	75.80%	72.51%	
ADR	8.39%	8.67%	6.57%	
Retrograde	16.24%	15.53%	20.92%	
Final crossing strategy				<0.0001
AWE	45.89%	51.95%	5.24%	
ADR	18.95%	20.96%	5.49%	
Retrograde	23.97%	27.09%	2.99%	
None	11.19%	0.00%	86.28%	
Balloon uncrossable lesions	10.62%	10.21%	29.37%	<0.0001
Balloon undilatable lesions	11.11%	10.74%	22.22%	0.0349
Access site				
Right femoral	78.96%	78.46%	82.24%	0.0798
Left femoral	54.29%	53.34%	60.58%	0.0060
Right radial	32.48%	33.12%	28.22%	0.0481
Left radial	18.67%	19.48%	13.38%	0.0031
Bifemoral approach	51.35%	50.42%	57.42%	0.0082
Biradial approach	14.09%	14.90%	8.76%	0.0009

ADR, antegrade dissection reentry; AWE, antegrade wire escalation.

Figure 4 Application of the hybrid approach in the study patients

Sequence of CTO crossing techniques used in the study patients and associated technical success.

CTO, chronic total occlusion; PCI, percutaneous coronary intervention.

The final successful crossing strategy was antegrade wire escalation (46%), antegrade dissection reentry (19%), and the retrograde approach (24%). The success of antegrade wire escalation decreased with lesion complexity (**Figure 5**), as classified with the J-CTO score (easy [J-CTO 0]: 88%; intermediate [J-CTO 1]: 72%; difficult [J-CTO 2]: 51% and very difficult [J-CTO ≥ 3]: 32% to 17%; $p < 0.0001$), and the PROGRESS-CTO score (55%, 43%, 42%, 39% and 43%, respectively for scores of 0, 1, 2, 3, and 4; $p < 0.0001$).

Figure 5 Technical success and crossing strategy utilization according to the J-CTO score.

Impact of CTO lesion complexity on technical success and use of various crossing strategies.

ADR, antegrade dissection reentry; AWE, antegrade wire escalation; CTO, chronic total occlusion; J, Japan.

The retrograde approach was more commonly required in complex lesions, as classified by the J-CTO score (3%, 9%, 20%, 35% to 44%, respectively for J-CTO scores of 0, 1, 2, and ≥ 3 , $p < 0.0001$), but less frequently in lesions as classified according to the PROGRESS-CTO score (23%, 27%, 23%, 15%, 6%, $p < 0.0001$) (**Figure 6**).

Figure 6 Successful crossing strategy utilization according to the PROGRESS-CTO score.

The impact of retrograde technique utilization is less potent with increasing CTO lesion complexity.

ADR, antegrade dissection reentry; **AWE**, antegrade wire escalation; **CTO**, chronic total occlusion; **PROGRESS-CTO**, PROspective Global REgistry for the Study of Chronic Total Occlusion Intervention.

Dual injection was used in 70% of all cases, and was more frequent in failed interventions (76% vs. 70%, $p=0.026$) and in complex lesions with high J-CTO scores (48% vs. 78%, $p<0.0001$). Radial access was used in 37% overall, with biradial approach in 14%, and in combination with femoral approach in 20% of the cases. Use of radial access was lower with increasing lesion complexity (easy: 50%, intermediate: 39%, difficult: 36%, very difficult: 38%, $p=0.003$), whereas the frequency of femoral (63%, 78%, 83%, 84%, $p<0.0001$) and bifemoral approach (28%, 43%, 51%, 57%, $p<0.0001$) increased with increasing lesion complexity. Median contrast volume, air kerma radiation dose, procedure and fluoroscopy time were 270 (200-360) ml, 2.9 (1.7-4.7) Gray, 123 (81-188) and 47.0 (28.6-77.0) minutes, respectively, and were higher in more complex lesions.

Less complex lesions required fewer stents ($p<0.0001$), however the frequency of non-CTO PCI was higher in those groups, and decreased with increasing lesion complexity ($p=0.08$).

Table 4. Procedural characteristics, classified according to PCI outcomes.

Procedural outcomes	Overall (n=3055)	Technical success (n=2657)	Technical failure (n=398)	p value
Dual injection	70.41%	69.57%	75.82%	0.0260
Radial access ‡	37.06%	37.70%	32.85%	0.0577
Femoral access	81.90%	81.37%	85.40%	0.0480
Procedure time (min) *	123 (81, 188)	121 (80, 184)	140 (85, 224)	<0.0003
Contrast volume (mL) *	270 (200, 360)	260 (200, 350)	300 (220, 400)	<0.0001
Fluoroscopy time (min) *	47.0 (28.6, 77.0)	45.0 (27.3, 73.7)	66.0 (39.0, 93.6)	<0.0001
Patient AK dose (Gray) *	2.9 (1.7, 4.7)	2.8 (1.6, 4.5)	3.9 (2.4, 6.0)	<0.0001
Number of stents †§	2.4±1.1	2.4±1.1	2.7±1.6	0.5559
Stent length (mm) †§	71.8±36.4	71.7±36.3	78.6±47.6	0.6599
Non-CTO lesion PCI	28.20%	28.39%	26.80%	0.5611
In-hospital MACE	3.04%	2.37%	7.54%	<0.0001

* median (interquartile range)

† mean ± standard deviation

‡ Radial access indicates any radial access site use in CTO PCI; including biradial and combined radial-femoral approaches

§ In successful cases stents were implanted in 97.10% vs. in failed procedures in 4.89% (related to perforation, investment procedure, donor vessel dissection, or stenting but 3≥ Thrombolysis In Myocardial Infarction flow grade), $p<0.0001$.

AK, air kerma; **PCI**, percutaneous coronary intervention; **MACE**: major adverse cardiac event.

Table 5. Procedural characteristics, classified according to PCI outcomes and lesion complexity (J-CTO score).

Procedural outcomes	Easy (0)	Intermediate (1)	Difficult (2)	Very difficult (3±)	p for trend
Dual injection	47.59%	64.19%	68.70%	77.89%	<0.0001
Radial access ‡	49.59%	39.09%	35.83%	38.17%	0.0030
Femoral access	63.27%	78.17%	82.65%	84.09%	<0.0001
Procedure time (min) *	77 (46, 117)	92 (58, 136)	115 (73, 172)	152 (102, 217)	<0.0001
Contrast volume (mL) *	220 (160, 300)	250 (190, 340)	260 (200, 350)	280 (202, 385)	<0.0001
Fluoroscopy time (min) *	27 (18, 39)	33 (19, 54)	42 (27, 66)	63 (39, 93)	<0.0001
Patient AK dose (Gray) *	1.8 (0.9, 3.3)	2.1 (1.2, 3.5)	2.5 (1.4, 4.4)	3.5 (2.1, 5.2)	<0.0001
Number of stents †§	1.8±0.9	2.0±1.0	2.3±1.1	2.8±1.2	<0.0001
Stent length (mm) †§	48.0±25.2	57.6±31.1	66.5±33.5	85.5±36.3	<0.0001
Non-CTO lesion PCI	32.75%	32.00%	29.19%	26.37%	0.0786
In-hospital MACE	1.36%	1.41%	3.01%	3.11%	0.0119
Technical success	96.90%	94.84%	89.14%	81.26%	<0.0001

* median (interquartile range)

† mean ± standard deviation

‡ Radial access indicates any radial access site use in CTO PCI; including biradial and combined radial-femoral approaches

AK, air kerma; **PCI**, percutaneous coronary intervention; **MACE**: major adverse cardiac event.

The procedural success and annual CTO PCI volume in the participating sites are shown in **Figure 7**.

Figure 7 Procedural success rate and annual CTO PCI volume in the participating centers.

Procedural success rate significantly differed in the participating sites (p for trend <0.0001) with increasing annual CTO PCI volume.

CTO, chronic total occlusion; **PCI**, percutaneous coronary intervention.

Higher median annually performed CTO PCI per center was associated with higher procedural success on both univariable and multivariable analysis (**Figure 8**) but not in-hospital MACE.

Figure 8 Baseline clinical and angiographic characteristics and procedural success.

Multivariable analysis of parameters associated with procedural success.

CI, confidence interval, **CVD**, cerebrovascular disease; **CTO**, chronic total occlusion; **LAD**, left anterior descending; **HF**, heart failure; **MI**, myocardial infarction; **OR**, odds ratio, **PCI**, percutaneous coronary intervention.

The incidence of in-hospital MACE was 3.04% [death (0.85%), acute myocardial infarction (1.08%), stroke (0.26%), emergency CABG (0.16%), urgent re-PCI (0.36%), and pericardial tamponade (0.85%)], and increased with increasing lesion complexity (**Table 5**). The prevalence of in-hospital MACE (major adverse cardiac event) were higher in failed procedures (7.54% vs. 2.37%, $p<0.0001$) (**Table 6**) and with more complex crossing techniques: antegrade

wire escalation, antegrade dissection reentry or retrograde crossing (1.09% vs. 2.96% vs. 5.61%, $p<0.0001$). Use of the retrograde approach was associated with higher incidence of complications, as shown in **Figure 9**. Median length of hospital stay was significantly higher in patients with vs. without in-hospital MACE (6 [2-9] days vs. 1 [1-2] days), $p<0.0001$).

Table 6. Procedural complications in the study patients.

Variable	Overall (n=3055)	Technical success (n=2657)	Technical failure (n=398)	p value
In-hospital MACE	3.04%	2.37%	7.54%	<i><0.0001</i>
Death	0.85%	0.68%	2.01%	<i>0.0143</i>
Acute MI	1.08%	0.79%	3.02%	<i>0.0005</i>
Stroke	0.26%	0.30%	0.00%	0.6072
Re-PCI	0.36%	0.38%	0.25%	1.0000
Emergency CABG	0.16%	0.08%	0.75%	<i>0.0179</i>
Pericardiocentesis	0.85%	0.53%	3.02%	<i><0.0001</i>
Perforation	4.48%	3.54%	10.80%	<i><0.0001</i>
Vascular access complication	1.44%	1.39%	1.76%	0.5674
Equipment loss	0.23%	0.19%	0.50%	0.2288
DV dissection/thrombosis	1.15%	1.05%	1.76%	0.2081
Bleeding	1.18%	1.20%	1.01%	1.0000
Aortocoronary dissection	0.10%	0.08%	0.25%	0.3422
Contrast induced nephropathy *	0.36%	0.38%	0.25%	1.0000

CABG, coronary artery bypass graft surgery; **DV**, donor vessel; **PCI**, percutaneous coronary intervention; **MACE**, major adverse cardiac event; **MI**, myocardial infarction.

* Contrast induced nephropathy is defined as a relative increase of $\geq 25\%$ or absolute increase of ≥ 0.5 mg/dL in pre-PCI serum creatinine at 48 h after PCI.

Figure 9 In-hospital major complications classified according to final successful crossing strategy.

Use of the retrograde approach was associated with higher overall in-hospital MACE and risk for perforation.

ADR, antegrade dissection and reentry; AWE, antegrade wire escalation; CABG, coronary artery bypass graft; MACE, major adverse cardiac event; MI, myocardial infarction; PCI, percutaneous coronary intervention.

DISCUSSION

To the best of our knowledge this is the largest study reported to date on CTO PCI using the hybrid approach, demonstrating high technical success rates (87%) with acceptable major complication rate (3.04%). These outcomes were achieved despite high lesion complexity and relatively low success of the initially selected CTO crossing strategy (55%) (**Figure 4**).

Previous smaller studies have provided similarly encouraging results. In an earlier report from the PROGRESS-CTO registry from 11 US centers, technical success was 91% and in-hospital MACE was 1.7% (29). Wilson et al reported following 79% initial success rate among 1156 patients from 7 centers enrolled in the UK Hybrid CTO Registry, with 90% final technical success rate after repeat procedures (30). The RECHARGE registry of 1253 CTO interventions performed in 1177 patients between 2014 and 2015 at 22 European centers, reported 86% procedural success rate and 2.6% major in-hospital complication rate (34). Vo et al reported a single operator pilot-study showing rapid increase in procedural success despite the low initial CTO-PCI experience (71). At two high-volume, experienced, centers Pershad et al showed significant increase in technical (79.4% to 95.4%, $p<0.001$) and procedural (77.9% to 88.3%) success rates after implementation of the hybrid algorithm, as compared with the pre-implementation period (62). Furthermore, in the Hybrid Video Registry analyzed 194 video-taped live case demonstrations reporting high success rates (92.8%), even in highly complex CTOs with acceptable procedure time and contrast volume (72).

In the present study we found that technical and procedural success remained high with reasonably low complication rates despite expansion of the registry in recent years. Antegrade wire escalation was more commonly applied as the initial crossing approach (74%) in less complex lesions (J-CTO score: 2.24 ± 1.24 , PROGRESS-CTO score: 1.32 ± 0.87) and was the most common final crossing strategy (in approximately half the cases). Antegrade dissection reentry and retrograde techniques were more likely to be used as initial strategy in cases with complex anatomy (J-CTO score 2.78 ± 1.21 and 3.32 ± 0.98 , respectively; PROGRESS-CTO score 1.38 ± 0.93 and 2.00 ± 0.89 , respectively), and were the final successful strategy in 22% and 28% of all cases.

Failure to cross with a guidewire was the most common reason for CTO PCI failure (in 86%). In 13% the procedure failed despite successful guidewire crossing (**Table 3**) due to balloon undilatable lesions (3.9%), inability to deliver stents (2.3%), final TIMI flow <3 (1.3%), residual

stenosis >30% (1.0%), and procedure-related complications (0.8%, 1 patient with donor vessel thrombosis, 1 patient with aortocoronary dissection, and 1 procedure related death due to pericardial tamponade and subsequent cardiogenic shock). The presence of balloon uncrossable (29% vs. 10%, $p<0.0001$), and undilatable lesions (22% vs. 11, $p=0.0109$) was higher in the failed CTO PCI group, highlighting the need for CTO PCI operators to have experience in treating these and other complex lesions, such as severe calcification and bifurcations (73-76).

As shown in prior studies higher annual CTO PCI volume was independently associated with higher success rates, reflecting the importance of center and operator experience in optimizing outcomes, even among complex lesion and patient subgroups (77). Our study confirms that annual CTO PCI volume significantly improves procedural success amongst various operators and lesion subsets (1.21 OR, CI 95%, 1.13 to 1.29, $p<0.0001$) (**Figure 8**). Attempting 2 or more CTO PCI during the same procedure (0.38 OR, CI 95% 0.18 to 0.77, $p=0.0074$) lowers the likelihood of procedural success, that is currently not recommended due to high potential risk for compromising larger myocardial territory that may result in higher procedure-related in-hospital mortality (78).

Based upon both univariate and multivariate analysis, several anatomic parameters were found to have significant impact on decreasing procedural success (calcification, tortuosity, proximal cap ambiguity) that have been linked with CTO PCI failure in prior studies (35,79,80), and also were implemented into various scores. Nevertheless, distal occlusion anatomy also determines procedural outcomes which is well represented in our study: bifurcation at distal cap decreases [0.62 OR, CI 95% 0.46 to 0.82, $p=0.0011$], and conversely, appropriate distal vessel landing zone increases the likelihood of success (1.40 OR, CI 95% 1.03 to 1.91, $p=0.0340$).

Furthermore, the prevalence of prior CABG was 32%, which is more common compared to European (15-22% (30,31,34)) or Japanese (8-12% (33,81)) registries. PCI in patients with prior bypass graft surgery is especially challenging due to multiple comorbidities, more complex coronary anatomy, and bypass graft failures requiring multiple re-interventions either in the grafts or native coronary arteries (commonly requiring CTO PCI) (82). Creative solutions, and comfortability with all available CTO PCI techniques are required in such challenging patient population (**Figure 10 and 11**) (83). Optimization of CTO PCI techniques, however, in post-bypass patients should be one of the major focus points of upcoming research as procedural success remains lower compared with patients without prior CABG (84).

Figure 10 Clinical case, illustrating the challenges of percutaneous coronary intervention (PCI) in a bypassed right coronary artery (RCA) chronic total occlusion (CTO). Successful CTO PCI was achieved with retrograde application of dual lumen microcatheter in a patient with Canadian Cardiovascular Society grade 2-3 angina and two prior failed CTO attempts and a prior coronary artery bypass graft surgery .

Panel A: Dual angiography showing RCA CTO with proximal blunt stump, distal cap at bifurcation and retrograde filling of SVG at the crux (yellow arrow). **Panel B:** Retrograde microcatheter tip injection reveals distal anatomy showing the SVG graft (yellow arrow) as well. **Panel C:** Successful wiring to distal cap, but failed distal cap puncture (wire advances only to SVG graft) **Panel D:** Retrograde dual lumen microcatheter (Sasuke [Asahi Intecc, Nagoya, Japan]) was used for distal cap puncturing via the over-the-wire (OTW) port (green arrowhead). **Panel E:** Antegrade and retrograde wire knuckling. **Panel F:** Multiple failed reverse CART (controlled antegrade and retrograde subintimal tracking) with increasing balloon sizes (red arrowheads). **Panel G:** Successful GuideLiner (Vacular Solution, Minneapolis, MN, USA) assisted reverse CART (green arrowheads). **Panel H:** Final result after 3 DES implantation.

Reproduced from reference (83)

Figure 11 Clinical case, illustrating the importance of changing crossing strategy to successfully recanalize a challenging chronic total occlusion. Percutaneous coronary intervention (PCI) of a distal right coronary artery (RCA) chronic total occlusion (CTO) in a patient with Canadian Cardiovascular Society grade 3 angina and four prior failed CTO attempts and two prior coronary artery bypass graft surgeries.

Panel A-B: Triple coronary injection showing distal RCA CTO immediately distal to the SVG anastomotic site. **Panel C:** Multiple attempts for septal collateral crossing (surfing and contrast guided techniques) succeeded in advancing a guidewire to the distal RCA, but failed microcatheter advancement. **Panel D:** Successful contrast guided epicardial collateral crossing (Suoh 03, Caravel microcatheter [both Asahi Intecc., Nagoya, Japan]). **Panel E:** Several attempts to advance the retrograde guidewire distal to the SVG anastomosis failed.

Panel F: Multiple attempts to recanalize the native RCA failed, as the antegrade guidewire could not be advanced distal to the SVG anastomosis. **Panel G:** The reverse controlled antegrade and retrograde subintimal tracking (reverse CART) was successfully used to cross the CTO and advance the retrograde guidewire into the SVG-RCA, followed by guidewire externalization. **Panel H:** Final angiographic result after stent implantation. The patient had significant symptom improvement.

The overall complication rate was 3.04%, and occurred less frequently in technically successful procedures (2.37% vs. 7.54%, $p<0.0001$). The risk for complications was higher in more complex lesions (easy [J-CTO score 0] 1.36% vs. very difficult [J-CTO score $3\leq$] 3.11%, $p=0.01$) and with use of advanced crossing techniques (which were more commonly used in more complex lesions). This highlights the importance of weighing the risks and benefits of the procedure, both during discussions with patients and family (to determine whether CTO PCI should be done), but also during the procedure itself: implementing more complex CTO crossing strategies (such as retrograde crossing via epicardial collaterals) may predispose to increased risk for complications, which may be justified in some patients due to significant potential benefit, but not in some others. Accordingly, adjustment and improvement of advanced CTO crossing techniques – such as antegrade dissection re-entry and retrograde approach – may further improve procedural outcomes in more challenging patient populations and complex lesion subsets.

Despite the encouraging findings from our study and other contemporary registries, the success rates of CTO interventions in unselected patient cohorts remain low. Hannan et al. (New York State PCI Registry, $n=4030$ patients) (36) reported 61.3% success rate with 1.07% complication rates (vs. 1.06% for non-CTO PCI cases, $p=0.95$). Ramunddal et al. showed similarly low procedural success rate (54.2%) among 6,442 patients undergoing CTO intervention in the Swedish Coronary Angiography and Angioplasty Registry (SCAAR) (85). Habara et al. compared the acute outcomes of 3,229 CTO interventions among 56 high and low volume centers in Japan showing higher overall success rate in high-volume centers (90.6% vs. 85.6% $p<0.001$), without significant difference in in-hospital MACE rates (0.45% vs. 0.62% $p=0.35$), mostly attributed to higher antegrade success rate (33). Hence, CTO PCI should be performed by experienced operators at dedicated centers in order to achieve optimal results. The CTO PCI success rates in all comer registries (54%-80%) are significantly lower than those achieved at experienced centers (85%-90%) (**Table 7**).

Table 7 Procedural outcomes of multicenter CTO and general PCI registries in recent years.

Authors	Center (n)	Case number (n)	Technical success	Procedural success	Overall MACE
DEDICATED CTO PCI REGISTRIES					
Christopoulos et al. (29) *	11	1,036	91%	90%	1.7%
Suzuki et al. (35) †	41	2,846	90%	89%	2.0%
Wilson et al. (30) *	7	1,156	90%	-	1.6%
Maeremans et al. (34) *	17	1,253	89%	86%	2.6%
Sapontis et al. (17) *	12	1,000	86%	85%	7.0%
Tajti et al. (86) *	20	3,122	87%	85%	3.0%
ALL COMER CTO PCI REGISTRIES					
Brilakis et al. (37)	-	22,365	-	59%	1.6%
Hannan et al. (36)	61	4,030	-	61%	1.1%
Ramunddal et al. (5)	-	6,442	-	54%	-
Tsai et al. (87)	79	2,394	80%	80%	4.3%

* Dedicated CTO PCI registries using the hybrid algorithm.

† 41 highly skilled Japanese operators (Japanese CTO-PCI Expert Registry).

MACE, major adverse cardiovascular event; **MI**, myocardial infarction.

Sharma et al. presented that procedural outcomes of CTO interventions among operators who had received proctorship in using the hybrid approach was higher (77.5% vs. 62.1% $p<0.0001$), especially in more complex cases (if J-CTO score ≥ 2 , the corresponding rates were 70.7% vs. 49.5%, respectively, $p=0.0003$) than those who did not receive mentorship (88).

Given the complexity of CTO PCI, becoming a successful CTO operator requires appropriate training and continued practice (**Figure 12**). Optimal training for CTO PCI remains controversial, as there are only few dedicated fellowship programs. Most operators learn CTO PCI after being in practice for a few years through participation in courses and proctoring. Developing experience in CTO PCI is important, both for achieving high success rates but also for minimizing the risk for complications and efficiently managing them if they occur (36,37).

Figure 12 The four stages of learning CTO PCI.

Experience in performing CTO interventions is associated with application of advanced techniques in more complex patient and lesion subsets.

CTO, chronic total occlusion; **PCI**, percutaneous coronary intervention.

Reproduced from reference (65).

Bridging the gap will likely require development of novel equipment and techniques as well as development of comprehensive, high-volume (77) CTO PCI programs (89) and continued education through live case demonstrations (90), online educational contents (such as case libraries – <https://www.ctomanual.org>, <https://www.pcimanual.org>; or online sharing of novel research achievements – <https://www.progresscto.org>), CTO PCI workshops and proctorships (88). CTO PCI is one of the key components of the growing complex, high-risk, PCI programs (91,92).

Our study has limitations. First, we did not have mid- and long-term follow up of the study patients. Second, there was no core laboratory assessment of the study angiograms or independent clinical event adjudication. Third, the procedures were performed in dedicated, high volume CTO centers by experienced operators, limiting the extrapolation to less experienced operators in low volume centers. Fourth, some centers only enrolled patients during part of the study period due to participation in other studies. Fifth, procedure related myocardial infarction events are site reported

without systematic post-procedural biomarker assessment, hence the rate of such complications may be underestimated.

CONCLUSIONS

- (i) Application of the hybrid approach resulted in high technical success rates (87%) and acceptable rate of major in-hospital complications (3%) across a large number of sites and operators in the US, Europe, and Russia.
- (ii) Changing crossing strategy was required in 41% of cases with the final success strategy being antegrade wire escalation in 52%, retrograde in 27%, and antegrade dissection reentry in 21%.
- (iii) Bridging the gap between what is currently achieved (50-60%) and what can be achieved (85-90%) in chronic total occlusion intervention should be a major focus of upcoming research and education efforts.
- (iv) CTO PCI can currently be achieved with high success and acceptable complication rates among various operators and patient populations in the US and Europe, highlighting the need for developing more CTO PCI centers of excellence in order to achieve the best possible clinical outcomes in this challenging patient and lesion group.

ACKNOWLEDGEMENTS

I would like to express my deepest gratitude to those who have supported me during my scientific work.

Imre Ungi, Associate Professor and Head of the Division of Invasive Cardiology Second Department of Internal Medicine and Cardiology Center. Thank you for your continuous support as a mentor, both in my clinical and research work. You gave me a secure background since medical school, and throughout my fellowships in Hungary and the United States.

Ferenc T. Nagy, Senior Lecturer, Division of Invasive Cardiology Second Department of Internal Medicine and Cardiology Center. Thank you for your valuable advices and management both in my clinical practice and research work.

Emmanouil S. Brilakis, Associate Professor of Medicine, University of Texas Southwestern Medical School and Director of the Center for Complex Coronary Interventions at the Minneapolis Heart Institute. I cannot thank him enough for his support, friendship, and tremendous opportunity to work in a highly stimulating research environment. Thank you for your indispensable scientific guidance throughout my post-graduate fellowship in the US and for showing me how to be efficient in academic life.

I would also like to express my acknowledgement to **Prof. Dr. Tamás Forster**, Head of the Second Department of Internal Medicine and Cardiology Center, for all his support.

I wish to express my special gratitude to my family. My mother, **Mónika**, my father, **László**, my sisters, **Franciska** and **Krisztina**, my grandparents, **Piroska**, **János** and **Márton**, my nephew **András** and my brother in law, **Zsolt**. You made all this possible with your enormous love and encouragement throughout my years both in Hungary and the United States. Thank you for your continuous support and for showing me how to live a valuable life.

Last but not least, I would like thank my colleagues at the Division of Interventional Cardiology of Szeged, and the Minneapolis Heart Institute Foundation and its employees for supporting my scientific work throughout career in Hungary and the United States.

REFERENCES

1. Brilakis ES. Manual of Coronary Chronic Total Occlusion Interventions: A Step-by-Step Approach. 2nd edition, Academic Press, 2017.
2. Fefer P, Knudtson ML, Cheema AN et al. Current perspectives on coronary chronic total occlusions: the Canadian Multicenter Chronic Total Occlusions Registry. *J Am Coll Cardiol* 2012;59:991-7.
3. Jeroudi OM, Alomar ME, Michael TT et al. Prevalence and management of coronary chronic total occlusions in a tertiary Veterans Affairs hospital. *Catheter Cardiovasc Interv* 2014;84:637-43.
4. Christofferson RD, Lehmann KG, Martin GV, Every N, Caldwell JH, Kapadia SR. Effect of chronic total coronary occlusion on treatment strategy. *Am J Cardiol* 2005;95:1088-91.
5. Ramunddal T, Hoebbers LP, Henriques JP et al. Chronic total occlusions in Sweden-a report from the Swedish Coronary Angiography and Angioplasty Registry (SCAAR). *PLoS One* 2014;9:e103850.
6. Azzalini L, Jolicoeur EM, Pighi M et al. Epidemiology, Management Strategies, and Outcomes of Patients With Chronic Total Coronary Occlusion. *Am J Cardiol* 2016;118:1128-1135.
7. Patel MR, Dehmer GJ, Hirshfeld JW, Smith PK, Spertus JA. ACCF/SCAI/STS/AATS/AHA/ASNC 2009 Appropriateness Criteria for Coronary Revascularization: A Report of the American College of Cardiology Foundation Appropriateness Criteria Task Force, Society for Cardiovascular Angiography and Interventions, Society of Thoracic Surgeons, American Association for Thoracic Surgery, American Heart Association, and the American Society of Nuclear Cardiology: Endorsed by the American Society of Echocardiography, the Heart Failure Society of America, and the Society of Cardiovascular Computed Tomography. *Circulation* 2009;119:1330-52.
8. Patel MR, Calhoon JH, Dehmer GJ et al. ACC/AATS/AHA/ASE/ASNC/SCAI/SCCT/STS 2016 Appropriate Use Criteria for Coronary Revascularization in Patients With Acute Coronary Syndromes: A Report of the American College of Cardiology Appropriate Use Criteria Task Force, American Association for Thoracic Surgery, American Heart Association, American Society of Echocardiography, American Society of

Nuclear Cardiology, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, and the Society of Thoracic Surgeons. *J Am Coll Cardiol* 2017;69:570-591.

9. Patel MR, Calhoon JH, Dehmer GJ et al. ACC/AATS/AHA/ASE/ASNC/SCAI/SCCT/STS 2017 Appropriate Use Criteria for Coronary Revascularization in Patients With Stable Ischemic Heart Disease: A Report of the American College of Cardiology Appropriate Use Criteria Task Force, American Association for Thoracic Surgery, American Heart Association, American Society of Echocardiography, American Society of Nuclear Cardiology, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, and Society of Thoracic Surgeons. *J Am Coll Cardiol* 2017;69:2212-2241.

10. Neumann FJ, Sousa-Uva M, Ahlsson A et al. 2018 ESC/EACTS Guidelines on myocardial revascularization. *Eur Heart J* 2019;40:87-165.

11. Henriques JP, Hoehers LP, Ramunddal T et al. Percutaneous Intervention for Concurrent Chronic Total Occlusions in Patients With STEMI: The EXPLORE Trial. *J Am Coll Cardiol* 2016;68:1622-1632.

12. Lee SW, Lee PH, Ahn JM et al. Randomized Trial Evaluating Percutaneous Coronary Intervention for the Treatment of Chronic Total Occlusion. *Circulation* 2019;139:1674-1683.

13. Werner GS, Martin-Yuste V, Hildick-Smith D et al. A randomized multicentre trial to compare revascularization with optimal medical therapy for the treatment of chronic total coronary occlusions. *Eur Heart J* 2018.

14. Mashayekhi K, Nuhrenberg TG, Toma A et al. A Randomized Trial to Assess Regional Left Ventricular Function After Stent Implantation in Chronic Total Occlusion: The REVASC Trial. *JACC Cardiovasc Interv* 2018;11:1982-1991.

15. Obedinskiy AA, Kretoy EI, Boukhris M et al. The IMPACTOR-CTO Trial. *JACC Cardiovasc Interv* 2018;11:1309-1311.

16. Han H, Chao H, Guerra A et al. Evolution of the American College of Cardiology/American Heart Association Clinical Guidelines. *J Am Coll Cardiol* 2015;65:2726-34.

17. Sapontis J, Salisbury AC, Yeh RW et al. Early Procedural and Health Status Outcomes After Chronic Total Occlusion Angioplasty: A Report From the OPEN-CTO Registry

(Outcomes, Patient Health Status, and Efficiency in Chronic Total Occlusion Hybrid Procedures). *JACC Cardiovasc Interv* 2017;10:1523-1534.

18. Christakopoulos GE, Christopoulos G, Carlino M et al. Meta-analysis of clinical outcomes of patients who underwent percutaneous coronary interventions for chronic total occlusions. *Am J Cardiol* 2015;115:1367-75.

19. Tomasello SD, Boukhris M, Giubilato S et al. Management strategies in patients affected by chronic total occlusions: results from the Italian Registry of Chronic Total Occlusions. *Eur Heart J* 2015.

20. Yang JH, Kim BS, Jang WJ et al. Optimal Medical Therapy vs. Percutaneous Coronary Intervention for Patients With Coronary Chronic Total Occlusion - A Propensity-Matched Analysis. *Circ J* 2016;80:211-7.

21. Jang WJ, Yang JH, Choi SH et al. Long-term survival benefit of revascularization compared with medical therapy in patients with coronary chronic total occlusion and well-developed collateral circulation. *JACC Cardiovasc Interv* 2015;8:271-9.

22. Bruckel JT, Jaffer FA, O'Brien C, Stone L, Pomerantsev E, Yeh RW. Angina Severity, Depression, and Response to Percutaneous Revascularization in Patients With Chronic Total Occlusion of Coronary Arteries. *J Invasive Cardiol* 2016;28:44-51.

23. Rossello X, Pujadas S, Serra A et al. Assessment of Inducible Myocardial Ischemia, Quality of Life, and Functional Status After Successful Percutaneous Revascularization in Patients With Chronic Total Coronary Occlusion. *Am J Cardiol* 2016;117:720-6.

24. Mashayekhi K, Neuser H, Kraus A et al. Successful Percutaneous Coronary Intervention Improves Cardiopulmonary Exercise Capacity in Patients With Chronic Total Occlusions. *J Am Coll Cardiol* 2017;69:1095-1096.

25. Abdullah SM, Hastings JL, Amsavelu S et al. Percutaneous Coronary Intervention of Coronary Chronic Total Occlusions Improves Peak Oxygen Uptake During Cardiopulmonary Exercise Testing. *J Invasive Cardiol* 2017;29:83-91.

26. Di Marco A, Paglino G, Oloriz T et al. Impact of a chronic total occlusion in an infarct-related artery on the long-term outcome of ventricular tachycardia ablation. *J Cardiovasc Electrophysiol* 2015;26:532-9.

27. Iannaccone M, D'Ascenzo F, Piazza F et al. Optimal medical therapy vs. coronary revascularization for patients presenting with chronic total occlusion: A meta-analysis of

randomized controlled trials and propensity score adjusted studies. *Catheter Cardiovasc Interv* 2019;93:E320-E325.

28. Brilakis ES, Grantham JA, Rinfret S et al. A percutaneous treatment algorithm for crossing coronary chronic total occlusions. *JACC Cardiovasc Interv* 2012;5:367-79.

29. Christopoulos G, Karpaliotis D, Alaswad K et al. Application and outcomes of a hybrid approach to chronic total occlusion percutaneous coronary intervention in a contemporary multicenter US registry. *Int J Cardiol* 2015;198:222-228.

30. Wilson WM, Walsh SJ, Yan AT et al. Hybrid approach improves success of chronic total occlusion angioplasty. *Heart* 2016;102:1486-93.

31. Galassi AR, Sianos G, Werner GS et al. Retrograde Recanalization of Chronic Total Occlusions in Europe: Procedural, In-Hospital, and Long-Term Outcomes From the Multicenter ERCTO Registry. *J Am Coll Cardiol* 2015;65:2388-400.

32. Okamura A, Yamane M, Muto M et al. Complications during retrograde approach for chronic coronary total occlusion: Sub-analysis of Japanese multicenter registry. *Catheter Cardiovasc Interv* 2016;88:7-14.

33. Habara M, Tsuchikane E, Muramatsu T et al. Comparison of percutaneous coronary intervention for chronic total occlusion outcome according to operator experience from the Japanese retrograde summit registry. *Catheter Cardiovasc Interv* 2016;87:1027-35.

34. Maeremans J, Walsh S, Knaapen P et al. The Hybrid Algorithm for Treating Chronic Total Occlusions in Europe: The RECHARGE Registry. *J Am Coll Cardiol* 2016;68:1958-1970.

35. Suzuki Y, Tsuchikane E, Katoh O et al. Outcomes of Percutaneous Coronary Interventions for Chronic Total Occlusion Performed by Highly Experienced Japanese Specialists: The First Report From the Japanese CTO-PCI Expert Registry. *JACC Cardiovasc Interv* 2017;10:2144-2154.

36. Hannan EL, Zhong Y, Jacobs AK et al. Patients With Chronic Total Occlusions Undergoing Percutaneous Coronary Interventions: Characteristics, Success, and Outcomes. *Circ Cardiovasc Interv* 2016;9:e003586.

37. Brilakis ES, Banerjee S, Karpaliotis D et al. Procedural Outcomes of Chronic Total Occlusion Percutaneous Coronary Intervention: A Report From the NCDR (National Cardiovascular Data Registry). *JACC Cardiovascular interventions* 2015;8:245-53.

38. Morino Y, Abe M, Morimoto T et al. Predicting successful guidewire crossing through chronic total occlusion of native coronary lesions within 30 minutes: the J-CTO (Multicenter CTO Registry in Japan) score as a difficulty grading and time assessment tool. *JACC Cardiovascular interventions* 2011;4:213-21.
39. Christopoulos G, Wyman RM, Alaswad K et al. Clinical Utility of the Japan-Chronic Total Occlusion Score in Coronary Chronic Total Occlusion Interventions: Results from a Multicenter Registry. *Circ Cardiovasc Interv* 2015;8:e002171.
40. Tanaka H, Morino Y, Abe M et al. Impact of J-CTO score on procedural outcome and target lesion revascularisation after percutaneous coronary intervention for chronic total occlusion: a substudy of the J-CTO Registry (Multicentre CTO Registry in Japan). *EuroIntervention* 2016;11:981-8.
41. Abe M, Morimoto T, Morino Y et al. Association between J-CTO score and long-term target lesion revascularization rate after successful chronic total coronary occlusion angioplasty (from the J-CTO Registry). *Catheter Cardiovasc Interv* 2019;93:1025-1032.
42. Christopoulos G, Kandzari DE, Yeh RW et al. Development and Validation of a Novel Scoring System for Predicting Technical Success of Chronic Total Occlusion Percutaneous Coronary Interventions: The PROGRESS CTO (Prospective Global Registry for the Study of Chronic Total Occlusion Intervention) Score. *JACC Cardiovasc Interv* 2016;9:1-9.
43. Maeremans J, Spratt JC, Knaapen P et al. Towards a contemporary, comprehensive scoring system for determining technical outcomes of hybrid percutaneous chronic total occlusion treatment: The RECHARGE score. *Catheter Cardiovasc Interv* 2017.
44. Alessandrino G, Chevalier B, Lefevre T et al. A Clinical and Angiographic Scoring System to Predict the Probability of Successful First-Attempt Percutaneous Coronary Intervention in Patients With Total Chronic Coronary Occlusion. *JACC Cardiovasc Interv* 2015;8:1540-8.
45. Szigyarto Z, Rampat R, Werner GS et al. Derivation and Validation of a Chronic Total Coronary Occlusion Intervention Procedural Success Score From the 20,000-Patient EuroCTO Registry: The EuroCTO (CASTLE) Score. *JACC Cardiovasc Interv* 2019;12:335-342.
46. Galassi AR, Boukhris M, Azzarelli S, Castaing M, Marza F, Tomasello SD. Percutaneous Coronary Revascularization for Chronic Total Occlusions: A Novel Predictive Score of Technical Failure Using Advanced Technologies. *JACC Cardiovasc Interv* 2016.

47. Ellis SG, Burke MN, Murad MB et al. Predictors of Successful Hybrid-Approach Chronic Total Coronary Artery Occlusion Stenting: An Improved Model With Novel Correlates. *JACC Cardiovasc Interv* 2017;10:1089-1098.
48. Karatasakis A, Danek BA, Karpaliotis D et al. Comparison of various scores for predicting success of chronic total occlusion percutaneous coronary intervention. *Int J Cardiol* 2016;224:50-56.
49. Tajti P, Burke MN, Karpaliotis D et al. Update in the Percutaneous Management of Coronary Chronic Total Occlusions. *JACC Cardiovasc Interv* 2018;11:615-625.
50. Tajti P, Xenogiannis I, Najjar H, Brilakis ES. Management of Balloon Undilatable CTOs. *Cardiac Interventions Today* 2018;(May/June 2018).
51. Rinfret S, Joyal D, Spratt JC, Buller CE. Chronic total occlusion percutaneous coronary intervention case selection and techniques for the antegrade-only operator. *Catheter Cardiovasc Interv* 2014.
52. Michael TT, Papayannis AC, Banerjee S, Brilakis ES. Subintimal dissection/reentry strategies in coronary chronic total occlusion interventions. *Circ Cardiovasc Interv* 2012;5:729-38.
53. Colombo A, Mikhail GW, Michev I et al. Treating chronic total occlusions using subintimal tracking and reentry: the STAR technique. *Catheter Cardiovasc Interv* 2005;64:407-11; discussion 412.
54. Whitlow PL, Burke MN, Lombardi WL et al. Use of a novel crossing and re-entry system in coronary chronic total occlusions that have failed standard crossing techniques: results of the FAST-CTOs (Facilitated Antegrade Steering Technique in Chronic Total Occlusions) trial. *JACC Cardiovasc Interv* 2012;5:393-401.
55. Werner GS. The BridgePoint devices to facilitate recanalization of chronic total coronary occlusions through controlled subintimal reentry. *Expert Rev Med Devices* 2011;8:23-9.
56. Rathore S, Katoh O, Matsuo H et al. Retrograde percutaneous recanalization of chronic total occlusion of the coronary arteries: procedural outcomes and predictors of success in contemporary practice. *Circ Cardiovasc Interv* 2009;2:124-32.
57. Saito S. Different strategies of retrograde approach in coronary angioplasty for chronic total occlusion. *Catheter Cardiovasc Interv* 2008;71:8-19.

58. Brilakis ES, Grantham JA, Thompson CA et al. The retrograde approach to coronary artery chronic total occlusions: a practical approach. *Catheter Cardiovasc Interv* 2012;79:3-19.
59. Joyal D, Thompson CA, Grantham JA, Buller CE, Rinfret S. The retrograde technique for recanalization of chronic total occlusions: a step-by-step approach. *JACC Cardiovasc Interv* 2012;5:1-11.
60. Michael TT, Mogabgab O, Fuh E et al. Application of the "hybrid approach" to chronic total occlusion interventions: a detailed procedural analysis. *J Interv Cardiol* 2014;27:36-43.
61. Christopoulos G, Menon RV, Karpaliotis D et al. Application of the "hybrid approach" to chronic total occlusions in patients with previous coronary artery bypass graft surgery (from a Contemporary Multicenter US registry). *Am J Cardiol* 2014;113:1990-4.
62. Pershad A, Eddin M, Girotra S, Cotugno R, Daniels D, Lombardi W. Validation and incremental value of the hybrid algorithm for CTO PCI. *Catheter Cardiovasc Interv* 2014;84:654-9.
63. Sabbagh AE, Banerjee S, Brilakis ES. Illustration of the 'hybrid' approach to chronic total occlusion crossing. *Interventional Cardiology* 2012;4:639-643.
64. Christopoulos G, Menon RV, Karpaliotis D et al. The Efficacy and Safety of the "Hybrid" Approach to Coronary Chronic Total Occlusions: Insights From a Contemporary Multicenter US Registry and Comparison With Prior Studies. *J Invasive Cardiol* 2014;26:427-32.
65. Tajti P, Ungi I. Az anterográd technikák szerepe a krónikus teljes koszorúér-elzáródás perkután revaszkularizációjában. *Cardiologia Hungarica* 2018;48:330-339.
66. Karacsonyi J, Tajti P, Rangan BV et al. Randomized Comparison of a CrossBoss First Versus Standard Wire Escalation Strategy for Crossing Coronary Chronic Total Occlusions: The CrossBoss First Trial. *JACC Cardiovasc Interv* 2018;11:225-233.
67. Azzalini L, Dautov R, Brilakis ES et al. Procedural and longer-term outcomes of wire- versus device-based antegrade dissection and re-entry techniques for the percutaneous revascularization of coronary chronic total occlusions. *Int J Cardiol* 2017;231:78-83.
68. Thygesen K, Alpert JS, Jaffe AS et al. Third universal definition of myocardial infarction. *J Am Coll Cardiol* 2012;60:1581-98.

69. Morino Y, Abe M, Morimoto T et al. Predicting successful guidewire crossing through chronic total occlusion of native coronary lesions within 30 minutes: the J-CTO (Multicenter CTO Registry in Japan) score as a difficulty grading and time assessment tool. *JACC Cardiovasc Interv* 2011;4:213-21.
70. Danek BA, Karatasakis A, Karpaliotis D et al. Development and Validation of a Scoring System for Predicting Periprocedural Complications During Percutaneous Coronary Interventions of Chronic Total Occlusions: The Prospective Global Registry for the Study of Chronic Total Occlusion Intervention (PROGRESS CTO) Complications Score. *J Am Heart Assoc* 2016;5.
71. Vo MN, McCabe JM, Lombardi WL, Ducas J, Ravandi A, Brilakis ES. Adoption of the hybrid CTO approach by a single non-CTO operator: procedural and clinical outcomes. *J Invasive Cardiol* 2015;27:139-44.
72. Daniels DV, Banerjee S, Alaswad K et al. Safety and efficacy of the hybrid approach in coronary chronic total occlusion percutaneous coronary intervention: The Hybrid Video Registry. *Catheter Cardiovasc Interv* 2017.
73. Patel SM, Pokala NR, Menon RV et al. Prevalence and treatment of "balloon-uncrossable" coronary chronic total occlusions. *J Invasive Cardiol* 2015;27:78-84.
74. Karacsonyi J, Karpaliotis D, Alaswad K et al. Prevalence, indications and management of balloon uncrossable chronic total occlusions: Insights from a contemporary multicenter US registry. *Catheter Cardiovasc Interv* 2017;90:12-20.
75. Ojeda S, Pan M, Gutierrez A et al. Bifurcation lesions involved in the recanalization process of coronary chronic total occlusions: Incidence, treatment and clinical implications. *Int J Cardiol* 2017;230:432-438.
76. Tajti P, Karpaliotis D, Alaswad K et al. Prevalence, Presentation and Treatment of 'Balloon Undilatable' Chronic Total Occlusions: Insights from a Multicenter US Registry. *Catheter Cardiovasc Interv* 2018.
77. Michael TT, Karpaliotis D, Brilakis ES et al. Temporal trends of fluoroscopy time and contrast utilization in coronary chronic total occlusion revascularization: insights from a multicenter united states registry. *Catheter Cardiovasc Interv* 2015;85:393-9.

78. Tajti P, Alaswad K, Karpalotis D et al. In-hospital Outcomes of Attempting More Than One Chronic Total Coronary Occlusion Through Percutaneous Intervention During the Same Procedure. *Am J Cardiol* 2018;[Article in Press].
79. Karatasakis A, Danek BA, Karpalotis D et al. Impact of Proximal Cap Ambiguity on Outcomes of Chronic Total Occlusion Percutaneous Coronary Intervention: Insights From a Multicenter US Registry. *J Invasive Cardiol* 2016;28:391-396.
80. Karacsonyi J, Karpalotis D, Alaswad K et al. Impact of Calcium on Chronic Total Occlusion Percutaneous Coronary Interventions. *Am J Cardiol* 2017;120:40-46.
81. Teramoto T, Tsuchikane E, Matsuo H et al. Initial success rate of percutaneous coronary intervention for chronic total occlusion in a native coronary artery is decreased in patients who underwent previous coronary artery bypass graft surgery. *JACC Cardiovascular interventions* 2014;7:39-46.
82. Brilakis ES, O'Donnell CI, Penny W et al. Percutaneous Coronary Intervention in Native Coronary Arteries Versus Bypass Grafts in Patients With Prior Coronary Artery Bypass Graft Surgery: Insights From the Veterans Affairs Clinical Assessment, Reporting, and Tracking Program. *JACC Cardiovasc Interv* 2016;9:884-93.
83. Tajti P, Abu-Fanne R, Ungi I, Katona A, Sasi V, Nagy FT. Kettős lumenű mikrokatéterek alkalmazása krónikus teljes koronária okklúzió katéteres revaszkularizációjában - egy komplex beavatkozás tanulságai. *Cardiologia Hungarica* 2019.
84. Tajti P, Karpalotis D, Alaswad K et al. In-Hospital Outcomes of Chronic Total Occlusion Percutaneous Coronary Interventions in Patients With Prior Coronary Artery Bypass Graft Surgery. *Circ Cardiovasc Interv* 2019;12:e007338.
85. Ramunddal T, Hoebbers LP, Henriques JP et al. Prognostic Impact of Chronic Total Occlusions: A Report From SCAAR (Swedish Coronary Angiography and Angioplasty Registry). *JACC Cardiovascular interventions* 2016;9:1535-44.
86. Tajti P, Karpalotis D, Alaswad K et al. The Hybrid Approach to Chronic Total Occlusion Percutaneous Coronary Intervention: Update From the PROGRESS CTO Registry. *JACC Cardiovasc Interv* 2018.
87. Tsai TT, Stanislawski MA, Shunk KA et al. Contemporary Incidence, Management, and Long-Term Outcomes of Percutaneous Coronary Interventions for Chronic

Coronary Artery Total Occlusions: Insights From the VA CART Program. *JACC Cardiovasc Interv* 2017;10:866-875.

88. Sharma V, Jadhav ST, Harcombe AA et al. Impact of proctoring on success rates for percutaneous revascularisation of coronary chronic total occlusions. *Open Heart* 2015;2:e000228.

89. Brilakis ES, Vo MN. How to develop a successful chronic total occlusion percutaneous coronary intervention program. *Cardiovasc Revasc Med* 2016;17:3-4.

90. Shimura T, Yamamoto M, Tsuchikane E et al. Safety of Live Case Demonstrations in Patients Undergoing Percutaneous Coronary Intervention for Chronic Total Occlusion. *Am J Cardiol* 2016;118:967-73.

91. Kirtane AJ, Doshi D, Leon MB et al. Treatment of Higher-Risk Patients With an Indication for Revascularization: Evolution Within the Field of Contemporary Percutaneous Coronary Intervention. *Circulation* 2016;134:422-31.

92. Kalra A, Bhatt DL, Pinto DS et al. Accreditation and funding for a 24-month advanced interventional cardiology fellowship program: A call-to-action for optimal training of the next generation of interventionalists. *Catheter Cardiovasc Interv* 2016;88:1010-1015.

I.

The Hybrid Approach to Chronic Total Occlusion Percutaneous Coronary Intervention

Update From the PROGRESS CTO Registry

Peter Tajti, MD,^{a,b} Dimitri Karpaliotis, MD,^c Khaldoun Alaswad, MD,^d Farouc A. Jaffer, MD, PhD,^e Robert W. Yeh, MD,^f Mitul Patel, MD,^g Ehtisham Mahmud, MD,^g James W. Choi, MD,^h M. Nicholas Burke, MD,^a Anthony H. Doing, MD,ⁱ Phil Dattilo, MD,ⁱ Catalin Toma, MD,^j A.J. Conrad Smith, MD,^j Barry Uretsky, MD,^k Elizabeth Holper, MD,^l R. Michael Wyman, MD,^m David E. Kandzari, MD,ⁿ Santiago Garcia, MD,^o Oleg Krestyaninov, MD,^p Dmitrii Khelinskii, MD,^p Michalis Koutouzis, MD,^q Ioannis Tsiafoutis, MD,^q Jeffrey W. Moses, MD,^c Nicholas J. Lembo, MD,^c Manish Parikh, MD,^c Ajay J. Kirtane, MD,^c Ziad A. Ali, MD,^c Darshan Doshi, MD,^c Bavana V. Rangan, BDS, MPH,^r Imre Ungi, MD, PhD,^b Subhash Banerjee, MD,^r Emmanouil S. Brilakis, MD, PhD^{a,r}

ABSTRACT

OBJECTIVES The aim of this study was to determine the techniques and outcomes of hybrid chronic total occlusion (CTO) percutaneous coronary intervention (PCI) in a diverse group of patients and operators on 2 continents.

BACKGROUND CTO PCI has been evolving with constant improvement of equipment and techniques.

METHODS Contemporary outcomes of CTO PCI were examined by analyzing the clinical, angiographic, and procedural characteristics of 3,122 CTO interventions performed in 3,055 patients at 20 centers in the United States, Europe, and Russia.

RESULTS The mean age was 65 ± 10 years, and 85% of the patients were men, with high prevalence of diabetes (43%), prior myocardial infarction (46%), prior coronary artery bypass graft surgery (33%), and prior PCI (65%). The CTO target vessels were the right coronary artery (55%), left anterior descending coronary artery (24%), and left circumflex coronary artery (20%). The mean J-CTO (Multicenter Chronic Total Occlusion Registry of Japan) and PROGRESS CTO (Prospective Global Registry for the Study of Chronic Total Occlusion Intervention) scores were 2.4 ± 1.3 and 1.3 ± 1.0 , respectively. The overall technical and procedural success rate was 87% and 85%, respectively, and the rate of in-hospital major complications was 3.0%. The final successful crossing strategy was antegrade wire escalation in 52.0%, retrograde in 27.1%, and antegrade dissection re-entry in 20.9%; >1 crossing strategy was required in 40.9%. Median contrast volume, air kerma radiation dose, and procedure and fluoroscopy time were 270 ml (interquartile range: 200 to 360 ml), 2.9 Gy (interquartile range: 1.7 to 4.7 Gy), 123 min (interquartile range: 81 to 188 min) and 47 min (interquartile range: 29 to 77 min), respectively.

CONCLUSIONS CTO PCI is currently being performed with high success and acceptable complication rates among various experienced centers in the United States, Europe, and Russia. (Prospective Global Registry for the Study of Chronic Total Occlusion Intervention [PROGRESS CTO]; [NCT02061436](#)) (J Am Coll Cardiol Intv 2018;11:1325–35)

© 2018 the American College of Cardiology Foundation. Published by Elsevier. All rights reserved.

From the ^aMinneapolis Heart Institute, Abbott Northwestern Hospital, Minneapolis, Minnesota; ^bUniversity of Szeged, Division of Invasive Cardiology, Second Department of Internal Medicine and Cardiology Center, Szeged, Hungary; ^cColumbia University, New York, New York; ^dHenry Ford Hospital, Detroit, Michigan; ^eMassachusetts General Hospital, Boston, Massachusetts; ^fBeth Israel Deaconess Medical Center, Boston, Massachusetts; ^gVA San Diego Healthcare System and University of California, San Diego, La Jolla, California; ^hBaylor Heart and Vascular Hospital, Dallas, Texas; ⁱMedical Center of the Rockies, Loveland, Colorado; ^jUniversity of Pittsburgh Medical Center, Pittsburgh, Pennsylvania; ^kVA Central Arkansas Healthcare System, Little Rock, Arkansas; ^lThe Heart Hospital Baylor Plano, Plano, Texas; ^mTorrance Memorial Medical Center, Torrance, California; ⁿPiedmont Heart Institute, Atlanta, Georgia; ^oVA Minneapolis Healthcare System and University of Minnesota, Minneapolis.

ABBREVIATIONS AND ACRONYMS

CABG = coronary artery
bypass graft

CCS = Canadian Cardiovascular
Society

CTO = chronic total occlusion

IQR = interquartile range

MACE = major adverse
cardiac event(s)

MI = myocardial infarction

PCI = percutaneous coronary
intervention

TIMI = Thrombolysis In
Myocardial Infarction

The hybrid approach to chronic total occlusion (CTO) percutaneous coronary intervention (PCI) advocates dual coronary injection, careful and structured review of the angiogram, and flexibility (1). Use of all crossing strategies (antegrade wire escalation [2], antegrade dissection re-entry [3-6], and the retrograde approach [7-10]) is encouraged (1), with initial and subsequent choices influenced by the CTO anatomic characteristics and the outcomes of the originally selected approach (1,11-18). Application of the hybrid approach to CTO PCI has been associated with good outcomes in U.S. and European registries, although

CTO PCI outcomes in nonselected populations have been less optimal, with a success rate of approximately 60% (19). We examined a contemporary, multicenter CTO PCI registry to determine the techniques and outcomes of hybrid CTO PCI in a diverse group of patients and operators on 2 continents.

SEE PAGE 1336

METHODS

We analyzed the clinical, angiographic, and procedural characteristics of 3,122 CTO PCIs performed in 3,055

patients enrolled in the PROGRESS CTO (Prospective Global Registry for the Study of Chronic Total Occlusion Intervention; [NCT02061436](#)) registry between January 2012 and November 2017 at 18 U.S., 1 European, and 1 Russian center ([Online Appendix](#)). Some centers enrolled patients during only part of the study period because of participation in other studies. The study was approved by the Institutional Review Board of each center.

DEFINITIONS. Coronary CTOs were defined as coronary lesions with TIMI (Thrombolysis In Myocardial Infarction) flow grade 0 of at least 3 months' duration. Estimation of the duration of occlusion was clinical, based on the first onset of angina, history of myocardial infarction (MI) in the target vessel territory, or comparison with a prior angiogram. Calcification was assessed by angiography as mild (spots), moderate (involving $\leq 50\%$ of the reference lesion diameter), or severe (involving $>50\%$ of the reference lesion diameter). Moderate proximal vessel tortuosity was defined as the presence of at least 2 bends $>70^\circ$ or 1 bend $>90^\circ$ and severe tortuosity as 2 bends $>90^\circ$ or 1 bend $>120^\circ$ in the CTO vessel. Blunt or no stump was defined as lack of tapering or lack of a funnel shape at the proximal cap. Interventional collateral vessels were defined as collateral vessels considered amenable to crossing by a guidewire and a

Minnesota; ^PMeshalkin Siberian Federal Biomedical Research Center, Ministry of Health of Russian Federation, Novosibirsk, Russian Federation; ^QKorgialeneio-Benakeio Hellenic Red Cross General Hospital of Athens, Athens, Greece; and the ^{VA} North Texas Health Care System and University of Texas Southwestern Medical Center, Dallas, Texas. The PROGRESS CTO registry has received support from the Abbott Northwestern Hospital Foundation. Dr. Karpalotis has received speaking honoraria from Abbott Vascular, Boston Scientific, Medtronic, and Vascular Solutions. Dr. Alaswad has received consulting fees from Terumo and Boston Scientific and is a consultant (without financial remuneration) for Abbott Laboratories. Dr. Jaffer is a consultant for Abbott Vascular and Boston Scientific and has received research grants from Canon, Siemens, and the National Institutes of Health. Dr. Yeh has received a Career Development Award (1K23HL118138) from the National Heart, Lung, and Blood Institute. Dr. Patel is a member of the Speakers Bureau for AstraZeneca. Dr. Mahmud has received consulting fees from Medtronic and Corindus; speaking fees from Medtronic, Corindus, and Abbott Vascular; educational program fees from Abbott Vascular; and clinical events committee fees from St. Jude Medical. Dr. Burke has received consulting and speaking honoraria from Abbott Vascular and Boston Scientific. Dr. Wyman has received honoraria and consulting and speaking fees from Boston Scientific, Abbott Vascular, and Asahi Intecc USA, Inc. Dr. Kandzari has received research and grant support and consulting honoraria from Boston Scientific and Medtronic Cardiovascular and research and grant support from Abbott Vascular. Dr. Garcia has received consulting fees from Medtronic. Dr. Moses is a consultant to Boston Scientific and Abiomed. Dr. Lembo is a member of the Speakers Bureau for Medtronic; and advisory boards for Abbott Vascular and Medtronic. Dr. Parikh is a member of the Speakers Bureau for Abbott Vascular, Medtronic, Cardiovascular Systems Inc., Boston Scientific, and Trireme; and on the advisory boards for Medtronic, Abbott Vascular, and Philips. Dr. Kirtane has received institutional research grants to Columbia University from Boston Scientific, Medtronic, Abbott Vascular, Abiomed, St. Jude Medical, Vascular Dynamics, GlaxoSmithKline, and Eli Lilly. Dr. Ali has received consulting fees and honoraria from St. Jude Medical and AstraZeneca Pharmaceuticals; has ownership interest, partnership, or principal interest in Shockwave Medical and VitaBx; and has received research grants from Medtronic and St. Jude Medical. Dr. Rangan has received research grants from InfraRedx and Spectranetics. Dr. Banerjee has received research grants from Gilead and The Medicines Company; has received consulting and speaking honoraria from Covidien and Medtronic; has ownership in MDCARE Global (spouse); and has intellectual property in HygeiaTel. Dr. Brilakis has received consulting and speaking honoraria from Abbott Vascular, ACIST Medical Systems, Amgen, Asahi Intecc USA, Inc., Cardiovascular Systems Inc., Elsevier, GE Healthcare, Medicare, Medtronic, and Nitiloop; has received research support from Boston Scientific and Osprey; is a member of the board of directors of Cardiovascular Innovations Foundation; and is a member of the board of trustees of the Society of Cardiovascular Angiography and Interventions. All other authors have reported that they have no relationships relevant to the contents of this paper to disclose.

Manuscript received January 16, 2018; revised manuscript received February 12, 2018, accepted February 27, 2018.

microcatheter by the operator. A procedure was defined as “retrograde” if an attempt was made to cross the lesion through a collateral vessel or bypass graft supplying the target vessel distal to the lesion; if not, the procedure was classified as “antegrade only.” Antegrade dissection re-entry was defined as antegrade PCI during which a guidewire was intentionally introduced into the subintimal space proximal to the lesion, or re-entry into the distal true lumen was attempted following intentional or inadvertent subintimal guidewire crossing.

Technical success was defined as successful CTO revascularization with achievement of <30% residual diameter stenosis within the treated segment and restoration of TIMI antegrade flow grade 3. Procedural success was defined as the achievement of technical success without any in-hospital complications. In-hospital major adverse cardiac events (MACE) included any of the following adverse events before hospital discharge: death, MI, recurrent symptoms requiring urgent repeat target vessel revascularization with PCI or coronary artery bypass graft (CABG) surgery, tamponade requiring either pericardiocentesis or surgery, and stroke. MI was defined using the third universal definition of MI (type 4a MI) (20). Major bleeding was defined as bleeding causing reduction in hemoglobin >3 g/dl or bleeding requiring transfusion or surgical intervention. The J-CTO (Multicenter Chronic Total Occlusion Registry of Japan) score was calculated as described by Morino et al. (21), the PROGRESS CTO score as described by Christopoulos et al. (22), and the PROGRESS CTO complications score as described by Danek et al. (23).

STATISTICAL ANALYSIS. Categorical variables were expressed as percentages and were compared using Pearson’s chi-square test or the Fisher exact test. Continuous variables are presented as mean ± SD or as median (interquartile range [IQR]) unless otherwise specified and were compared using the Student’s *t*-test and 1-way analysis of variance for normally distributed variables; the Wilcoxon rank sum test and the Kruskal-Wallis test were applied for nonparametric continuous variables as appropriate. Multivariate logistic regression with stepwise backward elimination was performed to examine the independent association between annual CTO PCI volume and procedural outcomes (procedural success and in-hospital MACE). Variables with univariate associations in the present study (*p* < 0.05) were entered into the model, as well as variables that have been previously linked with procedural outcomes of CTO PCI, including age, smoking, peripheral arterial disease, chronic lung disease, history of MI, stroke,

TABLE 1 Clinical Characteristics of the Study Population, Classified According to Technical Success

	Overall (N = 3,055)	Technical Success (n = 2,657)	Technical Failure (n = 398)	p Value
Age (yrs)	64.80 ± 10.09	64.6 ± 10.15	66.01 ± 9.63	0.0141
Male	85.25	84.69	88.95	0.0378
BMI (kg/m ²)	30.60 ± 6.14	30.50 ± 6.15	31.20 ± 6.02	0.0666
Smoking (current)	26.01	25.37	30.27	0.0561
Diabetes	43.02	43.53	39.65	0.1758
Dyslipidemia	92.20	92.11	92.75	0.6781
Hypertension	90.26	89.61	94.49	0.0044
Family history of CAD	33.35	33.44	32.80	0.8423
CCS angina classification				0.4771
Class ≤1	11.44	11.64	10.15	
Class ≥2	88.56	88.36	89.85	
Myocardial viability performed	24.99	24.28	29.32	0.0783
Prior MI	46.00	44.82	53.75	0.0023
Heart failure	30.56	29.71	36.25	0.0159
Prior valve surgery or procedure	3.17	3.06	3.89	0.4210
Prior PCI	65.29	64.49	70.62	0.0180
Prior CABG surgery	32.49	31.28	40.68	0.0003
Baseline creatinine (mg/dl)	1.01 (0.89–1.22)	1.01 (0.89–1.21)	1.07 (0.90–1.27)	0.1301
Currently on dialysis	2.67	2.50	3.80	0.1633
Prior CVD	11.70	11.51	12.90	0.4567
Prior PAD	15.02	14.53	18.29	0.0709
Chronic lung disease	14.20	13.80	16.81	0.1386
Left ventricular EF (%)	54 (42–60)	55 (44–60)	50 (40–60)	0.0357

Values are mean ± SD, %, or median (interquartile range).
BMI = body mass index; CABG = coronary artery bypass graft; CAD = coronary artery disease; CVD = cerebrovascular disease; EF = ejection fraction; MI = myocardial infarction; PCI = percutaneous coronary intervention; PAD = peripheral arterial disease.

PCI or CABG surgery, left ventricular ejection fraction, CTO target vessel, multiple CTO vessels treated during the same procedure, and CTO PCI only. All statistical analyses were performed using JMP version 13.0 (SAS Institute, Cary, North Carolina). A 2-sided *p* value of 0.05 was considered to indicate statistical significance.

RESULTS

CLINICAL AND ANGIOGRAPHIC CHARACTERISTICS.

The baseline clinical features of the study population are summarized in Table 1. Compared with patients who had CTO PCI fail, patients who had successful CTO PCI were younger and less likely to be men and to have hypertension. They were also less likely to have had an MI, congestive heart failure, prior CABG surgery, and prior PCI and had higher left ventricular ejection fraction. Most patients (88.56%) were symptomatic, having at least Canadian Cardiovascular Society (CCS) angina classification class II

TABLE 2 Angiographic Characteristics of Study Lesions, Classified According to Technical Success

	Overall (N = 3,122)	Technical Success (n = 2,711)	Technical Failure (n = 411)	p Value
Target vessel				0.0640
RCA	55.22	54.93	57.14	
LAD	23.81	24.57	18.80	
LCx	19.91	19.47	22.81	
Other	1.06	1.03	1.25	
CTO length (mm)	33.99 ± 24.16	33.43 ± 24.14	37.80 ± 23.99	0.0030
Vessel diameter (mm)	2.85 ± 0.51	2.86 ± 0.51	2.81 ± 0.47	0.1383
Proximal cap ambiguity	35.06	31.98	53.97	<0.0001
Side branch at proximal cap	49.91	47.56	64.14	<0.0001
Blunt stump/no stump	53.69	50.61	72.55	<0.0001
Interventional collateral vessels	56.72	58.80	44.19	<0.0001
Moderate/severe calcification	54.23	52.30	67.02	<0.0001
Moderate/severe tortuosity	34.96	33.43	45.21	<0.0001
In-stent restenosis	16.61	16.13	19.68	0.0878
Previously failed CTO PCI	20.20	19.21	26.70	0.0005
J-CTO score	2.43 ± 1.30	2.34 ± 1.29	3.07 ± 1.13	<0.0001
PROGRESS CTO score	1.32 ± 1.03	1.25 ± 1.01	1.77 ± 1.01	<0.0001
PROGRESS CTO complication score	3.07 ± 1.93	3.00 ± 1.91	3.54 ± 1.97	<0.0001

Values are % or mean ± SD.
CTO = chronic total occlusion; J-CTO = Multicenter Chronic Total Occlusion Registry of Japan; LAD = left anterior descending coronary artery; LCx = left circumflex coronary artery; LM = left main segment; PCI = percutaneous coronary intervention; PROGRESS CTO = Prospective Global Registry for the Study of Chronic Total Occlusion Intervention; RCA = right coronary artery.

(CCS class II, 24.97%; CCS class III, 53.93%; CCS class IV, 9.65%), and most had stable (64.33%) or unstable (18.20%) angina.

The angiographic characteristics of the study lesions are presented in **Table 2**. The CTO target lesions were located in the right coronary artery (55.22%), left anterior descending coronary artery (23.81%), and left circumflex coronary artery (19.91%). Failed CTO PCI was associated with longer lesion length (33.4 ± 24.1 mm vs. 37.9 ± 24.0 mm; $p = 0.0030$), proximal cap ambiguity (53.97% vs. 31.98%; $p < 0.0001$), moderate to severe calcification (72.55% vs. 52.30%; $p < 0.0001$), and tortuosity (45.21% vs. 33.43%; $p < 0.0001$) or previously failed recanalization attempt (26.70% vs. 19.21%; $p = 0.0005$). Failed CTO PCI cases also had higher J-CTO scores (2.34 ± 1.29 vs. 3.07 ± 1.13; $p < 0.0001$), PROGRESS CTO scores (1.25 ± 1.01 vs. 1.77 ± 1.13; $p < 0.0001$), and PROGRESS CTO complications scores (3.00 ± 1.91 vs. 3.54 ± 1.97; $p < 0.0001$).

PROCEDURAL OUTCOMES OF THE HYBRID APPROACH. Overall technical and procedural success was 87% and 85%, respectively, and the in-hospital major complications rate was 3.0%. The baseline technical and procedural characteristics are presented in **Tables 3 and 4**. Antegrade wire escalation was the most commonly used initial approach

TABLE 3 Techniques Used for Chronic Total Occlusion Percutaneous Coronary Intervention in the Study Patients

	Overall (N = 3,122)	Technical Success (n = 2,711)	Technical Failure (n = 411)	p Value
Crossing strategies used				
AWE	81.77	81.15	85.89	0.0204
ADR	31.68	29.69	44.77	<0.0001
Retrograde	38.57	35.96	55.72	<0.0001
First crossing strategy				0.0124
AWE	75.36	75.80	72.51	
ADR	8.39	8.67	6.57	
Retrograde	16.24	15.53	20.92	
Final crossing strategy				<0.0001
AWE	45.89	51.95	5.24	
ADR	18.95	20.96	5.49	
Retrograde	23.97	27.09	2.99	
None	11.19	0.00	86.28	
Balloon-uncrossable lesions	10.62	10.21	29.37	<0.0001
Balloon-undilatable lesions	11.11	10.74	22.22	0.0349
Access site				
Right femoral	78.96	78.46	82.24	0.0798
Left femoral	54.29	53.34	60.58	0.0060
Right radial	32.48	33.12	28.22	0.0481
Left radial	18.67	19.48	13.38	0.0031
Bifemoral approach	51.35	50.42	57.42	0.0082
Biradial approach	14.09	14.90	8.76	0.0009

Values are %.
ADR = antegrade dissection reentry; AWE = antegrade wire escalation.

(in 75%), especially for lower complexity CTOs (J-CTO score 2.28 ± 1.29 , PROGRESS CTO score 1.35 ± 1.05), whereas antegrade dissection re-entry (8%; J-CTO score 2.86 ± 1.16 , PROGRESS CTO score 1.50 ± 1.07) and the retrograde approach (16%; J-CTO score 3.12 ± 1.07 , PROGRESS CTO score 1.33 ± 0.96) were used for more complex lesions ($p < 0.0001$). The initial approach was successful in 55% of patients, whereas 41% of patients underwent further attempts that were technically successful in 79% (**Figure 1**).

The final successful crossing strategy was antegrade wire escalation (46%), antegrade dissection re-entry (19%), and the retrograde approach (24%). The success of antegrade wire escalation decreased with lesion complexity, as classified with the J-CTO score (easy [J-CTO score 0], 88%; intermediate [J-CTO score 1], 72%; difficult [J-CTO score 2], 51%; and very difficult [J-CTO score ≥ 3], 32% to 17%; $p < 0.0001$) and the PROGRESS CTO score (55%, 43%, 42%, 39%, and 43%, respectively for scores of 0, 1, 2, 3, and 4; $p < 0.0001$). The retrograde approach was more commonly required for complex lesions, as classified by the J-CTO score (3%, 9%, 20%, and 35% to 44%, respectively for J-CTO scores of 0, 1, 2,

TABLE 4 Procedural Characteristics, Classified According to Percutaneous Coronary Intervention Outcomes and Lesion Complexity (J-CTO Score)

	Overall	Technical Success	Technical Failure	p Value	Easy (0)	Intermediate (1)	Difficult (2)	Very Difficult (≥3)	p Value
Dual injection	70.41	69.57	75.82	0.0260	47.59	64.19	68.70	77.89	<0.0001
Radial access*	37.06	37.70	32.85	0.0577	49.59	39.09	35.83	38.17	0.0030
Femoral access	81.90	81.37	85.40	0.0480	63.27	78.17	82.65	84.09	<0.0001
Procedure time (min)	123 (81-188)	121 (80-184)	140 (85-224)	0.0003	77 (46-117)	92 (58-136)	115 (73-172)	152 (102-217)	<0.0001
Contrast volume (ml)	270 (200-360)	260 (200-350)	300 (220-400)	0.0001	220 (160-300)	250 (190-340)	260 (200-350)	280 (202-385)	<0.0001
Fluoroscopy time (min)	47.0 (28.6-77.0)	45.0 (27.3-73.7)	66.0 (39.0-93.6)	0.0001	27.0 (18.1-39.2)	33.1 (19.3-53.7)	41.9 (27.1-65.7)	63.1 (38.6-93.0)	<0.0001
Patient AK dose (Gy)	2.9 (1.7-4.7)	2.8 (1.6-4.5)	3.9 (2.4-6.0)	0.0001	1.8 (0.9-3.3)	2.1 (1.2-3.5)	2.5 (1.4-4.4)	3.5 (2.1-5.2)	<0.0001
Number of stents†	2.4 ± 1.1	2.4 ± 1.1	2.7 ± 1.6	0.5559	1.8 ± 0.9	2.0 ± 1.0	2.3 ± 1.1	2.8 ± 1.2	<0.0001
Stent length (mm)†	71.8 ± 36.4	71.7 ± 36.3	78.6 ± 47.6	0.6599	48.0 ± 25.2	57.6 ± 31.1	66.5 ± 33.5	85.5 ± 36.3	<0.0001
Non-CTO lesion PCI	28.20	28.39	26.80	0.5611	32.75	32.00	29.19	26.37	0.0786
In-hospital MACE	3.04	2.37	7.54	0.0001	1.36	1.41	3.01	3.11	0.0119
Technical success	86.84	—	—	—	96.90	94.84	89.14	81.26	<0.0001

Values are %, median (interquartile range), or mean ± SD. *Radial access indicates any radial access site use in CTO PCI; including biradial and combined radial-femoral approaches. †In successful cases stents were implanted in 97.10% versus in failed procedures in 4.89% (related to perforation, investment procedure, donor vessel dissection, or stenting but Thrombolysis In Myocardial Infarction flow grade ≥3) (p < 0.0001).

AK = air kerma; J-CTO = Multicenter Chronic Total Occlusion Registry of Japan; MACE = major adverse cardiac event(s); PCI = percutaneous coronary intervention.

FIGURE 1 Application of the Hybrid Approach in the Study Patients

Sequence of chronic total occlusion (CTO) crossing techniques used in the study patients and associated technical success. PCI = percutaneous coronary intervention.

FIGURE 2 Technical Success and Crossing Strategy Use According to J-CTO Score and the PROGRESS CTO Score

Impact of chronic total occlusion (CTO) lesion complexity, as assessed by the J-CTO score (A) and PROGRESS CTO score (B) on technical success and use of various crossing strategies. ADR = antegrade dissection re-entry; AWE = antegrade wire escalation; J-CTO = Multicenter Chronic Total Occlusion Registry of Japan; PROGRESS CTO = Prospective Global Registry for the Study of Chronic Total Occlusion Intervention.

and ≥ 3 ; $p < 0.0001$) but less frequently in lesions with higher PROGRESS CTO score (23%, 27%, 23%, 15%, and 6%; $p < 0.0001$) (Figures 2A and 2B).

Dual injection was used in 70% of all cases and was more frequent in failed interventions (76% vs. 70%; $p = 0.026$) and in complex lesions with high J-CTO scores (48% vs. 78%; $p < 0.0001$). Radial access was used in 37% overall, with a biradial approach in 14% and in combination with a femoral approach in 20% of cases. Use of radial access was lower with increasing lesion complexity (easy, 50%; intermediate, 39%;

difficult, 36%; very difficult, 38%; $p = 0.003$), whereas the frequency of femoral (63%, 78%, 83%, and 84%; $p < 0.0001$) and bifemoral (28%, 43%, 51%, and 57%; $p < 0.0001$) approaches increased with increasing lesion complexity. Median contrast volume, air kerma radiation dose, and procedural and fluoroscopy time were 270 ml (IQR: 200 to 360 ml), 2.9 Gy (IQR: 1.7 to 4.7 Gy), and 123 min (IQR: 81 to 188 min) and 47.0 min (IQR: 28.6 to 77.0 min), respectively, and were higher for more complex lesions (Table 4). Less complex lesions required fewer stents ($p < 0.0001$), but the frequency of non-CTO PCI was higher in those groups and decreased with increasing lesion complexity ($p = 0.08$).

The procedural success and annual CTO PCI volume at the participating sites are shown in Online Figure 1. Higher median annually performed CTO PCI per center was associated with higher procedural success in both univariate and multivariate analysis (Figure 3) but not in-hospital MACE.

The incidence of in-hospital MACE was 3.04% (death, 0.85%; acute MI, 1.08%; stroke, 0.26%; emergency CABG surgery, 0.16%; urgent repeat PCI, 0.36%; and pericardial tamponade, 0.85%) and increased with increasing lesion complexity (Table 4). The prevalence of in-hospital MACE was higher in failed procedures (7.54% vs. 2.37%; $p < 0.0001$) and with more complex crossing techniques: antegrade wire escalation, antegrade dissection re-entry, or retrograde crossing (1.09% vs. 2.96% vs. 5.61%; $p < 0.0001$). Use of the retrograde approach was associated with higher incidence of complications, as shown in Figure 4. Median length of hospital stay was significantly higher in patients with versus without in-hospital MACE (6 days [IQR: 2 to 9 days] vs. 1 day [IQR: 1 to 2 days]); $p < 0.0001$).

DISCUSSION

To the best of our knowledge, this is the largest study reported to date on CTO PCI using the hybrid approach, demonstrating a high technical success rate (88%) with an acceptable major complication rate (3.0%). These outcomes were achieved despite high lesion complexity and relatively low success of the initially selected CTO crossing strategy (55%) (Figure 5).

Previous smaller studies have provided similarly encouraging results. In an earlier report from the PROGRESS CTO registry from 11 U.S. centers, technical success was 91% and in-hospital MACE was 1.7% (11). Wilson et al. (17) reported a 79% initial success rate among 1,156 patients from 7 centers enrolled in the UK Hybrid CTO Registry, with a 90%

FIGURE 3 Baseline Clinical and Angiographic Characteristics and Procedural Success

final technical success rate after repeat procedures. The RECHARGE (Registry of Crossboss and Hybrid Procedures in France, the Netherlands, Belgium, and United Kingdom) registry of 1,253 CTO interventions performed in 1,177 patients between 2014 and 2015 at 22 European centers reported an 86% procedural success rate and a 2.6% major in-hospital complication rate (18). Vo et al. (24) reported a single-operator pilot study showing a rapid increase in procedural success despite the low initial CTO PCI experience. At 2 high-volume, experienced centers Pershad et al. (14) showed significant increase in technical (from 79.4% to 95.4%; $p < 0.001$) and procedural (from 77.9% to 88.3%) success rates after implementation of the hybrid algorithm, compared with the pre-implementation period. Furthermore, the Hybrid Video Registry analyzed 194 video-recorded live case demonstrations reporting a high success rate (92.8%), even in highly complex CTOs, with acceptable procedure time and contrast volume (25). As shown in prior studies, higher annual CTO PCI volume was independently associated with

higher success rates, reflecting the importance of center and operator experience in optimizing outcomes, especially among complex lesion and patient subgroups (26).

In the present study, we found that technical and procedural success remained high, with reasonably low complication rates, despite expansion of the registry in recent years. Antegrade wire escalation was more commonly applied as the initial crossing approach (74%) for less complex lesions (J-CTO score 2.24 ± 1.24 , PROGRESS CTO score 1.32 ± 0.87) and was the most common final crossing strategy (in approximately one-half of the cases). Antegrade dissection re-entry and retrograde techniques were more likely to be used as initial strategy in cases with complex anatomy (J-CTO scores 2.78 ± 1.21 and 3.32 ± 0.98 , respectively; PROGRESS CTO scores 1.38 ± 0.93 and 2.00 ± 0.89 , respectively) and were the final successful strategies in 22% and 28% of all cases, respectively.

Failure to cross with a guidewire was the most common reason for CTO PCI failure (in 86%). In

13%, the procedure failed despite successful guide-wire crossing (Table 3) because of balloon-undilatable lesions (3.9%), inability to deliver stents (2.3%), final TIMI flow grade <3 (1.3%), residual stenosis >30% (1.0%), and procedure-related complications (0.8%; 1 patient with donor vessel thrombosis, 1 patient with aortocoronary dissection, and 1 procedure related death due to pericardial tamponade and subsequent cardiogenic shock). The presence of balloon-uncrossable (29.4% vs. 10.2%; $p < 0.0001$) and balloon-undilatable lesions (22.2% vs. 10.7; $p = 0.0109$) was higher in the failed CTO PCI group, highlighting the need for CTO PCI operators to have experience in treating these and other complex lesions, such as severe calcification and bifurcations (27,28).

The overall complication rate was 3%, and complications occurred less frequently in technically successful procedures (2.2% vs. 7.9%; $p < 0.0001$). The risk for complications was higher in more complex lesions (easy [J-CTO score 0] 1.36% vs. very difficult [J-CTO score ≥ 3] 3.11%; $p = 0.01$) and with use of advanced crossing techniques (which were more commonly used for more complex lesions). This highlights the importance of weighing the risks and benefits of the procedure, both during discussions with patients and family (to determine whether CTO PCI should be done) and during the procedure itself: implementing more complex CTO crossing strategies (such as retrograde crossing via epicardial collateral

vessels) may predispose to increased risk for complications, which may be justified in some patients because of significant potential benefit, but not in some others.

Despite the encouraging findings of our study and other contemporary registries, the success rates of CTO interventions in unselected patient cohorts remain low. Hannan et al. (29) (New York State PCI Registry, $n = 4,030$ patients) reported a 61.3% success rate with a 1.07% complication rate (vs. 1.06% for non-CTO PCI cases, $p = 0.95$). Ramunddal et al. (30) showed a similarly low procedural success rate (54.2%) among 6,442 patients undergoing CTO intervention in SCAAR (Swedish Coronary Angiography and Angioplasty Registry). Habara et al. (31) compared the acute outcomes of 3,229 CTO interventions among 56 high- and low-volume centers in Japan, showing a higher overall success rate at high-volume centers (90.6% vs. 85.6%; $p < 0.001$), without a significant difference in in-hospital MACE rate (0.45% vs. 0.62%; $p = 0.35$), attributed mostly to a higher antegrade success rate. Sharma et al. (32) showed that procedural outcomes of CTO interventions among operators who had received proctorship in using the hybrid approach were better (77.5% vs. 62.1%; $p < 0.0001$), especially in more complex cases (for J-CTO scores ≥ 2 , the corresponding rates were 70.7% and 49.5%, respectively; $p = 0.0003$) than those who did not receive mentorship. Hence, CTO PCI should be performed by

FIGURE 5 Clinical Case, Illustrating the Importance of Changing Crossing Strategy to Successfully Recanalize a Challenging Chronic Total Occlusion

Percutaneous coronary intervention (PCI) of a distal right coronary artery (RCA) chronic total occlusion (CTO) in a patient with Canadian Cardiovascular Society class III angina and 4 prior failed CTO attempts and 2 prior coronary artery bypass graft surgeries. (A) Left internal mammary artery (LIMA) injection demonstrating septal and epicardial collateral channels from the left anterior descending coronary artery (LAD) to the RCA. (B) Triple coronary injection showing mild disease in the saphenous vein graft (SVG) to the RCA, distal RCA CTO immediately distal to the SVG anastomotic site, and mild disease in previously stented proximal and mid LAD along with competitive flow from the LIMA graft. (C) Multiple attempts for septal collateral crossing (using surfing and contrast guided techniques) succeeded in advancing a guidewire to the distal RCA, but a microcatheter could not be advanced over the guidewire. (D) Successful contrast-guided epicardial collateral crossing using a Suoh 03 guidewire over a Caravel microcatheter (Asahi Intecc, Nagoya, Japan). (E) Several attempts to advance the retrograde guidewire distal to the SVG anastomosis failed, despite using multiple guidewires. (F) Multiple attempts to recanalize the native RCA failed, as the antegrade guidewire could not be advanced distal to the SVG anastomosis. (G) The reverse controlled antegrade and retrograde tracking and dissection (CART) was successfully used to cross the CTO and advance the retrograde guidewire into the SVG-RCA, followed by guidewire externalization. (H) Final angiographic result after stent implantation. The patient had significant symptom alleviation.

experienced operators at dedicated centers to achieve optimal results.

STUDY LIMITATIONS. First, we did not have mid- and long-term follow-up of the study patients. Second, there was no core laboratory assessment of the study angiograms or clinical event adjudication. Third, the procedures were performed at dedicated, high-volume CTO centers by experienced operators, limiting the extrapolation to less experienced operators at low-volume centers.

CONCLUSIONS

CTO PCI can currently be achieved with high success and acceptable complication rates among various operators and patient populations in the United States and Europe, highlighting the need for developing more CTO PCI centers of excellence in order to

achieve the best possible clinical outcomes in this challenging patient and lesion group.

ACKNOWLEDGMENTS Study data were collected and managed using Research Electronic Data Capture electronic data capture tools hosted at the Minneapolis Heart Institute Foundation. Research Electronic Data Capture is a secure, Web-based application designed to support data capture for research studies, providing: 1) an intuitive interface for validated data entry; 2) audit trails for tracking data manipulation and export procedures; 3) automated export procedures for seamless data downloads to common statistical packages; and 4) procedures for importing data from external sources.

ADDRESS FOR CORRESPONDENCE: Dr. Emmanouil S. Brilakis, Minneapolis Heart Institute, 920 East 28th Street, #300, Minneapolis, Minnesota 55407. E-mail: esbrilakis@gmail.com.

PERSPECTIVES

WHAT IS KNOWN? Coronary CTO can be challenging to recanalize, with success rates of approximately 60% in unselected, all-comer populations.

WHAT IS NEW? Application of the hybrid approach resulted in a high technical success rate (87%) and an acceptable rate of major in-hospital complications (3%) across a large number of sites and operators in the United States, Europe, and Russia. Changing crossing strategy was

required in 41% of cases, with the final success strategy being antegrade wire escalation in 52%, retrograde in 27%, and antegrade dissection re-entry in 21%.

WHAT IS NEXT? Bridging the gap between what is currently achieved and what can be achieved in CTO intervention should be a major focus of upcoming research and education efforts.

REFERENCES

- Brilakis ES, Grantham JA, Rinfret S, et al. A percutaneous treatment algorithm for crossing coronary chronic total occlusions. *J Am Coll Cardiol Interv* 2012;5:367–79.
- Rinfret S, Joyal D, Spratt JC, Buller CE. Chronic total occlusion percutaneous coronary intervention case selection and techniques for the antegrade-only operator. *Catheter Cardiovasc Interv* 2015;85:408–15.
- Michael TT, Papayannis AC, Banerjee S, Brilakis ES. Subintimal dissection/reentry strategies in coronary chronic total occlusion interventions. *Circ Cardiovasc Interv* 2012;5:729–38.
- Colombo A, Mikhail GW, Michev I, et al. Treating chronic total occlusions using subintimal tracking and reentry: the STAR technique. *Catheter Cardiovasc Interv* 2005;64:407–11.
- Whitlow PL, Burke MN, Lombardi WL, et al. Use of a novel crossing and re-entry system in coronary chronic total occlusions that have failed standard crossing techniques: results of the FAST-CTOs (Facilitated Antegrade Steering Technique in Chronic Total Occlusions) trial. *J Am Coll Cardiol Interv* 2012;5:393–401.
- Werner GS. The BridgePoint devices to facilitate recanalization of chronic total coronary occlusions through controlled subintimal reentry. *Expert Rev Med Devices* 2011;8:23–9.
- Rathore S, Katoh O, Matsuo H, et al. Retrograde percutaneous recanalization of chronic total occlusion of the coronary arteries: procedural outcomes and predictors of success in contemporary practice. *Circ Cardiovasc Interv* 2009;2:124–32.
- Saito S. Different strategies of retrograde approach in coronary angioplasty for chronic total occlusion. *Catheter Cardiovasc Interv* 2008;71:8–19.
- Brilakis ES, Grantham JA, Thompson CA, et al. The retrograde approach to coronary artery chronic total occlusions: a practical approach. *Catheter Cardiovasc Interv* 2012;79:3–19.
- Joyal D, Thompson CA, Grantham JA, Buller CE, Rinfret S. The retrograde technique for recanalization of chronic total occlusions: a step-by-step approach. *J Am Coll Cardiol Interv* 2012;5:1–11.
- Christopoulos G, Karpaliotis D, Alaswad K, et al. Application and outcomes of a hybrid approach to chronic total occlusion percutaneous coronary intervention in a contemporary multicenter US registry. *Int J Cardiol* 2015;198:222–8.
- Michael TT, Mogabgab O, Fuh E, et al. Application of the “hybrid approach” to chronic total occlusion interventions: a detailed procedural analysis. *J Interv Cardiol* 2014;27:36–43.
- Christopoulos G, Menon RV, Karpaliotis D, et al. Application of the “hybrid approach” to chronic total occlusions in patients with previous coronary artery bypass graft surgery (from a Contemporary Multicenter US registry). *Am J Cardiol* 2014;113:1990–4.
- Pershad A, Eddin M, Girotra S, Cotugno R, Daniels D, Lombardi W. Validation and incremental value of the hybrid algorithm for CTO PCI. *Catheter Cardiovasc Interv* 2014;84:654–9.
- Sabbagh AE, Banerjee S, Brilakis ES. Illustration of the “hybrid” approach to chronic total occlusion crossing. *Interventional Cardiology* 2012;4:639–43.
- Christopoulos G, Menon RV, Karpaliotis D, et al. The efficacy and safety of the “hybrid” approach to coronary chronic total occlusions: insights from a contemporary multicenter US registry and comparison with prior studies. *J Invasive Cardiol* 2014;26:427–32.
- Wilson WM, Walsh SJ, Yan AT, et al. Hybrid approach improves success of chronic total occlusion angioplasty. *Heart* 2016;102:1486–93.
- Maeremans J, Walsh S, Knaapen P, et al. The hybrid algorithm for treating chronic total occlusions in Europe: the RECHARGE registry. *J Am Coll Cardiol* 2016;68:1958–70.
- Brilakis ES, Banerjee S, Karpaliotis D, et al. Procedural outcomes of chronic total occlusion percutaneous coronary intervention: a report from the NCDR (National Cardiovascular Data Registry). *J Am Coll Cardiol Interv* 2015;8:245–53.
- Thygesen K, Alpert JS, Jaffe AS, et al. Third universal definition of myocardial infarction. *J Am Coll Cardiol* 2012;60:1581–98.
- Morino Y, Abe M, Morimoto T, et al. Predicting successful guidewire crossing through chronic total occlusion of native coronary lesions within 30 minutes: the J-CTO (Multicenter CTO Registry in Japan) score as a difficulty grading and time assessment tool. *J Am Coll Cardiol Interv* 2011;4:213–21.
- Christopoulos G, Kandzari DE, Yeh RW, et al. Development and validation of a novel scoring system for predicting technical success of chronic total occlusion percutaneous coronary interventions: the PROGRESS CTO (Prospective Global Registry for the Study of Chronic Total Occlusion Intervention) Score. *J Am Coll Cardiol Interv* 2016;9:1–9.
- Danek BA, Karatasakis A, Karpaliotis D, et al. Development and validation of a scoring system for predicting periprocedural complications during percutaneous coronary interventions of chronic total occlusions: the Prospective Global Registry for the Study of Chronic Total Occlusion Intervention (PROGRESS CTO) complications score. *J Am Heart Assoc* 2016;5:e004272.
- Vo MN, McCabe JM, Lombardi WL, Ducas J, Ravandi A, Brilakis ES. Adoption of the hybrid CTO approach by a single non-CTO operator: procedural and clinical outcomes. *J Invasive Cardiol* 2015;27:139–44.
- Daniels DV, Banerjee S, Alaswad K, et al. Safety and efficacy of the hybrid approach in coronary chronic total occlusion percutaneous coronary intervention: the Hybrid Video Registry. *Catheter Cardiovasc Interv* 2018;91:175–9.
- Michael TT, Karpaliotis D, Brilakis ES, et al. Temporal trends of fluoroscopy time and contrast utilization in coronary chronic total occlusion revascularization: insights from a multicenter united states registry. *Catheter Cardiovasc Interv* 2015;85:393–9.
- Patel SM, Pokala NR, Menon RV, et al. Prevalence and treatment of “balloon-uncrossable” coronary chronic total occlusions. *J Invasive Cardiol* 2015;27:78–84.
- Karacsonyi J, Karpaliotis D, Alaswad K, et al. Prevalence, indications and management of balloon uncrossable chronic total occlusions:

insights from a contemporary multicenter US registry. *Catheter Cardiovasc Interv* 2017;90:12–20.

29. Hannan EL, Zhong Y, Jacobs AK, et al. Patients With chronic total occlusions undergoing percutaneous coronary interventions: characteristics, success, and outcomes. *Circ Cardiovasc Interv* 2016;9:e003586.

30. Ramunddal T, Hoebors LP, Henriques JP, et al. Prognostic impact of chronic total occlusions: a report from SCAAR (Swedish Coronary

Angiography and Angioplasty Registry). *J Am Coll Cardiol Interv* 2016;9:1535–44.

31. Habara M, Tsuchikane E, Muramatsu T, et al. Comparison of percutaneous coronary intervention for chronic total occlusion outcome according to operator experience from the Japanese retrograde summit registry. *Catheter Cardiovasc Interv* 2016;87:1027–35.

32. Sharma V, Jadhav ST, Harcombe AA, et al. Impact of proctoring on success rates for percutaneous revascularisation of coronary

chronic total occlusions. *Open Heart* 2015;2:e000228.

KEY WORDS chronic total occlusion, outcomes, percutaneous coronary intervention, techniques

APPENDIX For a list of centers participating in the present study and a supplemental figure, please see the online version of this paper.

II.

Az anterográd technikák szerepe a krónikus teljes koszorúér-elzáródás perkután revaszkularizációjában

Tajti Péter^{1,2}, Ungi Imre¹

¹Szegedi Tudományegyetem, Invazív Kardiológiai Részleg, II. Sz. Belgyógyászati Klinika és Kardiológiai Központ, Szeged

²Minneapolis Heart Institute, Abbott Northwestern Hospital, Minneapolis, Minnesota, Egyesült Államok

Levelezési cím:

Dr. Ungi Imre PhD, Szegedi Tudományegyetem, Invazív Kardiológiai Részleg, II. sz. Belgyógyászati Klinika és Kardiológiai Központ, 6725 Szeged, Semmelweis u. 6., e-mail: ungi.imre@gmail.com

A krónikus teljes koszorúér-elzáródás perkután koronáriaintervenciója rohamos fejlődésen ment keresztül az elmúlt évtized alatt, amelynek következtében a dedikált CTO-centrumok beavatkozási sikeraránya jelentősen javult alacsony szövődésménnyel. Sikeres CTO PCI a megfelelően megválasztott esetekben jelentős javuláshoz segíti a betegeket, ugyanakkor a CTO-revaszkularizációnak továbbra is a legnagyobb kihívása az okklúzió megnyitása. A széles körben alkalmazott technikák magukba foglalják az anterográd dróteszkalációt, az anterográd disszekció és re-entry, valamint a retrográd technikákat. Anterográd dróteszkaláció továbbra is a leggyakrabban alkalmazott, és legtöbb esetben sikeres technika, az anterográd disszekció re-entry és a retrográd technikákat főleg komplex léziók esetén alkalmazzuk. Cikkünk célja, hogy átfogó képet adjunk a jelenkori anterográd technikákról és eszközökről, amelyek elősegítik a vezetődrót átjutását az okklúzión.

Kulcsszavak: krónikus teljes koszorúér-elzáródás, perkután koronária-intervenció, stabil koronáriabetegség, komplex koronária-intervenció

The Role of Antegrade Techniques in Current Practice of Percutaneous Coronary Interventions for Chronic Total Occlusions

Chronic total occlusion (CTO) percutaneous coronary intervention (PCI) has been rapidly evolving with high success and acceptable complication rates currently being achieved at dedicated high-volume centers. Successful CTO PCI in appropriately selected patients can provide significant clinical benefits, however crossing the occlusion with a guidewire can be challenging. The following techniques can be used for CTO crossing: antegrade wire escalation, antegrade dissection/re-entry and the retrograde approach. Antegrade wire escalation remains the most frequently used technique and is the most common final successful technique, especially in less complex CTOs. Antegrade dissection re-entry and the retrograde approach are more technically demanding, but are often necessary to achieve crossing, in complex CTOs. In this manuscript, we provide a comprehensive overview of antegrade wire crossing of coronary CTOs.

Keywords: chronic total occlusion, percutaneous coronary intervention, stable coronary artery disease, complex coronary intervention

A krónikus teljes koszorúér-elzáródás (CTO) definíció szerint a koronária lumenének teljes (TIMI [Thrombolysis In Myocardial Infarction] flow 0) elzáródása, amely fennállásának időtartama minimum három hónap koronarográfiával, vagy annak hiányában a klinikai kép alapján alátámasztva. A CTO incidenciája 18-52% közötti azon betegeken, akik diagnosztikus szívkatéterezéssel esnek át koronáriabetegség tisztázása céljából

(1–4), ugyanakkor ezen betegek nagyobb része konzervatív gyógyszeres terápiában vagy műtéti revaszkularizációban részesül, csupán kis hányadukban végeznek perkután koronáriaintervenciót (PCI) (5). Ennek az alacsony aránynak lehetséges magyarázatai között szerepel a korlátozott intervenció tapasztalat CTO PCI terén, a magasabb költségek, a relatíve hosszú beavatkozások miatti katéteres laborzsúfoltság, az átlagos in-

tervenciókét meghaladó szövődmenyráta, és a továbbra is ellentmondásos kép a CTO-revaszkularizációnak klinikai hasznáról.

Az elmúlt években a CTO PCI technikája jelentős fejlődésen ment keresztül bizonyítva, hogy specializált centrumokban az esetek 85-90%-ában sikeres revaszkularizáció érhető el alacsony periprocedurális kompli-

káció rátával (3%) (6–9). A legnagyobb technikai kihívást továbbra is a vezetődrót átjuttatása jelenti a distalis valódi lumen felé (10, 11), ami nem specializált centrumokban csak 50-60%-os sikerarányt eredményez (4, 12, 13).

Az anginas panaszokkal, illetve szívelégtelenség tüneteivel élő betegek nagy száma miatt a krónikus teljes

1. ÁBRA. Kockázat-haszon arány klinikai felbecslése a krónikus teljes koszorúér-elzáródások perkután koronária-intervenciója során (14)

koszorúér-elzáródás perkután revaszkularizációja iránti igény emelkedik, amely több dedikált CTO-centrum kialakítását igényli hazánkban is. Az elmúlt évek technikai fejlődése elsősorban a jelenkori anterográd CTO rekanalizációs technikák elterjedését eredményezte, amelyek rutinszerű elsajátítása fontos sarokköve a sikeres CTO-program elindításának.

Esetválasztás

Egy adott centrum és operatőr felkészültségével arányos komplexitású esetek megválasztása kritikus az intervenció sikere és biztonsága szempontjából, így a kockázat-haszon arány mérlegelése az első fontos lépés az intervenciót megelőzően (1. ábra) (14). Az elzáródás PCI-vel megoldhatóságának felmérését 4 fő kérdés köré kell csoportosítanunk:

- milyen klinikai javulás várható a sikeres CTO PCI elvégzését követően,
- mekkora a valószínűsége a beavatkozás sikerének az angiográfiás kép és a rendelkezésre álló eszköztár alapján,
- az operatőrök képzettsége megfelelő-e (2. ábra), valamint

- a beavatkozási szövődmények esélyének mérlegelése.

CTO komplexitás

Számos pontozási rendszert validáltak és közöltek az elmúlt évek alatt a CTO-intervenciók várható kimenetelének megítélésére (J-CTO Score [15], CL Score [16], PROGRESS CTO Score [17], ORA Score [18], RECHARGE Score [19], és az Ellis Score [20]). Ezek a pontrendszerek különböző klinikai és angiográfiás faktorokat ötvöznek az eset komplexitásának és a beavatkozás nehézségének becslésére (1. táblázat) (15–21). Jelenleg a legáltalánosabban elterjedt és validált pontrendszer a J-CTO Score (22–24), amely segítségével megjósolható a vezetődrót distalis lumenbe juttatásának esélye 30 percen belül 5 alapvető paraméter figyelembevételével (lézióhossz >20 mm, 45°-os lézió tortuozitás, meszesedés angiográfiás jelei, a proximális sapka alakja [V-alakú vagy tompa], és korábbi sikertelen CTO intervenció).

A CTO pontrendszerek alkalmazása fontos lehet több szempontból is:

- felméri a speciális rekanalizációs technikák szükségességét (anterográd disszekció és re-entry [ADR] és/vagy retrográd technikák) (3. ábra);

1. TÁBLÁZAT. A jelenleg elérhető CTO pontozási rendszerek

Tényezők	J-CTO score (15)	CL score (16)	PROGRESS CTO score (17)	ORA score (18)	RECHARGE score (19)	Ellis core (20)	PROGRESS CTO Complications score (21)
Esetszám (n)	494	1,657	781	1,073	1,253	456	1589
Végpont	Drót eszkálcio <30 min	Angiográfiás siker	Angiográfiás siker	Angiográfiás siker	Angiográfiás siker	Angiográfiás siker	Periprocedurális komplikációk
Életkor (év)	–	–	–	+ (≥75)	+ (>65)	–	+ (>65)
Korábbi CABG	–	+	–	–	+	–	–
Korábbi sikertelen kísérlet	+	–	–	–	–	–	–
Proximális sapka	+ (Tompá)	+ (Tompá)	+ (Homályos)	+ (Ostialis)	+	+ (Homályos, ostialis)	–
Tortuozitás	+ (>45°, okkludált szegmentben)	–	+ (okkludált szegmenttől proximálisant)	–	+	+	–
Meszesedés	+	+	–	–	+	+	–
Lézió hossz	+ (≥20 mm)	+ (≥20 mm)	–	–	+	+	+ (>23 mm)
Okkludált ér	–	+ (nem-LAD)	+ (LCX)	–	–	+ (Kedvezőtlen distalis érszakasz)	–
Kollaterálisok minősége	–	–	+ (Intervenció)	+ (Rentrop <2)	–	+	–
Egyéb	–	Korábbi miokardiális infarktus	–	–	BMI >30, nem proximális lokáció	CTO intervenció tapasztalat	Retrográd technika alkalmazása

[†]A CTO-tól proximálisan lévő érszakasznak minimum 2 görbülete a 70°-ot, vagy 1 görbülete a 90°-ot meghaladja.

BMI, body mass index; CABG, coronary artery bypass grafting; LAD, left anterior descending artery; LCX, circumflex artery.

- beavatkozás sikeres kimenetelének predikciója a rendelkezésre álló eszközök és intervenciók tapasztalati adottságok alapján (25), valamint
- az intervenció időigényének becslésére (több magas komplexitású eset egy napra történő tervezése jelentősen megterhelheti a centrum emberi erőforrásait) (26).

A CTO pontrendszerek rutinszerű alkalmazásának kritikus szerepe van az anterográd technikák sikerének becslésében továbbá a beavatkozások megtervezésében, valamint a beteg előzetes tájékoztatásában is.

3. ÁBRA. Sikeres revaszkularizációs technikák megoszlása a lézió komplexitásának függvényében a J-CTO score alapján (6)

A technika megválasztása

Az optimális elsődleges és az azt követő intervenciók stratégiák kiválasztása a koronária, illetve az okklúzió anatómiájának alapján kulcsfontosságú a beavatkozás sikeréhez (27). Igen lényeges az angiográfiás kép alapos tanulmányozása (szükség szerint akár 15-20 percre) az intervenciót megelőzően. A krónikus teljes okklúzió patomorfológiai (4. ábra) felépítése, azaz a proximális sapka alakja és mésztartalma, a mikrocatornák jelenléte, valamint a distális sapka morfológiája és a distális érszakasz minősége alapvető az intervenció megtervezéséhez. Az okklúzió anatómiájának precíz és megfelelő vizualizációjához az esetek legnagyobb hányadában szimultán ipsilaterális és kontralaterális kontraszt-injekció („dual angiography”) szükséges, amelynek során nagyítás és a képmező mozgatása nélkül megfelelő hosszúságú felvételeket kell készítenünk. Alternatív lehetőség a mikrokateéterbe adott kontrasztanyag („tip injection”) a distális érszakasz vizualizálására (5. ábra), főleg ha elsősorban ipsilaterális epikardiális kollaterálisok dominálnak –

4. ÁBRA. A krónikus teljes koszorúér-elzáródások morfológiájának sematikus felépítése

5. ÁBRA. Ipsilateralis szimultán kontraszt injekció vezérelte anterográd CTO-revaszkularizáció

ugyanakkor kontralaterális, kollaterális hálózat feltérképezése esetén a konvencionális kontralaterális injekció eredményesebb képet adhat.

Jelenleg két algoritmus érhető el a klinikai gyakorlatban, amely segítséget nyújt a CTO rekanalizációs stratégia megválasztásában:

- a hibrid algoritmus (27) és az
- Asia Pacific algoritmus (28).

A hibrid algoritmus (6. ábra) bilaterális injekció segítségével 4 anatómiai tulajdonságra fókuszálva (proximális sapka alakja, distális érszakasz minősége, intervenciós kollaterálisok jelenléte, és a lézió hosszúsága [$>$ vagy <20 mm]) tesz javaslatot a kezdő stratégia megválasztására (anterográd/retrográd, vezetődrót-eszkaláció/disszekció re-entry) (29). A hibrid algoritmus lényege a szükség esetén gyors és rugalmas intervenció – anterográd és retrográd – stratégiák alkalmazása

(a kezdő technika csupán az esetek 50-60%-ában eredményes), amely emeli revaszkularizációk sikerességét (6).

Az Asia Pacific algoritmus elsősorban az intra-plakk rekanalizációs szemléletmódot képviseli, azaz kevésbé támaszkodik a disszekció-re-entry technikákra. Ez a stratégia – szemben a hibrid algoritmussal – intravaszkuláris ultrahang (IVUS) vagy CT-angiográfia (CTA) gyakori alkalmazását igényli (28).

Bilaterális kontraszt injekció alkalmazása kritikus az okklúzió és a kollaterális hálózat anatómiájának megítélésében, amelyet szinte minden CTO PCI esetében rutinszerűen ajánlott elvégezni. Algoritmikus stratégiaválasztás és a rendelkezésre álló technikák alternáló használata növeli a CTO-intervenció sikerességét valamint segíti az ad hoc döntések elkerülését a beavatkozás alatt.

6. ÁBRA. A hibrid algoritmus (14)

Anterográd dróteszkaláció (antegrade wire escalation [AWE]) technikák

Az anterográd dróteszkaláció a jelenkori CTO intervenciók gyakorlatban az alapvető, és legelterjedtebben alkalmazott technika, amelynek hatékonysága főleg kevésbé komplex okklúziók esetén jelentős. Ismerete és biztonságos alkalmazása a különböző bonyolultabb technikák (ADR és retrográd) elsajátításának is alapját képezi (3. ábra). Az AWE-technika alkalmazása optimális a rövid (<20 mm), egyenes, és mikrocsonnákat tartalmazó elzáródások esetén (4. ábra).

Eszközök és a behatolási kapu megválasztása

A megfelelő katéterszupport biztosítása rendkívül fontos a CTO-intervenciókban, hiszen ez jelentősen képes javítani a vezetődrót penetrációs képességét. A megfelelő szupport biztosítására számos technika áll rendelkezésre:

- nagy átmérőjű vezetőkatéterek,
- mikrokatéterek,
- extenziós katéterek vagy
- a különböző horgonyzó technikák.

Radiális vs. femoralis behatolás

A koszorúér-intervenciók többségében ma általánosan elfogadott a radiális behatolás elsődleges alkalmazása. CTO PCI esetén nem ennyire egyértelmű a kérdés megválaszolása, hiszen általánosan elfogadott tény, hogy az elérhető és a beteg által tolerált legnagyobb belső átmérőjű, vezetőkatéter a preferálandó, amely számos esetben femoralis behatolást tesz szükségessé. A 7-8 F (French) transfemoralis vezetőkatéterek használata a leggyakrabban javasolt (lehetőleg hosszú, 45 cm-es sheath használatával), amely optimális szupportot biztosít, ugyanakkor e katéterekkel a vaszkuláris szövődmények esélye is magasabb (30). Ultrahang-vezérelte punkció (31) és/vagy fluoroszkópiás ellenőrzés mellett (32) csökkenthető a perifériás komplikációk kialakulásának esélye, ha az operátor a femoralis artéria használata mellett dönt. Ezzel szemben számos CTO-operátor kombinált radialis-femoralis vagy bilaterális radiális beavatkozást alkalmaz, amely a 7 F „slender sheath” (Terumo, Somerset, New Jersey) alkalmazásával elegendő belső diamétert és támasztékot biztosíthat. A „hibrid vaszkuláris” szemlélet elterjedése, azaz a kombinált radialis és femoralis behatolás jó megoldás az esetek többségében, mivel a nagy átmérőjű femoralis vezetőkatéter megfelelő anterográd szupportot ad, míg a radialis feljuttatott katéter (kisebb átmérőjű sheath [6-7 F]) elegendő teret biztosít a retrográd eszközök felvezetéséhez (33).

Mikrokatéterek

Mikrokatéterek a CTO-intervencióban minden esetben alkalmazandóak, mint a legegyszerűbb, de egyben leghatékonyabb megoldás a vezetődrót penetrációs ké-

pességének javítására. A mikrokatéterek a vezetődrót végének formálását/újra formálását, és kicserélését is lehetővé teszik a pozíció elvesztése nélkül. Jelen gyakorlatban az over-the-wire (OTW) ballonkatéterek alkalmazása mikrokatéterek helyett nem ajánlott, mivel a sugárfogó marker a ballon közepén helyezkedik el, ezáltal megnehezíti a precíz pozicionálást, valamint az OTW-ballonok fizikai tulajdonságai sem alkalmasak az okklúzió történő áthatolásra.

Extenziós katéterek

Ezen eszközök használata (Guidezilla [Boston Scientific, Natick, MA, USA]; GuideLiner [Teleflex, Wayne, PA, USA]) – amelyet az angol nomenklatura „mother-and-child” technikaként tart számon (34) – növeli azt a koaxiális erőt, amelyet a drót kifejthet a kalcifikált szövetre. Az extenziós katéterek mindemellett elősegítik a különböző eszközök lejuttatását is distalis léziók vagy kanyarulatok proximális koronáriák esetén, így csökkentik a koszorúér-szájadékok disszekciójának a mély vezetőkatéter inkubációból származó veszélyét. Extrém távolságok, vagy tortuozitás esetén többszöri extenziós katéter használata is lehetséges – amely teleszkópos technikaként is ismert (angol nomenklaturában „mother-daughter-granddaughter” technika) (35) – azonban ez mindenképp 8 F vezetőkatéter használatához kötött.

Horgonyzó technikák

Számos horgonyzó technika elérhető, amelyek célja ugyancsak a vezetőkatéter szupportjának növelése. A horgonyzás általános koncepciója, hogy egy második levezetett vezetődróton keresztül ballont fújunk fel, amely a vezetőkatétert az ér szájadékában „lehorgonyozza”. E technikának több formája létezik: az okklúziótól proximálisan elhelyezkedő oldalágban (oldalág horgony) (36); az okklúziótól proximálisan a főérben (koaxiális horgony) (37) – a ballon rögzíti a mikrokatétert is az érfalhoz); az okklúziótól distálisan (distalis horgony) (38) a vezető drót átjutását követően, amely elsősorban a stent lejuttatását segítheti); vagy a subintimalis térben (subintimalis horgony) (39). A horgonyzás alkalmazása esetén figyelembe kell vennünk az oldalágakban okozott érsérülés vagy az okklúzió miatti hosszan tartó iszkémia veszélyét.

Megfelelő támaszték biztosítása esszenciális a vezetődrót penetrációs képességének javítása érdekében, amelyhez a lehető legnagyobb átmérőjű katéterek használata javasolt. Mikrokatéterek minden CTO-intervenció esetében rutinszerűen alkalmazandóak. Egyéb technikák, mint például az extenziós katéterek vagy horgonyzási technikák további segítséget nyújthatnak a koaxiális erő növelése érdekében.

A vezetődrót előkészítése és drótozási technikák

Az anterográd drót eszkalációs technika 4 fő fázisból áll, amely az

- okklúzió proximális sapkájához való eljutást,
- a proximális sapka penetrációját,
- a drót okklúzióban történő navigációját, és
- a distalis valódi lumenbe jutást jelenti.

A CTO megközelítése

A speciális CTO-vezetődrót és a mikrokatóter atraumatikus lejuttatása a proximális sapkához reguláris intervenciók drótok segítségével történik. Ezzel elkerülhetjük a proximális érszakasz sérülését, továbbá a dedikált CTO-drót görbületének útközbéli torzulását. A proximális sapkát elérve a mikrokatóter e drótra rávezethető, és ott az eredeti drót a speciális CTO-vezetődrótra kicserélhető. Eközben mikrokatóter tip-injekció egyes esetekben további információt nyújthat a proximális sapka anatómiájáról.

A proximális sapka penetrációja

A vezetődrót megválasztását elsősorban a proximális sapka anatómiája határozza meg. Abban az esetben, ha az ér lefutása jól követhető (rövid, egyenes érszakasz), merev kúpos végű drótok a választandók (Gaia 2nd, vagy igen kemény sapka esetén Confianza Pro 12 [Asahi Intecc. Nagoya, Japán]), ugyanakkor, ha az ér architektúrája nem egyértelmű, kevésbé merev, polimer bevonatú (Fielder XT család [Asahi Intecc. Nagoya, Japán], vagy Pilot 200 [Abbott Laboratories, Chicago, US]) vagy polimer bevonat nélküli drótok (Ultimate Bros 3, Miracle [Asahi Intecc. Nagoya, Japán]) képesek az ér lefutását megfelelően követni. Az ideális vezetődrót görbület a proximális sapka penetrációjához az 1 mm hosszú 30–45° szöget bezáró, amely maximális penetrációs kapacitás mellett a mikrocatornak útvonalát is megfelelően követni tudja csökkentve a subintimalis térbe jutás esélyét. A proximális sapka extrém kalcifikációja esetén a drót rotáció nélküli szűrő jellegű előre mozgatása (penetrációs technika) is eredményes lehet.

A vezetődrót navigációja az okklúzióban

A proximális sapka sikeres áttörését követően az okkludált szegmensten belül történő manipuláció eltérő technikát igényel, amely gyakori drótcserével is járhat (eszkaláció/deeszkaláció). A „sliding” technika, amely mérsékelt rotációs mozgás és penetrációval kombinált drótozást jelent, polimer bevonatú drótokkal eredményes lehet, ugyanakkor rezisztens léziók esetén erőteljesebb rotációs mozgás „drilling technika” válhat szükségessé. A „drilling-technikához” a Gaia-vezetődrót család az ideális választás (Gaia 1st, 2nd, 3rd) mivel ezen drótok mikrostruktúrája megfelelő manőverezési képességet biztosít még a hosszú és tortuózus érszakaszok esetén is minimalizálva a perforáció esélyét.

Behatolás a valódi lumenbe

Miután a vezetődrót elérte a distalis sapka síkját, mérsekelt (Gaia család, Ultimate Bros 3, Miracle 6) vagy kifejezetten merev (Confianza Pro 12) végű drót

használata eredményezheti a valódi lumenbe jutást, ugyanakkor a distalis drót pozícionak a megerősítése nagyon lényeges, hiszen esetleges perforáció dróttal többnyire nem veszélyes, ugyanakkor mikrokatóterrel vagy ballonnal perikardiális tamponádot okozhat. A drót valódi lumen pozíciójának megerősítése többnyire kontralaterális angiográfiával történik. Ha a bejuttatott mikrokatóterből vért tudunk aspirálni, vagy nyomásgörbe detektálható a mikrokatóteren keresztül, az szintén kisegítő információt nyújthat az intraluminalis pozícióról.

„Trapping” technika

Az esetek jelentős többségében több, különböző penetrációs készségű vezetődróra is szükség lehet. Ilyen esetekben szükséges lehet a mikrokatóter eltávolítása az első drót helyben hagyásával. A „trapping” technika során egy segédballont fűjünk fel a vezetőkatéteren belül (2,0 mm átmérőjű 6 F-hez, 2,5 mm átmérőjű 7 F-hez). Ez hozzápréseli a vezetődrótot a katéter falához stabilizálva annak pozícióját a mikrokatóter eltávolításához. Erre a célra fejlesztett speciális ballonok előnye (Trapper, Boston Scientific, Natick, MA, USA), hogy igen alacsony profiljuk kevés helyet foglal el, továbbá megegyező hosszúságúak a vezetőkatéterével, így fluoroszkópia nélkül is biztonságosan behelyezhetők a koronária sérülésének elkerülésével.

Vezetődrótok tudatos megválasztása, előkészítése, és a drótozási manőverek megfelelő alkalmazása alapvető fontosságú a sikeres CTO PCI érdekében. A drótpozíció megtartása kritikus a dróteszkaláció és deeszkaláció során, ezért a trapping technika elsajátítása és megfelelő alkalmazása fontos részét képezi az anterográd CTO-operatőrök gyakorlatának.

Speciális anterográd drótozási technikák

Ideális esetben az anterográd eszkálcációs technika sikeres (7. ábra, A panel), mindazonáltal az esetek jelentékeny hányadában a drót a subintimalis térbe jut, ahonnan igen nehéz és sokszor csak nagy gyakorlattal lehetséges a visszavezetésük a lumen felé. E célból a drót végét újra kell formáznunk, ekkor 70–90°-os 2–3 mm hosszú görbületet képezünk. Ki kell emelnünk ugyanakkor, hogy subintimalis pozíció esetén a legjobb eredményt a dedikált anterográd disszekció re-entry technika alkalmazása adja erre specializált eszközökkel (CrossBoss mikrokatóter, Stingray ballon [mindkettő Boston Scientific, Natick, MA, USA]) (40). A csupán dróttal történő ún. random re-entry technika azzal a veszéllyel jár, hogy nagy subintimalis haematomát képezünk, ezután már egyre nehezebb a valódi lumenbe jutás. A hosszú szakaszon subintimalis stentelés pedig magas restenosis és reokklúziós rátát okozhat (41, 42). A dedikált re-entry eszközökön kívül is lehetséges azonban a distalis valódi lumenbe visszajutás a következő eljárások alkalmazásával.

7. ÁBRA. Speciális anterográd drótozási technikák: eszkálációs technika (A); paralleldrótozás (B); „libikóka technika” (C); kettős lumenű mikrokatóéter technika (D); mikrocsatorna („Carlino”) technika (E)

Paralleldrótozási technika (43)

Abban az esetben, ha az első vezetődrt helyzete subintimalisnak sejtethető, annak helyben hagyása (mikrokatóéter nélkül) mellett egy második vezetődrt szimultán felvezetése – parallel helyzetben az elsőhöz képest (7. ábra, B panel) – elősegítheti második vezetődrt intraluminalis vezetését azáltal, hogy a helyben hagyott első drót kiegyenesíti az adott érszakaszt, valamint meggátolja hogy a második drót ugyanazt a fals utat kövesse. Kettőnél több vezetődrt is használható, ugyanakkor ez jelentősen megnehezítheti az angiográfiás tájékozódást.

„Libikóka technika” (43)

A libikóka technika egy módosított paralleldrótozási eljárás, amely során az első vezetődrt és mikrokatóéter a subintimalis térben marad, mellé pedig egy második mikrokatóéter-vezetődrt együttest vezetünk az okklúzióba. Ezen két mikrokatóéter segítségével felváltva manipulálhatunk a két dróttal (7. ábra C panel). Ilyen esetben

második drótként érdemes merev, de könnyen irányítható eszközöket választani, mint a Gaia-vezetődrtok, vagy a Miracle 6, illetve 12.

Kettős lumenű mikrokatóéterek (44)

A kettős lumenű mikrokatóéterek alkalmazása kezdeti subintimalis drótpozíció felvétele során ideális opcióként szolgálnak a második vezetődrt helyes irányba juttatásához (7. ábra D-panel). Ezen mikrokatóéterek monorail- (distalis) és over-the-wire (proximális) portot tartalmaznak. Ha a primer vezetődrt subintimalis pozícióba jut, a mikrokatóétert trapping technikával kettős lumenű katóéterre cserélhetjük, majd az oldalsó proximális porton keresztül (over-the-wire) a szekunder vezetődrt intraluminalis drót manipulációra használható. A kettős lumenű mikrokatóéterek használata ugyancsak jelentős szerepet játszik a bifurkációs okklúziók oldalágának rekanalizációjában.

„Szezám tárulj technika” (45)

Ezen módszer csak olyan CTO-anatómia esetében alkalmazható, ahol az okklúzió közvetlenül distalisan kezdődik az oldalágtól. A „szezám tárulj technika” lényege, hogy az oldalágba jutott vezetődrtön ballonnal tágítást végzünk, amely módosítja az okklúzió architektúráját elősegítve a második vezetődrt könnyebb bevezetését a distalis valódi lumen felé. Mindazonáltal ez a technika csak egy speciális koronária-anatómia esetén alkalmazható, valamint a subintimalis drótozás esélye is magasabb („side-BASE technika”) (46), amely így jelentős mértékben korlátozza ennek a technikának széles körű alkalmazhatóságát.

Mikrocsatorna-technika („Carlino technika”) (47)

A mikrocsatorna-technika lényege a mikrokatóéter okklúzióba vezetése, majd finoman 0,5-1,0 ml kontrasztanyag injektálása az okklúzió testébe, amely kontrollált hidraulikus disszekciót okoz a mikrocsatorna mentén elősegítve az okklúzió „felpuhítását”, valamint a koronária lefutásának feltérképezését (7. ábra, E panel). A Carlino-technika eredendően anterográd drót eszkáláció elősegítésére lett fejlesztve, azonban ez gyakran a subintimalis tér megnyitásával is járhat, amely a dedikált disszekciós re-entry technikák alkalmazását követeli meg (Stingray-ballon), így ez a technika csak megfelelő intervenciós gyakorlattal alkalmazható biztonságosan.

Komplex anterográd technikák alkalmazásának el-sajátítása jelentős tapasztalatot és megfelelő eszköztárat igényel a biztonságos használathoz, azonban e módszerek hasznos stratégiaként szolgálnak, ha a vezetődrt subintimalis pozícióba jutott. Subintimalis drótozás, és dedikált re-entry eszközök nélküli anterográd disszekció és re-entry technika (klasszikus STAR-technika) klinikai alkalmazását nem javasoljuk a mai intervenciós gyakorlatban a magas restenosis és reokklúziós ráta miatt.

A rekanalizáció befejező lépései

Amint a vezetődrót a distalis valódi lumenbe jutott, és annak intraluminalis pozíciójáról meggyőződünk, a mikrokatóteron át a drót cseréje következik hagyományos vezetődrótra, célszerűen extra szupport drótra (Sion Blue ES, BHW, Grand Slam). Ennek a lépésnek jelentősége, hogy a CTO-vezetődrótok distalis vége jó penetrációs képességű, így könnyen okozhatnak distalis perforációt. E ponttól az intervenció az angioplasztika általánosan ismert lépéseinek megfelelően zajlik (ballon predilatáció és stentimplantáció), ugyanakkor az esetek kis százalékában a ballon felvezetése („balloon unrossable” léziók [az esetek 6-8%-ában, sikeres distalis drótozást követően]) (48, 49) és a lézióknak megfelelő expanziója („balloon undilatable” léziók [esetek 12%-a sikeres distalis drótozást követően]) (50) sikertelen, amely megakadályozhatja a rekanalizált érszakasz stentelését. A ballonnal átjárhatatlan vagy tágíthatatlan léziók megoldása gyakran speciális technikákat igényelnek – mint például „grenadoplasztika” (azaz kicsiny ballon szándékosan előidézett ruptúrája a proximális sapka szintjében), lézerkatóter, rotációs arterectomia, vagy különböző „cutting” ballonok (spirális vagy paralel vágópengékkel felszerelt ballonok). Ezen eszközök, eljárások helyes algoritmus szerinti alkalmazása az esetek nagy részében lehetővé teszi az intervenciók biztonságos és hatékony befejezését (29).

A speciális CTO-vezetődrótok cseréje fontos lépés a ballon és a stent levezetését megelőzően, ezzel minimálisra csökkentjük a vezetődrót-okozta traumatikus distalis érsérülés lehetőségét. Az angioplasztika az esetek nagy többségében ezután konvencionális módon kivitelezhető, azonban anterográd rekanalizációk esetében a nagyobb és körkörös plakktömeg miatt egyéb plakk-modifikációs technikák használata is szükséges lehet a ballon vagy stentek kihelyezéséhez és megfelelő expanziójához.

Következtetések

Anterográd CTO-rekanalizációs technikák széles körben és hatékonyan alkalmazhatóak a mai CTO-intervenciók gyakorlatban, így ezen szemlélet és gyakorlat megszerzése és alkalmazása a CTO-tréning egyik legfontosabb lépése, továbbá alapul szolgál a komplex retrográd és anterográd disszekciós re-entry technikák elsajátításához is.

Irodalom

1. Fefer P, Knudtson ML, Cheema AN, et al. Current perspectives on coronary chronic total occlusions: the Canadian Multicenter Chronic Total Occlusions Registry. *J Am Coll Cardiol* 2012; 59: 991–7. doi: 10.1016/j.jacc.2011.12.007
2. Jeroudi OM, Alomar ME, Michael TT, et al. Prevalence and ma-

nagement of coronary chronic total occlusions in a tertiary Veterans Affairs hospital. *Catheter Cardiovasc Interv* 2014; 84: 637–43. doi: 10.1002/ccd.25264

3. Christofferson RD, Lehmann KG, Martin GV, Every N, Caldwell JH, Kapadia SR. Effect of chronic total coronary occlusion on treatment strategy. *Am J Cardiol* 2005; 95: 1088–91. doi: 10.1016/j.amjcard.2004.12.065
4. Ramunddal T, Hoebers LP, Henriques JP, et al. Chronic total occlusions in Sweden – a report from the Swedish Coronary Angiography and Angioplasty Registry (SCAAR). *PLoS One* 2014; 9: e103850. doi: 10.1371/journal.pone.0103850
5. Azzalini L, Jolicœur EM, Pighi M, et al. Epidemiology, Management Strategies, and Outcomes of Patients With Chronic Total Coronary Occlusion. *Am J Cardiol* 2016; 118: 1128–1135. doi: 10.1016/j.amjcard.2016.07.023
6. Tajti P, Karpalotis D, Alaswad K, et al. The Hybrid Approach to Chronic Total Occlusion Percutaneous Coronary Intervention: Update From the PROGRESS CTO Registry. *JACC Cardiovasc Interv* 2018. doi: 10.1016/j.jcin.2018.02.036
7. Wilson WM, Walsh SJ, Yan AT, et al. Hybrid approach improves success of chronic total occlusion angioplasty. *Heart* 2016; 102: 1486–93. doi: 10.1136/heartjnl-2015-308891
8. Maeremans J, Walsh S, Knaepen P, et al. The Hybrid Algorithm for Treating Chronic Total Occlusions in Europe: The RECHARGE Registry. *J Am Coll Cardiol* 2016; 68: 1958–1970. doi: 10.1016/j.jacc.2016.08.034
9. Habara M, Tsuchikane E, Muramatsu T, et al. Comparison of percutaneous coronary intervention for chronic total occlusion outcome according to operator experience from the Japanese retrograde summit registry. *Catheter Cardiovasc Interv* 2016; 87: 1027–35. doi: 10.1002/ccd.26354
10. Werner G. A Randomized Multicentre Trial to Evaluate the Utilization of Revascularization or Optimal Medical Therapy for the Treatment of Chronic Total Coronary Occlusions (EuroCTO). *EuroPCR* 2017. Paris, 2017. doi: 10.1093/eurheartj/ehy220
11. Sapontis J, Salisbury AC, Yeh RW, et al. Early Procedural and Health Status Outcomes After Chronic Total Occlusion Angioplasty: A Report From the OPEN-CTO Registry (Outcomes, Patient Health Status, and Efficiency in Chronic Total Occlusion Hybrid Procedures). *JACC Cardiovasc Interv* 2017; 10: 1523–1534. doi: 10.1016/j.jcin.2017.05.065
12. Hannan EL, Zhong Y, Jacobs AK, et al. Patients With Chronic Total Occlusions Undergoing Percutaneous Coronary Interventions: Characteristics, Success, and Outcomes. *Circ Cardiovasc Interv* 2016; 9: e003586. doi: 10.1161/CIRCINTERVENTIONS.116.003586
13. Brilakis ES, Banerjee S, Karpalotis D, et al. Procedural Outcomes of Chronic Total Occlusion Percutaneous Coronary Intervention: A Report From the NCDR (National Cardiovascular Data Registry). *JACC Cardiovasc Interv* 2015; 8: 245–53. doi: 10.1016/j.jcin.2014.08.014
14. Tajti P, Burke MN, Karpalotis D, et al. Update in the Percutaneous Management of Coronary Chronic Total Occlusions. *JACC Cardiovasc Interv* 2018; 11: 615–625. doi: 10.1016/j.jcin.2017.10.052
15. Morino Y, Abe M, Morimoto T, et al. Predicting successful guidewire crossing through chronic total occlusion of native coronary lesions within 30 minutes: the J-CTO (Multicenter CTO Registry in Japan) score as a difficulty grading and time assessment tool. *JACC Cardiovasc Interv* 2011; 4: 213–21. doi: 10.1016/j.jcin.2010.09.024
16. Alessandrino G, Chevalier B, Lefevre T, et al. A Clinical and Angiographic Scoring System to Predict the Probability of Successful First-Attempt Percutaneous Coronary Intervention in Patients With Total Chronic Coronary Occlusion. *JACC Cardiovasc Interv* 2015; 8: 1540–8. doi: 10.1016/j.jcin.2015.07.009
17. Christopoulos G, Kandzari DE, Yeh RW, et al. Development and Validation of a Novel Scoring System for Predicting Technical Success of Chronic Total Occlusion Percutaneous Coronary Interventions: The PROGRESS CTO (Prospective Global Registry for the Study of Chronic Total Occlusion Intervention) Score. *JACC Cardiovasc Interv* 2016; 9: 1–9. doi: 10.1016/j.jcin.2015.09.022

18. Galassi AR, Boukhris M, Azzarelli S, Castaing M, Marza F, Tomasello SD. Percutaneous Coronary Revascularization for Chronic Total Occlusions: A Novel Predictive Score of Technical Failure Using Advanced Technologies. *JACC Cardiovasc Interv* 2016. doi: 10.1016/j.jcin.2016.01.036
19. Maeremans J, Spratt JC, Knaapen P, et al. Towards a contemporary, comprehensive scoring system for determining technical outcomes of hybrid percutaneous chronic total occlusion treatment: The RECHARGE score. *Catheter Cardiovasc Interv* 2017. doi: 10.1002/ccd.27092
20. Ellis SG, Burke MN, Murad MB, et al. Predictors of Successful Hybrid-Approach Chronic Total Coronary Artery Occlusion Stenting: An Improved Model With Novel Correlates. *JACC Cardiovasc Interv* 2017; 10: 1089–1098. doi: 10.1016/j.jcin.2017.03.016
21. Danek BA, Karatasakis A, Karpaliotis D, et al. Development and Validation of a Scoring System for Predicting Periprocedural Complications During Percutaneous Coronary Interventions of Chronic Total Occlusions: The Prospective Global Registry for the Study of Chronic Total Occlusion Intervention (PROGRESS CTO) Complications Score. *J Am Heart Assoc* 2016; 5. doi: 10.1161/JAHA.116.004272
22. Tanaka H, Morino Y, Abe M, et al. Impact of J-CTO score on procedural outcome and target lesion revascularisation after percutaneous coronary intervention for chronic total occlusion: a substudy of the J-CTO Registry (Multicentre CTO Registry in Japan). *EuroIntervention* 2016; 11: 981–8. doi: 10.4244/EIJV11I9A202
23. Christopoulos G, Wyman RM, Alaswad K, et al. Clinical Utility of the Japan-Chronic Total Occlusion Score in Coronary Chronic Total Occlusion Interventions: Results from a Multicenter Registry. *Circ Cardiovasc Interv* 2015; 8: e002171. doi: 10.1161/CIRCINTERVENTIONS.114.002171
24. Nombela-Franco L, Urena M, Jerez-Valero M, et al. Validation of the J-chronic total occlusion score for chronic total occlusion percutaneous coronary intervention in an independent contemporary cohort. *Circ Cardiovasc Interv* 2013; 6: 635–43. doi: 10.1161/CIRCINTERVENTIONS.113.000447
25. Azzalini L, Brilakis ES. Ipsilateral vs. contralateral vs. no collateral (antegrade only) chronic total occlusion percutaneous coronary interventions: What is the right choice for your practice? *Catheter Cardiovasc Interv* 2017; 89: 656–657. doi: 10.1002/ccd.27005
26. Karatasakis A, Danek BA, Karpaliotis D, et al. Comparison of various scores for predicting success of chronic total occlusion percutaneous coronary intervention. *Int J Cardiol* 2016; 224: 50–56. doi: 10.1016/j.ijcard.2016.08.317
27. Brilakis ES, Grantham JA, Rinfret S, et al. A percutaneous treatment algorithm for crossing coronary chronic total occlusions. *JACC Cardiovasc Interv* 2012; 5: 367–79. doi: 10.1016/j.jcin.2012.02.006
28. Harding SA, Wu EB, Lo S, et al. A New Algorithm for Crossing Chronic Total Occlusions From the Asia Pacific Chronic Total Occlusion Club. *JACC Cardiovasc Interv* 2017; 10: 2135–2143. doi: 10.1016/j.jcin.2017.06.071
29. Brilakis ES. *Manual of Coronary Chronic Total Occlusion Interventions: A Step-by-Step Approach*. 2nd ed. 2017.
30. Kinnaird T, Anderson R, Ossei-Gerning N, et al. Vascular Access Site and Outcomes Among 26,807 Chronic Total Coronary Occlusion Angioplasty Cases From the British Cardiovascular Interventions Society National Database. *JACC Cardiovasc Interv* 2017; 10: 635–644. doi: 10.1016/j.jcin.2016.11.055
31. Marquis-Gravel G, Tremblay-Gravel M, Levesque J, et al. Ultra-sound guidance versus anatomical landmark approach for femoral artery access in coronary angiography: A randomized controlled trial and a meta-analysis. *J Interv Cardiol* 2018; 31: 496–503. doi: 10.1111/joic.12492
32. Fairley SL, Lucking AJ, McEntegart M, et al. Routine Use of Fluoroscopic-Guided Femoral Arterial Puncture to Minimise Vascular Complication Rates in CTO Intervention: Multi-centre UK Experience. *Heart Lung Circ* 2016; 25: 1203–1209. doi: 10.1016/j.hlc.2016.04.006
33. Rinfret S, Joyal D, Nguyen CM, et al. Retrograde recanalization of chronic total occlusions from the transradial approach: early Canadian experience. *Catheter Cardiovasc Interv* 2011; 78: 366–74. doi: 10.1002/ccd.23140
34. Takahashi S, Saito S, Tanaka S, et al. New method to increase a backup support of a 6 French guiding coronary catheter. *Catheter Cardiovasc Interv* 2004; 63: 452–6. doi: 10.1002/ccd.20223
35. Finn MT, Green P, Nicholson W, et al. Mother-Daughter-Granddaughter Double GuideLiner Technique for Delivering Stents Past Multiple Extreme Angulations. *Circ Cardiovasc Interv* 2016; 9. doi: 10.1161/CIRCINTERVENTIONS.116.003961
36. Fujita S, Tamai H, Kyo E, et al. New technique for superior guiding catheter support during advancement of a balloon in coronary angioplasty: the anchor technique. *Catheter Cardiovasc Interv* 2003; 59: 482–8. doi: 10.1002/ccd.10551
37. Fang HY, Wu CC, Wu CJ. Successful transradial antegrade coronary intervention of a rare right coronary artery high anterior downward takeoff anomalous chronic total occlusion by double-anchoring technique and retrograde guidance. *Int Heart J* 2009; 50: 531–8. doi: 10.1536/ihj.50.531
38. Mahmood A, Banerjee S, Brilakis ES. Applications of the distal anchoring technique in coronary and peripheral interventions. *J Invasive Cardiol* 2011; 23: 291–4.
39. Michael TT, Banerjee S, Brilakis ES. Subintimal distal anchor technique for „balloon-uncrossable” chronic total occlusions. *J Invasive Cardiol* 2013; 25: 552–4.
40. Danek BA, Karatasakis A, Karpaliotis D, et al. Use of antegrade dissection re-entry in coronary chronic total occlusion percutaneous coronary intervention in a contemporary multicenter registry. *Int J Cardiol* 2016; 214: 428–437. doi: 10.1016/j.ijcard.2016.03.215
41. Valenti R, Vergara R, Migliorini A, et al. Predictors of reocclusion after successful drug-eluting stent-supported percutaneous coronary intervention of chronic total occlusion. *J Am Coll Cardiol* 2013; 61: 545–50. doi: 10.1016/j.jacc.2012.10.036
42. Azzalini L, Dautov R, Brilakis ES, et al. Procedural and longer-term outcomes of wire- versus device-based antegrade dissection and re-entry techniques for the percutaneous revascularization of coronary chronic total occlusions. *Int J Cardiol* 2017; 231: 78–83. doi: 10.1016/j.ijcard.2016.11.273
43. Rathore S, Matsuo H, Terashima M, et al. Procedural and in-hospital outcomes after percutaneous coronary intervention for chronic total occlusions of coronary arteries 2002 to 2008: impact of novel guidewire techniques. *JACC Cardiovasc Interv* 2009; 2: 489–97. doi: 10.1016/j.jcin.2009.04.008
44. Chiu CA. Recanalization of difficult bifurcation lesions using adjunctive double-lumen microcatheter support: two case reports. *J Invasive Cardiol* 2010; 22: E99–103.
45. Saito S. Open Sesame Technique for chronic total occlusion. *Catheter Cardiovasc Interv* 2010; 75: 690–4. doi: 10.1002/ccd.22316
46. Roy J, Hill J, Spratt JC. The “side-BASE technique”: Combined side branch anchor balloon and balloon assisted sub-intimal entry to resolve ambiguous proximal cap chronic total occlusions. *Catheter Cardiovasc Interv* 2017. doi: 10.1002/ccd.27422
47. Carlino M, Latib A, Godino C, Cosgrave J, Colombo A. CTO recanalization by intraocclusion injection of contrast: the micro-channel technique. *Catheter Cardiovasc Interv* 2008; 71: 20–6. doi: 10.1002/ccd.21396
48. Patel SM, Pokala NR, Menon RV, et al. Prevalence and treatment of “balloon-uncrossable” coronary chronic total occlusions. *J Invasive Cardiol* 2015; 27: 78–84.
49. Karacsonyi J, Karpaliotis D, Alaswad K, et al. Prevalence, indications and management of balloon uncrossable chronic total occlusions: Insights from a contemporary multicenter US registry. *Catheter Cardiovasc Interv* 2016. doi: 10.1002/ccd.26780
50. Tajti P, Karpaliotis D, Alaswad K, et al. Prevalence, Presentation and Treatment of “Balloon Undilatable” Chronic Total Occlusions: Insights from a Multicenter US Registry. *Catheter Cardiovasc Interv* 2018; 91: 657–666. doi: 10.1002/ccd.27510

III.

Kettős lumenű mikrokatóterek alkalmazása krónikus teljes koronária-okklúzió katéteres revaszkularizációjában – Egy komplex beavatkozás tanulságai

Tajti Péter¹, Rami Abu Fanne², Ungi Imre¹, Katona András¹, Sasi Viktor¹, Nagy Ferenc Tamás¹

¹Szegedi Tudományegyetem, Invazív Kardiológiai Részleg, II. sz. Belgyógyászati Klinika és Kardiológiai Központ, Szeged

²Hillel Yaffe Medical Center, Hadera, Izrael

Levelezési cím:

Dr. Tajti Péter

Szegedi Tudományegyetem, Invazív Kardiológiai Részleg, II. sz. Belgyógyászati Klinika és Kardiológiai Központ
6725 Szeged, Semmelweis u. 8.

E-mail: ptajti@gmail.com

A krónikus, teljes koszorúér-okklúzió (CTO) perkután intervenció (PCI) sikeres és biztonságos kivitelezhetősége jelentősen javult az arra felkészült centrumokban az elmúlt években. Ugyanakkor az anamnézisben szereplő aorto-koronáriás bypass graft (ACBG) műtét jelentős mértékben csökkenti a CTO PCI sikerességét miközben emeli a beavatkozáshoz társuló szövődmények gyakoriságát. A komplex koronária anatómiájából adódó nehézségek megoldása gyakran előrehaladott technikák ismeretét és eszközök alkalmazását követeli meg az effektív és biztonságos intervenciók kivitelezéséhez ezen speciális betegpopulációban. Jelen esetbemutatásunkban egy korábban két alkalommal sikertelen anterográd CTO-rekanalizáción átesett poszt-ACBG-beteg komplex retrográd intervencióját ismertetjük. A beavatkozás során a komplex jobb koronária-bifurkáció és a korábban erre varrott, majd elzáródott saphena véna graft helyzete miatt egyedülálló módon retrográd és anterográd kettős lumenű mikrokatóterrel facilitált rekanalizációs technikát alkalmaztunk, amely eredményeképpen sikeres CTO-megnyitást végeztünk.

Kulcsszavak: kettős lumenű mikrokatóter, aorto-koronáriás bypass graft műtét, krónikus teljes okklúzió, perkután koronária-intervenció

Application of dual lumen microcatheters in chronic total occlusion percutaneous coronary interventions – lessons learned from a complex case

The success rate and safety of percutaneous coronary intervention (PCI) of chronic total coronary occlusions (CTO) in dedicated CTO centers has significantly improved recently. History of prior coronary artery bypass graft surgery (CABG) in patients undergoing CTO PCI is associated with lower procedural success and higher rates of major procedural complications. Increased complexity of coronary anatomy requires the use of advanced interventional techniques and devices in order to achieve successful CTO revascularization safely and effectively in such a challenging patient population. In our current case we present a complex right coronary CTO recanalization in a patient with a history of CABG surgery who had undergone two previous failed antegrade CTO attempts in an outside hospital. Given the complexity of the distal right coronary artery bifurcation and pre-branch saphenous vein graft (SVG) anastomosis we used the dual lumen microcatheter in a unique way both retrogradely and antegradely to preserve side-branches and achieve success.

Keywords: dual lumen microcatheter, coronary artery bypass graft operation, chronic total occlusion, percutaneous coronary intervention

Bevezetés

Korszerű eszközök, újabb technikák és algoritmusok megjelenése a krónikus teljes koronária-okklúziók (CTO) perkután koronária-intervenciójában (PCI) a sikerráta jelentős növekedését eredményezte (85-90%) tapasztalt operatőrök kezében (1–3). Ugyanakkor a korábban aorto-koronáriás bypass graft (ACBG) műtéten áteset betegek esetében továbbra is nagy technikai kihívást jelent a krónikus okklúziók revaszkularizációja. Ennek hátterében elsősorban többszörös társbetegségek jelenléte, illetve anatómiai sajátosságok állnak (anasztomózis lokalizációja, disztális bifurkáció potenciális jelenléte, meszesedés foka, a perikardiális hegesedés miatti epikardiális koronáriák merevsége, koronária-tortuozitás, „tenting” jelenség [zsugorodott SVG által felhúzott érlefutás mindig a graftba vezet mind az antero-, mind a retrográd eszközöket és néha lehetetlenné teszi az éles kanyar bevételét a „sátor” tetején]) (4, 5). Jelen esetünkön keresztül ismertetjük a kettős lumenű mikrokatóter (a rapid exchange [RX] disztális port és proximális over-the-wire [OTW] oldalsó port struktúra szimultán 2 vezetődrót bevezetését teszi lehetővé) hasznosságát anterográd és retrográd drótozás esetében, amely komplex anatómiai helyzetekben nyújthat effektív megoldást speciális CTO-intervenciókban (poszt-ACBG, bifurkáció).

Esetismertetés

A 75 éves férfi beteg kórelőzményében hipertenzió, dyslipidaemia, inzulinval kezelte 2-es típusú diabetes mellitus szerepelt. Revaszkularizációs anamnéziséből kiemelendő 1999 RCA (right coronary artery) PCI (bare metal stent, BMS), 2000-ben RCA stentokklúzió és reziduális szűkületek miatti ACBG-műtét (LIMA-LAD, SVG-CXOM, SVG-RCA, SVG-I. diagonális ág), valamint 2016-ban az RCA véna graft okklúziója miatti graft PCI (drug eluting stent, DES). Rekurráló panaszok miatt 2018-ban elvégzett koronarográfia során okkludált véna graftok igazolódtak (SVG-RCA, SVG-diagonális ág, SVG-CXOM), amely miatt első lépésben CX PCI történt (DES). Kardiális mágneses rezonancia (cMR) által igazolt hátsófalvi viabilitás igazolása után és optimális gyógyszeres terápia beállítása ellenére perzisztáló panaszok (CCS 2-3 angina ekvivalens fulladás) miatt hazánk egy másik centrumában két alkalommal is sikertelen anterográd RCA-rekanalizációs kísérletet végeztek. Ezt követően vettük fel a betegünket ismételt jobb koronária-revaszkularizációs kísérlet elvégzése céljából.

Bal arteria radialis és jobb arteria femoralis punkcióját követően 7 French EBU 4.0 valamint 7 French AL 1.0 katéterekkel kanuláltuk a bal közös törzset, illetve a jobb koronária-szájadékot. Bilaterális injekció alapján (1. ábra A panel) aneurizmatikusan tágult proximális

1. ÁBRA. Kettős lumenű mikrokatóter retrográd alkalmazása krónikus teljes koronária-okklúzió (CTO) perkután intervenciója (PCI) során. **A:** Bilaterális injekció reprezentálja a crux-ig terjedő disztális jobb koronária krónikus okklúziójának disztális és proximális anatómiáját. **B:** Retrográd mikrokatóter (Corsair Pro [Asahi Intecc. Nagoya, Japan]) tip injekciója. **C:** Sikeres disztális CTO-sapkához történő drótozás (Sion [Asahi Intecc. Nagoya, Japan], Corsair Pro). **D:** Saphena graftba parkolt Confianza Pro 12 vezetődrót [Asahi Intecc. Nagoya, Japán]. **E:** Retrográd kettős lumenű mikrokatóter (Sasuke [Asahi Intecc. Nagoya, Japán]) alkalmazása a disztális sapka over-the-wire (OTW) porton keresztül történő punkciójához. **F:** Anterográd és retrográd „nukle” drótok (Fielder XT, Pilot 200 [Asahi Intecc. Nagoya, Japan]) koaxiális pozícióban. **G:** Sikertelen reverz CART (controlled antegrade and retrograde subintimal tracking) retrográd disszekció és re-entry technika a vertikális jobb koronáriaszakaszon. **H:** GuideLiner (Vascular Solutions, Minneapolis, MN, USA) reverz CART technika vezérelte vezetődrót belepés az anterográd vezetőkatéterbe

RCA, tompa okklúziós sapka, valamint kielégítő disztális retrográd telődés (Rentrop 2-3, dominánsan szeptális kollaterálisokon keresztül) volt detektálható. Kezdő stratégiaként anterográd drót eszkalációt kíséreltünk meg Gaia 2 és Fielder XT (mindkettő Asahi Intecc, Nagoya, Japán) vezetődrótok és Corsair Pro (Asahi Intecc, Nagoya, Japán) mikrokatóter szupportjával. Rövid próbálkozások után azonban szubintimális drótpozíció volt sejtethető, így „knuckle” drót segítségével tompa preparálással jutottunk a disztális jobb koronáriszakaszba, majd egy rendkívül kemény, áthatolhatatlan szegmentum miatt retrográd technikára tértünk át. Kontrasztanyag-vezérelt drótozás révén (mikrokatóter tip injekciók [1. ábra B panel]) sikeresen passzáltuk a szeptális kollaterális csatornát (Sion [Asahi Intecc, Nagoya, Japán] Corsair Pro). Többszöri próbálkozás ellenére sem sikerült azonban a disztális sapkát áttörni (1. ábra C panel) (Fielder XT, Gaia 3, Sion), mivel a már említett „tenting” jelenség miatt ismételt a disztális jobb koronária-bifurkáció előtt anasztomizált okkludált véna graftba jutottunk (Confianza Pro 12 [Asahi Intecc, Nagoya, Japán]). A komplex anatómia megoldásaként a véna graftba parkolt vezetődrót (1. ábra D panel) felhasználva Sasuke kettős lumenű mikrokatótert juttattunk (Asahi Intecc,

Nagoya, Japán) abba, majd annak oldal-portján (OTW port) keresztül megfelelő szupportot és irányt tudtunk biztosítani a második retrográd drót sikeres disztális sapka punkciójához (1. ábra E panel) (Miracle 6 [Asahi Intecc, Nagoya, Japán]). Következő lépésben a Sasuke mikrokatótert Corsair Pro-ra cserélve retrográd Pilot 200 (Asahi Intecc, Nagoya, Japán) „knuckle” drót segítségével megfelelő koaxiális drótpozíciót vettünk fel az anterográd drót magasságában (1. ábra F panel). GuideLinerrel (Vascular Solutions, Minneapolis, MN, USA) asszisztált reverz CART (controlled antegrade and retrograde subintimal tracking) technikával sikeresen az anterográd vezető katéterbe jutottunk (1. ábra G és H panel). Ezt drót externalizáció követte, azaz 330 cm-es RG3 (Asahi Intecc, Nagoya, Japán) drótot juttatunk a retrográd vezető katéter felől a mikrokatóter lumenen át az anterográd vezető katéterbe, majd ezt anterográd vezetődrótként használva ballon predilatációt végeztünk az okkludált szegmentumban. Kontralaterális angiogramok és szimultán disztális RCA-ballon infláció útján győződünk meg arról, hogy a cruxban retrográd disszekció alakult ki. Ezzel a vizualizációs technikával elkerülhettük az anterográd disszekciós tasak hidraulikus expanszióját. Tekintettel a disszekcióra és az oldalág megőrzés

2. ÁBRA. Kettős lumenű mikrokatóter anterográd alkalmazása krónikus teljes koronária-okklúzió (CTO) perkután intervenciója (PCI) során. **A:** Kissing-ballon tágítás a poszterolaterális oldalág kettős lumenű Sasuke (Asahi Intecc, Nagoya, Japan) való újradrótózást követően. **B:** GuideLiner (Vascular Solutions, Minneapolis, MN, USA) facilitált angiográfia, amely perzisztáló crux disszekciót mutat. **C:** Sasuke asszisztált poszterior descendens ág drótozás majd externalizált dróteltávolítás. **D:** „Crossover” stentelés a poszterolaterális ág fele. **E:** Proximális stent-implantáció reziduális proximális disszekció miatt. **F:** Sikeres jobb koronária CTO-intervenció 3 gyógyszerkibocsájtó stent implantációjával

fontosságára, a poszterolaterális (PL) ág drótozásához ismételten a Sasuke kettős lumenű mikrokatótert használtuk, majd kissing dilatációkat végeztünk a crux bifurkációjában (2. ábra A panel). A perzisztáló jobb koronária crux disszekció miatt (2. ábra B panel) ismételten a kettős lumenű mikrokatótert alkalmaztuk – ezúttal az anterográd PL-drótra vezetve – és az OTW-lumenen át újra drótozást végeztünk a poszterior descendens (PD) ágba (2. ábra C panel). Ezt követően a retrográd externalizációs drótot már biztonsággal eltávolíthattuk, hiszen mindkét ágban biztosan intraluminális anterográd drót állt rendelkezésünkre. A PL-ág felé „crossover” stentelést végeztünk (2. ábra D panel), majd a Sasuke mikrokatóter segítségével a stent cellákon át a PD-ágot ismételten drótoztuk. Utolsó lépésben kissing utótágítást végeztünk és két addicionális gyógyszerkibocsátó stentet helyeztünk be (2. ábra E panel), jó angiográfiás végeredménnyel (2. ábra F panel) (kontrasztanyag 320 ml, procedurális idő 257 min, sugár dózis 2,114 Gray). A beteget, 48 órás eseménytelen obszervációját követően, stabil és panaszmentes állapotban otthonába bocsájtottuk.

Megbeszélés

Irodalmi ismereteink szerint esetünk egyedülállóan mutatja be retrográd kettős lumenű mikrokatóter többszörös alkalmazását korábban ACBG-műtéten átesett beteg CTO rekanalizációja céljából (3. ábra). Ennek során 1. az eszközt retrográd úton szeptális kollaterálison át a disztális főágba hatolásra;

2. anterográd irányban a disszekált PL-ág lumenének elérésére;
3. anterográd irányban a PD-ág lumenének elérésére;
4. crossover stenten át a PD-ág újradrótozására használtuk.

Esetünk azt példázza, hogy komplex technikák alkalmazása fontos sarokpontja a krónikus okklúziók megnyitásában az oldalágvesztés elkerülésének, mely meghatározó lehet a hosszú távú poszt-PCI-mortalitás csökkentésében.

Számos tanulmány vizsgálta az ACBG-műtött betegekben végzett CTO-rekanalizációk eredményeit rövid és hosszú távon. *Christopoulos és munkatársai* 496 betegben vizsgálták a hibrid algoritmus alkalmazását CTO PCI-ben post-ACBG-betegekben (n=176, 35%), ACBG nélküli csoporttal összehasonlítva (n=320) (4). Az ACBG-betegekben gyakrabban volt szükség retrográd technikák alkalmazására (39% vs. 24%, p<0,001), valamint az intervenciós sikerarány (88% vs. 93%, p=0,04) csökkenő tendenciát mutatott, megfelelően reprezentálva ezen klinikai esetek komplexitását. Ugyanakkor a beavatkozást követően nem mutatkozott számottevő különbség a szövődményeket illetően (major adverse cardiac event, MACE; 2,1% vs. 1,1%, p=0,40). *Azzalini és munkatársai* hasonló tendenciájú klinikai sikerarányt találtak 2058 esetszámú (ACBG [n=401] 82% vs. non-ACBG [n=1657] 88%, p=0,001) heterogén betegpopulációban (multicentrikus vizsgálat) (5). Ezen tanulmány alapján – a korábbiakkal ellentétben – szignifikánsan magasabb periprocedurális komplikáció ráta (MACE, 3,7% vs. 1,5%, p=0,0004), és medián 377 napos utánkövetési idővel magasabb kar-

3. ÁBRA. Retrográd es anterográd kettős lumenű mikrokatóter facilitált jobb koronária krónikus teljes okklúzió perkután intervenciójának vázlatos lépései (CART: controlled antegrade and retrograde subintimal tracking; MC: mikrokatóter; PD: posterior descendens; PL: poszterolaterális; RCA: right coronary artery; SVG: saphena véna graft)

4. ÁBRA. A kettős lumenű mikrokatóter felhasználási lehetőségei bifurkáció drótozás esetén („over-the-wire” port [OTW, sárga], „rapid-exchange” port [RX, kék]). **A:** Anterográd bifurkáció-drótozás. **B:** Anterográd bifurkáció (újra) drótozás „stent struktok”-on keresztül. **C:** Retrográd parallel drótozás szubintimálisan parkolt első retrográd dróton (RX) felvezetett kettős lumenű mikrokatóter „over the wire” (OTW) portját felhasználva. **D:** Retrográd kettős lumenű mikrokatóter felhasználása SVG-graftban parkolt RX-drót, és OTW-oldalport drót segítségével. **E:** Retrográd poszterior descendens (PD) poszterolaterális (PL) drótozás RX-porton, majd OTW-porton keresztüli disztális sapkapunkció. **F:** Disztális re-entry-t követően a valódi lumenben parkolt RX-drót, és reverz OTW-drót segítségével történő bifurkációs oldalágdrótozás anterográd úton

diális mortalitás (3,8% vs. 1,9%, $p=0,02$), „target” lézió re-intervenció (11,5% vs. 6,6%, $p<0,001$), és „target” lézió területi miokardiális infarktuszban (2,0% vs. 0,7%, $p=0,04$) mutatkozott az ACBG-csoportban. Mindemellett az ACBG-műtöttek CTO-intervenciójához felhasznált kontrasztanyag-mennyiség, sugárdózis és procedurális idő is szignifikánsan meghaladja a non-ACBG CTO-intervenciókét (3, 4). Ezen legfrissebb tanulmányok alátámasztják, hogy előzetes ACBG-műtét után elzáródott SVG-ok mellett a CTO perkután intervenciója rendkívül komplex beavatkozás. Európai statisztikai adatok alapján ritkábban találkozunk ACBG utáni CTO-val (15-22%) (1, 2, 6) mint az Egyesült Államokban (28-37%) (3, 4).

A jelentős oldalágak megőrzése valamint a periprocedurális infarktus elkerülése CTO PCI során kritikus fontosságú a betegek hosszú távú mortalitásának csökkentésében (7, 8). Az okklúzió szegmentumában eredő nagy oldalág (okklúziótól proximálisan, az okklúzió magasságában, valamint okklúziótól disztálisan) elfordulási incidenciája 26,5–33,0% közöttre tehető a perkután intervenciók kísérleten áteső betegek körében (9). Tekintettel arra, hogy minden 3. esetben a bifurkáció is érintett speciális „oldalág-mentési” technikákra lehet szükség. A proximális oldalág elsősorban horgonyzási technikák alkalmazásában lehet releváns, valamint preventív drótozás válhat szükségessé az anterográd drót eszkaláció megkezdése előtt, ugyanakkor oldalágvesztés lehetőségével ritkán, csak retrográd, súlyosan

propagálódó disszekció esetén érdemes számolnunk. Ezzel szemben az okklúzió testében, valamint az elzáródás disztális sapkájánál eredő oldalág megőrzése jelentős mértékben csökkenti a beavatkozás klinikai sikerarányát (3).

Bemutatott esetünk jól szemlélteti az ACBG és bifurkációk együttese által okozott technikai akadályokat CTO PCI során, amelyek azonban a megfelelő eszközök használatával áthidalhatók. A retrográd mikrokatóterek használata jelentős mértékben függ a kollaterális méretétől és annak tortuozitásától (szeptális, epikardiális, vagy saphena véna graft), ezért a megfelelő mikrokatóter választásában a tip profiljának és átmerőjének fontos szerepe van. Habár több kettős lumenű mikrokatóter érhető el világszerte (Twin-Pass, Twin-Pass Torque [mindkettő Teleflex, Wayne, PA, USA], Crusade [Kaneka, Tokió, Japán], Fine Duo [Terumo, Tokió, Japán], NHancer [IMDS, Roden, Hollandia]), az általunk hazánkban elsőként alkalmazott Sasuke az egyedüli biztonságosan alkalmazható alacsony tip-profilú katéter, amely a szeptális kollaterálisok többségén is biztonságosan átvezethető (tip átmerő 1,5 Fr vs. 1,8 Fr [Fine Duo] vs. 2,0 Fr [Twin-Pass] vs. 2,1 Fr [Twin-Pass Torque] vs. 2,2 Fr [Crusade], 2,3 [NHancer]). Epikardiális kollaterálisokon keresztüli applikációja jelenleg nem került leírásra, és ennek magasabb perforáció rizikó miatt a használatát jelenleg nem javasoljuk. Ezen speciális esetünkben egyedülállóan „kihorgonyoztuk” a retrográd mikrokatótert az okkludált SVG disztális sza-

kaszán, amely kellő szupportot adott a kemény disztális sapka második dróttal történő penetrációjához (4. ábra D panel). Mindemellett a retrográd kettős lumenű mikrokatóter alkalmazható sikeres PL-PD, vagy PD-PL-ág drótozásakor (véna graft involválása nélkül) a disztális sapka punkciójához (4. ábra E panel). A kettős lumenű mikrokatóter alkalmazása retrográd parallel drótozásra is használható, mely szubintimális retrográd drótpozíció mellett az OTW-port használatával a második vezetődrót révén az intraluminális pozíció visszanyerhető (10) (4. ábra C panel). Ugyanakkor kiemelendő, hogy a kettős lumenű mikrokatóterek alkalmazása elsősorban disztális tip profiljuk miatt dominánsan anterográd drótozáshoz használatosak (4. ábra A panel). Főbb alkalmazási indikációs köréhez tartozik továbbá a szubintimális drótpozíciót követően egy második vezetődróttal történő „reentry” a disztális valódi lumenbe, valamint okklúziós test, illetve disztális sapkabifurkáció esetén anterográd drótozáshoz (4. ábra F panel). Két stentes bifurkációs technika alkalmazásakor a kettős lumenű mikrokatóter szinten kiválóan alkalmas a stent cellákon keresztüli vagy éles szögben eredő oldalágba bejutás megkönnyítésére (4. ábra B panel).

Következtetések

A kettős lumenű mikrokatóterek hasznos eszközként szolgálhatnak a bifurkációt érintő és/vagy elzáródott graft által megnehezített natív krónikus teljes koronária-okklúzió intervenciókban, megkönnyítve az anterográd és retrográd drótozást az oldalág megörzését.

Irodalom

1. Wilson WM, Walsh SJ, Yan AT, et al. Hybrid approach improves success of chronic total occlusion angioplasty. *Heart* 2016; 102:

1486–93. DOI:10.1136/heartjnl-2015-308891

2. Maeremans J, Walsh S, Knaapen P, et al. The Hybrid Algorithm for Treating Chronic Total Occlusions in Europe: The RECHARGE Registry. *J Am Coll Cardiol* 2016; 68: 1958–1970. DOI: 10.1016/j.jacc.2016.08.034

3. Tajti P, Karpalotis D, Alaswad K, et al. The Hybrid Approach to Chronic Total Occlusion Percutaneous Coronary Intervention: Update From the PROGRESS CTO Registry. *JACC Cardiovasc Interv* 2018. DOI:10.1016/j.jcin.2018.02.036

4. Christopoulos G, Menon RV, Karpalotis D, et al. Application of the "hybrid approach" to chronic total occlusions in patients with previous coronary artery bypass graft surgery (from a Contemporary Multicenter US registry). *Am J Cardiol* 2014; 113: 1990–4. DOI:10.1016/j.amjcard.2014.03.039

5. Azzalini L, Ojeda S, Karatasakis A, et al. Long-Term Outcomes of Percutaneous Coronary Intervention for Chronic Total Occlusion in Patients Who Have Undergone Coronary Artery Bypass Grafting vs Those Who Have Not. *Can J Cardiol* 2018; 34: 310–318. DOI:10.1016/j.cjca.2017.12.016

6. Galassi AR, Sianos G, Werner GS, et al. Retrograde Recanalization of Chronic Total Occlusions in Europe: Procedural, In-Hospital, and Long-Term Outcomes From the Multicenter ERCTO Registry. *J Am Coll Cardiol* 2015; 65: 2388–400. DOI: 10.1016/j.jacc.2015.03.566

7. Jang WJ, Yang JH, Choi SH, et al. Association of periprocedural myocardial infarction with long-term survival in patients treated with coronary revascularization therapy of chronic total occlusion. *Catheter Cardiovasc Interv* 2016; 87: 1042–9. DOI:10.1002/ccd.26286

8. Galassi AR, Boukhris M, Tomasello SD, et al. Incidence, treatment, and in-hospital outcome of bifurcation lesions in patients undergoing percutaneous coronary interventions for chronic total occlusions. *Coron Artery Dis* 2015; 26: 142–9. DOI: 10.1097/MCA.000000000000194

9. Ojeda S, Pan M, Gutierrez A, et al. Bifurcation lesions involved in the recanalization process of coronary chronic total occlusions: Incidence, treatment and clinical implications. *Int J Cardiol* 2017; 230: 432–438. DOI: 10.1016/j.ijcard.2016.12.088

10. Tanabe G, Oikawa Y, Yajima J, Matsuno S, Kano H, Yamashita T. Retrograde parallel wire technique using a dual lumen catheter can be useful for percutaneous coronary intervention with chronic total occlusion. *J Cardiol Cases* 2018; 17: 25–28. DOI: 10.1016/j.jccase.2017.08.014